

## REFERENCES

1. Steindler A. Kinesiology of the Human Body under Normal and Pathological Conditions. *J Am Med Assoc.* 1955;158(17):1573.
2. Bakhtiary a H, Fatemi E. Open versus closed kinetic chain exercises for patellar chondromalacia. *Br J Sports Med.* 2008;42(December):99-102; discussion 102.
3. Harvie D, Oleary T, Kumar S. A systematic review of randomized controlled trials on exercise parameters in the treatment of patellofemoral pain: What works? *J Multidiscip Healthc.* 2011;4:383-392.
4. Dolak KL, Silkman C, Medina McKeon J, Hosey RG, Lattermann C, Uhl TL. Hip strengthening prior to functional exercises reduces pain sooner than quadriceps strengthening in females with patellofemoral pain syndrome: a randomized clinical trial. *J Orthop Sports Phys Ther.* 2011;41(8):560-570.
5. Nakagawa TH, Muniz TB, Baldon RDM, Dias Maciel C, de Menezes Reiff RB, Serrão FV. The effect of additional strengthening of hip abductor and lateral rotator muscles in patellofemoral pain syndrome: a randomized controlled pilot study. *Clin Rehabil.* 2008;22(12):1051-1060.
6. Ismail MM, Gamaleldin MH, Hassa KA. Closed Kinetic Chain exercises with or without additional hip strengthening exercises in management of Patellofemoral pain syndrome: A randomized controlled trial. *Eur J Phys Rehabil Med.* 2013;49(5):687-698.
7. Fukuda TY. Hip Posterolateral Musculature Strengthening in Sedentary Women With Patellofemoral Pain Syndrome: A Randomized Controlled Clinical Trial With 1-Year Follow-up. *J Orthop Sports Phys Ther.* 2012;42(10):823-830.
8. Khayambashi K, Fallah A, Movahedi A, Bagwell J, Powers C. Posterolateral hip muscle strengthening versus quadriceps strengthening for patellofemoral pain: A comparative control trial. *Arch Phys Med Rehabil.* 2014;95(5):900-907.
9. Baldon RDM, Serrão FV, Scattone Silva R, Piva SR. Effects of functional stabilization training on pain, function, and lower extremity biomechanics in women with patellofemoral pain: a randomized clinical trial. *J Orthop Sports Phys Ther.* 2014;44(4):240-A8.
10. Moyano FR, Valenza MC, Martin LM, Caballero YC, Gonzalez-Jimenez E, Demet GV. Effectiveness of different exercises and stretching physiotherapy on pain and movement in patellofemoral pain syndrome: a randomized controlled trial. *Clin Rehabil.* 2013;27(5):409-417.
11. Alba-Martín P, Gallego-Izquierdo T, Plaza-Manzano G, Romero-Franco N, Núñez-Nagy S, Pecos-Martín D. Effectiveness of therapeutic physical exercise in the treatment of patellofemoral pain syndrome: a systematic review. *J Phys Ther Sci.* 2015;27(7):2387-2390.
12. Mason M, Keays SL, Newcombe PA. The Effect of Taping, Quadriceps Strengthening and Stretching Prescribed Separately or Combined on Patellofemoral Pain. *Physiother Res Int.* 2011;16(2):109-119.
13. Peeler J, Anderson JE. Effectiveness of static quadriceps stretching in individuals with patellofemoral joint pain. *Clin J Sport Med.* 2007;17(4):234-241.
14. White LC, Dolphin P, Dixon J. Hamstring length in patellofemoral pain syndrome. *Physiotherapy.* 2009;95(1):24-28.
15. Tyler TF, Nicholas SJ, Mullaney MJ, McHugh MP. The role of hip muscle function in the treatment of patellofemoral pain syndrome. *Am J Sports Med.* 2006;34(4):630-636.
16. Willett GM, Paladino JB, Barr KM., Korta JN., Karst GM. Medial and Lateral Quadriceps Muscle Activity During Weight-Bearing Knee Extension Exercise. *J Sport Rehabil.* 1998;7(4):248.
17. C.M. G, P.H. K, C.G. M. Targeting quadriceps inhibition with electromyographic biofeedback: A neuroplastic approach. *Crit Rev Biomed Eng.* 2013;41(2):125-135.
18. Hopkins JT, Ingersoll CD. Arthrogenic muscle inhibition: A limiting factor in joint rehabilitation. *J Sport Rehabil.* 2000;9(2):135-159.
19. Escamilla RF, Zheng N, Macleod TD, et al. Patellofemoral joint force and stress during the wall squat and one-leg squat. *Med Sci Sports Exerc.* 2009;41(4):879-888.
20. Escamilla RF. Squat Exercise. *Med Sci Sports Exerc.* 2001;33(1):127-141.
21. Escamilla RF, Fleisig GS, Zheng N, et al. Effects of technique variations on knee biomechanics during the squat and leg press. *Med Sci Sports Exerc.* 2001;33(9):1552-1566.
22. Escamilla RF, Zheng N, Macleod TD, et al. Patellofemoral joint force and stress between a short- and long-step forward lunge. *J Orthop Sports Phys Ther.* 2008;38(11):681-690.
23. Fry AC, Smith JC, Schilling BK. Effect of knee position on hip and knee torques during the barbell squat. *J Strength Cond Res.* 2003;17(4):629-633.
24. List R, Gülay T, Stoop M, Lorenzetti S. Kinematics of the trunk and the lower extremities during restricted and unrestricted squats. *J Strength Cond Res.* 2013;27(6):1529-1538.
25. Tang SFT, Chen CK, Hsu R, Chou SW, Hong WH, Lew HL. Vastus medialis obliquus and vastus lateralis activity in open and closed kinetic chain exercises in patients with patellofemoral pain syndrome: An electromyographic study. *Arch Phys Med Rehabil.* 2001;82(10):1441-1445.
26. Escamilla RF. Knee biomechanics of the dynamic squat exercise. *Med Sci Sport Exerc.* 2001;33(1):127-141.
27. Bell DR, Oates DC, Clark MA, Padua DA. Two- and 3-dimensional knee valgus are reduced after an exercise intervention in young adults with demonstrable valgus during squatting. *J Athl Train.* 2013;48(4):442-449.

28. Kulas AS, Hortobagyi T, Devita P. Trunk position modulates anterior cruciate ligament forces and strains during a single-leg squat. *Clin Biomech.* 2012;27(1):16-21.
29. Lynn SK, Noffal GJ. Lower Extremity Biomechanics During a Regular and Counterbalanced Squat. *J Strength Cond Res.* 2012;26(9):2417-2425.
30. Irish SE, Millward AJ, Wride J, Haas BM, Shum GLK. The effect of closed-kinetic chain exercises and open-kinetic chain exercise on the muscle activity of vastus medialis oblique and vastus lateralis. *J Strength Cond Res.* 2010;24(5):1256-1262.
31. McLaughlin J, DeMaio M, Noyes FR, Mangine RE. Rehabilitation after meniscus repair. *Orthopedics.* 1994;17(5):463-471.
32. Cavanaugh JT, Killian SE. Rehabilitation following meniscal repair. *Curr Rev Musculoskelet Med.* 2012;5(1):46-58.
33. Pabian P, Hanney WJ. Functional Rehabilitation after Medial Meniscus Repair in a High School Football Quarterback: A Case Report. *N Am J Sports Phys Ther.* 2008;3(3):161-169.
34. Barber FA. Accelerated rehabilitation for meniscus repairs. *Arthrosc J Arthrosc Relat Surg.* 1994;10(2):206-210.
35. Stiene HA, Brosky T, Reinking MF, Nyland J, Mason MB. A comparison of closed kinetic chain and isokinetic joint isolation exercise in patients with patellofemoral dysfunction. *J Orthop Sports Phys Ther.* 1996;24(3):136-141.
36. Palmitier RA, An KN, Scott SG, Chao EYS. Kinetic Chain Exercise in Knee Rehabilitation. *Sport Med.* 1991;11(6):402-413.
37. Rasch PJ, Morehouse LE. Effect of static and dynamic exercises on muscular strength and hypertrophy. *J Appl Physiol.* 1957;11:29-34.
38. Rutherford OM. Muscular Coordination and Strength Training: Implications for Injury Rehabilitation. *Sport Med.* 1988;5(3):196-202.
39. Witvrouw E, Lysens R, Bellemans J, Peers K, Vanderstraeten G. Open versus closed kinetic chain exercises for patellofemoral pain. A prospective, randomized study. *Am J Sports Med.* 2000;28(5):687-694.
40. Huberti HH, Hayes WC. Patellofemoral contact pressures. The influence of q-angle and tendofemoral contact. *J Bone Joint Surg Am.* 1984;66(5):715-724.
41. Hungerford DS, Barry M. Biomechanics of the patellofemoral joint. *Clin Orthop Relat Res.* 1979;(144):9-15.
42. Steinkamp LA, Dillingham MF, Markel MD, Hill JA, Kaufman KR. Biomechanical considerations in patellofemoral joint rehabilitation. *Am J Sport Med.* 1993;21(3):438-444.
43. Woodall WR, Welsh J. A biomechanical basis for rehabilitation programs involving the patellofemoral joint. *J Orthop Sports Phys Ther.* 1990;11(11):535-542.
44. Schoenfeld BJ. Squatting kinematics and kinetics and their application to exercise performance. *J Strength Cond Res.* 2010;24(12):3497-3506.
45. Escamilla RF. Knee biomechanics of the dynamic squat exercise. *Med Sci Sports Exerc.* 2001;33(1):127-141.
46. Isear J a, Erickson JC, Worrell TW. EMG analysis of lower extremity muscle recruitment patterns during an unloaded squat. *Med Sci Sports Exerc.* 1997;29(4):532-539.
47. Ninos JC, Irrgang JJ, Burdett R, Weiss JR. Electromyographic analysis of the squat performed in self-selected lower extremity neutral rotation and 30o of lower extremity turn-out from the self-selected neutral position. *J Orthop Sport Phys Ther.* 1997;25(5):307-315.
48. Stuart MJ, Meglan D a, Lutz GE, Growney ES, An KN. Comparison of intersegmental tibiofemoral joint forces and muscle activity during various closed kinetic chain exercises. *Am J Sports Med.* 1996;24:792-799.
49. Escamilla RF, Fleisig GS, Zheng N, Barrentine SW, Wilk KE, Andrews JR. Biomechanics of the knee during closed kinetic chain and open kinetic chain exercises. *Med Sci Sports Exerc.* 1998;30(4):556-569.
50. Mellor R, Hodges PW. Motor unit synchronization of the vasti muscles in closed and open chain tasks. *Arch Phys Med Rehabil.* 2005;86(4):716-721.
51. Stensdotter AK, Hodges PW, Mellor R, Sundelin G, H??ger-Ross C. Quadriceps Activation in Closed and in Open Kinetic Chain Exercise. *Med Sci Sports Exerc.* 2003;35(12):2043-2047.
52. Hug F, Hodges PW, Tucker K. Muscle force cannot be directly inferred from muscle activation: illustrated by the proposed imbalance of force between the vastus medialis and vastus lateralis in people with patellofemoral pain. *J Orthop Sports Phys Ther.* 2015;45(5):360-365.
53. Levinger P, Gilleard W, Coleman C. Femoral medial deviation angle during a one-leg squat test in individuals with patellofemoral pain syndrome. *Phys Ther Sport.* 2007;8(4):163-168.
54. Ageberg E, Bennell KL, Hunt M a, Simic M, Roos EM, Creaby MW. Validity and inter-rater reliability of medio-lateral knee motion observed during a single-limb mini squat. *BMC Musculoskelet Disord.* 2010;11(1):265.
55. Hwangbo P-N. The effects of squatting with visual feedback on the muscle activation of the vastus medialis oblique and the vastus lateralis in young adults with an increased quadriceps angle. *J Phys Ther Sci.* 2015;27(5):1507-1510.
56. Salsich GB. The Effects of Movement Pattern Modification on Lower Extremity Kinematics and Pain in Females With Patellofemoral Pain. *J Orthop Sports Phys Ther.* 2012;42(12):1017-1024.

57. Farrokhi S, Pollard CD, Souza RB, Chen Y-J, Reischl S, Powers CM. Trunk position influences the kinematics, kinetics, and muscle activity of the lead lower extremity during the forward lunge exercise. *J Orthop Sports Phys Ther.* 2008;38(7):403-409.
58. Shields RK, Madhavan S, Gregg E, et al. Neuromuscular Control of the Knee During a Resisted Single-Limb Squat Exercise. *Am J Sports Med.* 2005;33(10):1520-1526.
59. Toutoungi DE, Lu TW, Leardini A, Catani F, O'Connor JJ. Cruciate ligament forces in the human knee during rehabilitation exercises. *Clin Biomech* (Bristol, Avon). 2000;15(3):176-187.
60. Witvrouw E, Danneels L, Van Tiggelen D, Willems TM, Cambier D. Open versus closed kinetic chain exercises in patellofemoral pain: a 5-year prospective randomized study. *Am J Sports Med.* 2004;32(5):1122-1130.
61. Ernst GP, Saliba E, Diduch DR, Hurwitz SR, Ball DW. Lower extremity compensations following anterior cruciate ligament reconstruction. *Phys Ther.* 2000;80(3):251-260.
62. Salem GJ, Salinas R, Harding FV. Bilateral kinematic and kinetic analysis of the squat exercise after anterior cruciate ligament reconstruction. *Arch Phys Med Rehabil.* 2003;84(8):1211-1216.
63. Youdas JW, Hollman JH, Hitchcock JR, Hoyme GJ, Johnsen JJ. Comparison of hamstring and quadriceps femoris electromyographic activity between men and women during a single-limb squat on both a stable and labile surface. *J Strength Cond Res.* 2007;21(1):105-111.
64. Dye SF. The pathophysiology of patellofemoral pain: a tissue homeostasis perspective. *Clin Orthop Relat Res.* 2005;(436):100-110.
65. Wojtys EM, Beaman DN, Glover RA, Janda D. Innervation of the human knee joint by substance-P fibers. *Arthroscopy.* 1990;6(4):254-263.
66. Sanchis-Alfonso V, Roselló-Sastre E. Immunohistochemical analysis for neural markers of the lateral retinaculum in patients with isolated symptomatic patellofemoral malalignment. A neuroanatomic basis for anterior knee pain in the active young patient. *Am J Sports Med.* 28(5):725-731.
67. Sanchis-Alfonso V. [Patellofemoral pain]. *Der Orthopäde.* 2008;37(9):835-836, 838-840.
68. Rathleff MS, Roos EM, Olesen JL, Rasmussen S, Arendt-Nielsen L. Lower mechanical pressure pain thresholds in female adolescents with patellofemoral pain syndrome. *J Orthop Sports Phys Ther.* 2013;43(6):414-421.
69. Heijne A, Fleming BC, Renstrom PA, Peura GD, Beynonn BD, Werner S. Strain on the anterior cruciate ligament during closed kinetic chain exercises. *Med Sci Sports Exerc.* 2004;36(6):935-941.
70. Escamilla RF, Zheng N, MacLeod TD, et al. Patellofemoral compressive force and stress during the forward and side lunges with and without a stride. *Clin Biomech* (Bristol, Avon). 2008;23(8):1026-1037.
71. Begalle RL, DiStefano LJ, Blackburn T, Padua DA. Quadriceps and hamstrings coactivation during common therapeutic exercises. *J Athl Train.* 2012;47(4):396-405.
72. Seeley MK, Park J, King D, Ty Hopkins J. A novel experimental knee-pain model affects perceived pain and movement biomechanics. *J Athl Train.* 2013;48(3):337-345.
73. Willy RW, Scholz JP, Davis IS. Mirror gait retraining for the treatment of patellofemoral pain in female runners. *Clin Biomech* (Bristol, Avon). 2012;27(10):1045-1051.
74. Proske U, Morgan DL. Muscle damage from eccentric exercise: Mechanism, mechanical signs, adaptation and clinical applications. *J Physiol.* 2001;537(2):333-345.
75. McHugh MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand J Med Sci Sports.* 2003;13(2):88-97.
76. Kirkwood RN, Culham EG, Costigan P. Hip moments during level walking, stair climbing, and exercise in individuals aged 55 years or older. *Phys Ther.* 1999;79(4):360-370.
77. Wang MY, Flanagan S, Song JE, Greendale GA, Salem GJ. Lower-extremity biomechanics during forward and lateral stepping activities in older adults. *Clin Biomech.* 2003;18(3):214-221.
78. Chinkulprasert C, Vachalathiti R, Powers CM. Patellofemoral joint forces and stress during forward step-up, lateral step-up, and forward step-down exercises. *J Orthop Sports Phys Ther.* 2011;41(4):241-248.
79. Chang YJ, Chou CC, Chan HL, et al. Increases of quadriceps inter-muscular cross-correlation and coherence during exhausting stepping exercise. *Sensors (Switzerland).* 2012;12(12):16353-16367.
80. Enoka RM. Eccentric contractions require unique activation strategies by the nervous system. *J Appl Physiol.* 1996;4(4):2339-2346.
81. Pogosyan A, Gaynor LD, Eusebio A, Brown P. Boosting Cortical Activity at Beta-Band Frequencies Slows Movement in Humans. *Curr Biol.* 2009;19(19):1637-1641.
82. Gilbertson T, Lalo E, Doyle L, Di Lazzaro V, Cioni B, Brown P. Existing motor state is favored at the expense of new movement during 13-35 Hz oscillatory synchrony in the human corticospinal system. *J Neurosci.* 2005;25(34):7771-7779.
83. Androulidakis AG, Doyle LMF, Gilbertson TP, Brown P. Corrective movements in response to displacements in visual feedback are more effective during periods of 13-35 Hz oscillatory synchrony in the human corticospinal system. *Eur J Neurosci.* 2006;24(11):3299-3304.

84. Lalo E, Gilbertson T, Doyle L, Di Lazzaro V, Cioni B, Brown P. Phasic increases in cortical beta activity are associated with alterations in sensory processing in the human. *Exp Brain Res*. 2007;177(1):137-145.
85. Bigland-Ritchie BR, Dawson NJ, Johansson RS, Lippold OC. Reflex origin for the slowing of motoneurone firing rates in fatigue of human voluntary contractions. *J Physiol*. 1986;379:451-459.
86. Arabadzchiev TI, Dimitrov G V., Dimitrova NA. Intracellular action potential generation and extinction strongly affect the sensitivity of M-wave characteristic frequencies to changes in the peripheral parameters with muscle fatigue. *J Electromyogr Kinesiol*. 2005;15(2):159-169.
87. De Luca CJ, Erim Z. Common drive in motor units of a synergistic muscle pair. *J Neurophysiol*. 2002;87(4):2200-2204.
88. Miller R, Giannini D, Milner-Brown H, et al. Effects of fatiguing exercise on high-energy phosphates, force, and EMG: Evidence for three phases of recovery. *Muscle* 1987;10(9):810-821.
89. Taylor AD, Bronks R, Smith P, Humphries B. Myoelectric evidence of peripheral muscle fatigue during exercise in severe hypoxia: some references to m. vastus lateralis myosin heavy chain composition. *Eur J Appl Physiol Occup Physiol*. 1997;75(2):151-159.
90. Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev*. 2001;81(4):1725-1789.
91. Huang YM, Hsu MJ, Lin CH, Wei SH, Chang YJ. The non-linear relationship between muscle voluntary activation level and voluntary force measured by the interpolated twitch technique. *Sensors*. 2010;10(1):796-807.
92. Kettunen J a, Harilainen A, Sandelin J, et al. Knee arthroscopy and exercise versus exercise only for chronic patellofemoral pain syndrome: a randomized controlled trial. *BMC Med*. 2007;5:38.
93. Enoka RM, Duchateau J. Muscle fatigue: what, why and how it influences muscle function. *J Physiol*. 2008;586:11-23.
94. Cairns SP, Knicker AJ, Thompson MW, Sjøgaard G. Evaluation of models used to study neuromuscular fatigue. *Exerc Sport Sci Rev*. 2005;33(1):9-16.
95. Weeks BK, Carty CP, Horan SA. Effect of sex and fatigue on single leg squat kinematics in healthy young adults. *BMC Musculoskelet Disord*. 2015;16(1):271.
96. Schwellnus MP. Cause of exercise associated muscle cramps (EAMC)--altered neuromuscular control, dehydration or electrolyte depletion? *Br J Sports Med*. 2009;43:401-408.
97. Weeks BK, Carty CP, Horan SA. Effect of sex and fatigue on single leg squat kinematics in healthy young adults. *BMC Musculoskelet Disord*. 2015;16(1):271.
98. Schwellnus MP. Cause of exercise associated muscle cramps (EAMC)--altered neuromuscular control, dehydration or electrolyte depletion? *Br J Sports Med*. 2009;43:401-408.



## REFERENCES

1. Kaya D, Doral MN, Callaghan M. How can we strengthen the quadriceps femoris in patients with patellofemoral pain syndrome? *Muscles Ligaments Tendons J.* 2012;2(19):25-32.
2. Herrington L, Al-Sherhi A. A controlled trial of weight-bearing versus non-weight-bearing exercises for patellofemoral pain. *J Orthop Sports Phys Ther.* 2007;37(4):155-160.
3. Natri A, Kannus P, Järvinen M. Which factors predict the long-term outcome in chronic patellofemoral pain syndrome? A 7-yr prospective follow-up study. *Med Sci Sports Exerc.* 1998;30(11):1572-1577.
4. Harvie D, O'leary T, Kumar S. A systematic review of randomized controlled trials on exercise parameters in the treatment of patellofemoral pain: What works? *J Multidiscip Healthc.* 2011;4:383-392.
5. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther.* 2009;39(1):12-19.
6. Dolak KL, Silkman C, Medina McKeon J, Hosey RG, Lattermann C, Uhl TL. Hip strengthening prior to functional exercises reduces pain sooner than quadriceps strengthening in females with patellofemoral pain syndrome: a randomized clinical trial. *J Orthop Sports Phys Ther.* 2011;41(8):560-570.
7. Nakagawa TH, Muniz TB, Baldon RDM, Dias Maciel C, de Menezes Reiff RB, Serrão FV. The effect of additional strengthening of hip abductor and lateral rotator muscles in patellofemoral pain syndrome: a randomized controlled pilot study. *Clin Rehabil.* 2008;22(12):1051-1060.
8. Ismail MM, Gamaleldein MH, Hassa KA. Closed Kinetic Chain exercises with or without additional hip strengthening exercises in management of Patellofemoral pain syndrome: A randomized controlled trial. *Eur J Phys Rehabil Med.* 2013;49(5):687-698.
9. Fukuda TY. Hip Posterolateral Musculature Strengthening in Sedentary Women With Patellofemoral Pain Syndrome: A Randomized Controlled Clinical Trial With 1-Year Follow-up. *J Orthop Sports Phys Ther.* 2012;42(10):823-830.
10. Khayambashi K, Fallah A, Movahedi A, Bagwell J, Powers C. Posterolateral hip muscle strengthening versus quadriceps strengthening for patellofemoral pain: A comparative control trial. *Arch Phys Med Rehabil.* 2014;95(5):900-907.
11. Baldon RDM, Serrão FV, Scattone Silva R, Piva SR. Effects of functional stabilization training on pain, function, and lower extremity biomechanics in women with patellofemoral pain: a randomized clinical trial. *J Orthop Sports Phys Ther.* 2014;44(4):240-A8.
12. Lack S, Barton C, Sohan O, Crossley K, Morrissey D. Proximal muscle rehabilitation is effective for patellofemoral pain: a systematic review with meta-analysis. *Br J Sport Med.* 2015;49:1365-1376.
13. Ugalde V, Brockman C, Bailowitz Z, Pollard CD. Single Limb Squat Test and its Relationship to Dynamic Knee Valgus and Injury Risk Screening. *PM R.* 2015.7(3):229-35
14. Dimattia M a, Livengood AL, Uhl TL, Mattacola CG, Malone TR. What Are the Validity of the Single-Leg-Squat Test and Its Relationship to Hip-Abduction Strength? *J Sport Rehabil.* 2005;14(2):108-124.
15. Souza RB, Powers CM. Predictors of hip internal rotation during running: an evaluation of hip strength and femoral structure in women with and without patellofemoral pain. *Am J Sports Med.* 2009;37(3):579-587.
16. Hodges PW, Richardson CA. The influence of isometric hip adduction on quadriceps femoris activity. *Scand J Rehabil Med.* 1993;25(2):57-62.
17. Hanten WP, Schulthies SS. Exercise effect on electromyographic activity of the vastus medialis oblique and vastus lateralis muscles. *Phys Ther.* 1990;70(9):561-565.
18. Peng H Te, Kernozek TW, Song CY. Muscle activation of vastus medialis obliquus and vastus lateralis during a dynamic leg press exercise with and without isometric hip adduction. *Phys Ther Sport.* 2013;14(1):44-49.
19. Powers CM, Ward SR, Fredericson M, Guillet M, Shellock FG. Patellofemoral kinematics during weight-bearing and non-weight-bearing knee extension in persons with lateral subluxation of the patella: a preliminary study. *J Orthop Sports Phys Ther.* 2003;33(11):677-685.
20. Hollman JH, Ginos BE, Kozuchowski J, Vaughn AS, Krause D a, Youdas JW. Relationships between knee valgus, hip-muscle strength, and hip-muscle recruitment during a single-limb step-down. *J Sport Rehabil.* 2009;18(1):104-117.
21. Willson JD, Davis IS. Utility of the frontal plane projection angle in females with patellofemoral pain. *J Orthop Sports Phys Ther.* 2008;38(10):606-615.
22. Salsich GB. The Effects of Movement Pattern Modification on Lower Extremity Kinematics and Pain in Females With Patellofemoral Pain. *J Orthop Sports Phys Ther.* 2012;42(12):1017-1024.
23. Moyano FR, Valenza MC, Martin LM, Caballero YC, Gonzalez-Jimenez E, Demet GV. Effectiveness of different exercises and stretching physiotherapy on pain and movement in patellofemoral pain syndrome: a randomized controlled trial. *Clin Rehabil.* 2013;27(5):409-417.
24. Alba-Martín P, Gallego-Izquierdo T, Plaza-Manzano G, Romero-Franco N, Núñez-Nagy S, Pecos-Martín D. Effectiveness of therapeutic physical exercise in the treatment of patellofemoral pain syndrome: a systematic review. *J Phys Ther Sci.* 2015;27(7):2387-2390.
25. Mason M, Keays SL, Newcombe PA. The Effect of Taping, Quadriceps Strengthening and Stretching Prescribed Separately or Combined on Patellofemoral Pain. *Physiother Res Int.* 2011;16(2):109-119.

26. Peeler J, Anderson JE. Effectiveness of static quadriceps stretching in individuals with patellofemoral joint pain. *Clin J Sport Med.* 2007;17(4):234-241.
27. White LC, Dolphin P, Dixon J. Hamstring length in patellofemoral pain syndrome. *Physiotherapy.* 2009;95(1):24-28.
28. Mandelbaum BR, Silvers HJ, Watanabe DS, et al. Effectiveness of a neuromuscular and proprioceptive training program in preventing anterior cruciate ligament injuries in female athletes: 2-year follow-up. *Am J Sports Med.* 2005;33(7):1003-1010.
29. Caplan N, Kader DF. Knee injury patterns among men and women in collegiate basketball and soccer: NCAA data and review of literature. In: *Classic Papers in Orthopaedics.*; 2014:153-155.
30. Agel J, Arendt E a, Bershadsky B. Anterior cruciate ligament injury in national collegiate athletic association basketball and soccer: a 13-year review. *Am J Sports Med.* 2005;33(4):524-530.
31. Harmon KG, Ireland ML. Gender differences in noncontact anterior cruciate ligament injuries. *Clin Sports Med.* 2000;19(2):287-302.
32. Myer GD, Ford KR, Hewett TE. Rationale and clinical techniques for anterior cruciate ligament injury prevention among female athletes. *J Athl Train.* 2004;39(4):352-364.
33. Hewett TE, Stroupe a L, Nance T a, Noyes FR. Plyometric training in female athletes. Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996;24(6):765-773.
34. Li G, Rudy TW, Sakane M, Kanamori A, Ma CB, Woo SLY. The importance of quadriceps and hamstring muscle loading on knee kinematics and in-situ forces in the ACL. *J Biomech.* 1999;32(4):395-400.
35. Besier TF, Lloyd DG, Ackland TR. Muscle activation strategies at the knee during running and cutting maneuvers. *Med Sci Sports Exerc.* 2003;35(1):119-127.
36. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33(4):492-501.
37. Hewett TE, Paterno M V, Myer GD. Strategies for enhancing proprioception and neuromuscular control of the knee. *Clin Orthop Relat Res.* 2002;(402):76-94.
38. Ford KR, Myer GD, Schmitt LC, Uhl TL, Hewett TE. Preferential quadriceps activation in female athletes with incremental increases in landing intensity. *J Appl Biomech.* 2011;27(3):215-222.
39. Youdas JW, Hollman JH, Hitchcock JR, Hoyme GJ, Johnsen JJ. Comparison of hamstring and quadriceps femoris electromyographic activity between men and women during a single-limb squat on both a stable and labile surface. *J Strength Cond Res.* 2007;21(1):105-111.
40. Landry SC, McKean KA, Hubley-Kozey CL, Stanish WD, Deluzio KJ. Unanticipated running and cross-cutting maneuvers demonstrate neuromuscular and lower limb biomechanical differences between elite adolescent male and female soccer players. *J Biomech.* 2007;40:S379.
41. Malinzak RA, Colby SM, Kirkendall DT, Yu B, Garrett WE. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech.* 2001;16(5):438-445.
42. Chappell JD, Limpisvasti O. Effect of a neuromuscular training program on the kinetics and kinematics of jumping tasks. *Am J Sports Med.* 2008;36(6):1081-1086.
43. Aerts I, Cumps E, Verhagen E, Meeusen R. Efficacy of a 3 month training program on the jump-landing technique in jump-landing sports. Design of a cluster randomized controlled trial. *BMC Musculoskelet Disord.* 2010;11(1):281.
44. Fong CM, Blackburn JT, Norcross MF, McGrath M, Padua DA. Ankle-dorsiflexion range of motion and landing biomechanics. *J Athl Train.* 2011;46(1):5-10.
45. Hoch MC, Farwell KE, Gaven SL, Weinhandl JT. Weight-Bearing Dorsiflexion Range of Motion and Landing Biomechanics in Individuals With Chronic Ankle Instability. *J Athl Train.* 2015;50(5):150611075825000.
46. Kosashvili Y, Fridman T, Backstein D, Safir O, Bar Ziv Y. The correlation between pes planus and anterior knee or intermittent low back pain. *Foot ankle Int / Am Orthop Foot Ankle Soc [and] Swiss Foot Ankle Soc.* 2008;29(9):910-913.
47. De Oliveira Silva D, Briani RV, Pazzinatto MF, et al. Reliability and differentiation capability of dynamic and static kinematic measurements of rearfoot eversion in patellofemoral pain. *Clin Biomech.* 2015;30(2):144-148.
48. Caylor D, Fites R, Worrell TW. The relationship between quadriceps angle and anterior knee pain syndrome. *J Orthop Sports Phys Ther.* 1993;17(1):11-16.
49. Emami MJ, Ghahramani MH, Abdinejad F, Namazi H. Q-angle: An invaluable parameter for evaluation of anterior knee pain. *Arch Iran Med.* 2007;10(1):24-36.
50. Park SK, Stefanyshyn DJ. Greater Q angle may not be a risk factor of Patellofemoral Pain Syndrome. *Clin Biomech.* 2011;26(4):392-396.
51. Sheehan FT, Derasari A, Fine KM, Brindle TJ, Alter KE. Q-angle and J-sign: Indicative of maltracking subgroups in patellofemoral pain. *Clin Orthop Relat Res.* 2010;468(1):266-275.
52. Souza RB, Draper CE, Fredericson M, Powers CM. Femur rotation and patellofemoral joint kinematics: a weight-bearing magnetic resonance imaging analysis. *J Orthop Sports Phys Ther.* 2010;40(5):277-285.
53. Puniello MS. Iliotibial band tightness and medial patellar glide in patients with patellofemoral dysfunction. *J Orthop Sports Phys Ther.* 1993;17(3):144-148.
54. Winslow, J Yoder E. Patellofemoral pain in female ballet dancers: correlation with iliotibial band tightness and tibial external rotation. *J Orthop Sports Phys Ther.* 1995;22:18-21.

55. Merican AM, Amis AA. Iliotibial band tension affects patellofemoral and tibiofemoral kinematics. *J Biomech.* 2009;42(10):1539-1546.
56. Patil S, White L, Jones a, Hui a C. Idiopathic anterior knee pain in the young. A prospective controlled trial. *Acta Orthop Belg.* 2010;76(3):356-359.
57. Patil S, Dixon J, White LC, Jones AP, Hui ACW. An electromyographic exploratory study comparing the difference in the onset of hamstring and quadriceps contraction in patients with anterior knee pain. *Knee.* 2011;18(5):329-332.
58. Clark DI, Downing N, Mitchell J, Coulson L, Syzpryt EP, Doherty M. Physiotherapy for anterior knee pain: a randomised controlled trial. *Ann Rheum Dis.* 2000;59(9):700-704.
59. Lin P-L, Shih Y-F, Chen W-Y, Ma H-L. Neurodynamic responses to the femoral slump test in patients with anterior knee pain syndrome. *J Orthop Sports Phys Ther.* 2014;44(5):350-357.
60. Lai WH, Shih YF, Lin PL, Chen WY, Ma HL. Normal neurodynamic responses of the femoral slump test. *Man Ther.* 2012;17(2):126-132.
61. Clark DA, Simpson DL, Eldridge J, Colborne GR. Patellar instability and quadriceps avoidance affect walking knee moments. *Knee.* 2016;23(1):78-84.
62. Juhn MS. Patellofemoral pain syndrome: a review and guidelines for treatment. *Am Fam Physician.* 1999;60(7):2012-2022.
63. Herrington L, Nester C. Q-angle undervalued? The relationship between Q-angle and medio-lateral position of the patella. *Clin Biomech.* 2004;19(10):1070-1073.
64. Biedert RM, Warnke K. Correlation between the Q angle and the patella position: A clinical and axial computed tomography evaluation. *Arch Orthop Trauma Surg.* 2001;121(6):346-349.
65. Freedman BR, Brindle TJ, Sheehan FT. Re-evaluating the functional implications of the Q-angle and its relationship to in-vivo patellofemoral kinematics. *Clin Biomech.* 2014;29(10):1139-1145.
66. Bolgla L, Malone T. Exercise Prescription and Patellofemoral Pain : Evidence for Rehabilitation. *J Sport Rehabil.* 2005;14(1):72-88.
67. Bolgla L a, Boling MC. An update for the conservative management of patellofemoral pain syndrome: a systematic review of the literature from 2000 to 2010. *Int J Sports Phys Ther.* 2011;6(2):112-125.
68. Barton CJ, Lack S, Hemmings S, Tufail S, Morrissey D. The "Best Practice Guide to Conservative Management of Patellofemoral Pain": incorporating level 1 evidence with expert clinical reasoning. *Br J Sports Med.* 2015;49(14):923-934.
69. Ho KY, Blanchette MG, Powers CM. The influence of heel height on patellofemoral joint kinetics during walking. *Gait Posture.* 2012;36(2):271-275.
70. Lenhart RL, Thelen DG, Wille CM, Chumanov ES, Heiderscheit BC. Increasing running step rate reduces patellofemoral joint forces. *Med Sci Sports Exerc.* 2014;46(3):557-564.
71. Willson JD, Ratcliff OM, Meardon SA, Willy RW. Influence of step length and landing pattern on patellofemoral joint kinetics during running. *Scand J Med Sci Sport.* 2015;25(6):736-743.
72. Chumanov ES, Wille CM, Michalski MP, Heiderscheit BC. Changes in muscle activation patterns when running step rate is increased. *Gait Posture.* 2012;36(2):231-235.
73. Heiderscheit BC, Chumanov ES, Michalski MP, Wille CM, Ryan MB. Effects of step rate manipulation on joint mechanics during running. *Med Sci Sports Exerc.* 2011;43(2):296-302.
74. Vannatta CN, Kernozek TW. Patellofemoral joint stress during running with alterations in foot strike pattern. *Med Sci Sports Exerc.* 2015;47(5):1001-1008.
75. Bonacci J, Vicenzino B, Spratford W, Collins P. Take your shoes off to reduce patellofemoral joint stress during running. *Br J Sports Med.* 2014;48(6):425-428.
76. McCarthy C, Fleming N, Donne B, Blanksby B. Barefoot running and hip kinematics: Good news for the knee? *Med Sci Sports Exerc.* 2015;47(5):1009-1016.
77. Cheung RTH, Ng GYF, Chen BFC. Association of footwear with patellofemoral pain syndrome in runners. *Sports Med.* 2006;36(3):199-205.
78. Cheung RTH, Ng GYF. Motion control shoe affects temporal activity of quadriceps in runners. *Br J Sports Med.* 2009;43(12):943-947.
79. Neptune RR, Wright IC, Van Den Bogert AJ. The influence of orthotic devices and vastus medialis strength and timing on patellofemoral loads during running. *Clin Biomech.* 2000;15(8):611-618.
80. Teng H-L, Powers CM. Sagittal plane trunk posture influences patellofemoral joint stress during running. *J Orthop Sports Phys Ther.* 2014;44(10):785-792.
81. Sussman DH, Alrowayeh H, Walker ML. Patellofemoral Joint Compressive Forces During Backward and Forward Running At the Same Speed. *J Musculoskelet Res.* 2000;4(2):107-118.
82. Flynn TW, Soutas-Little RW. Patellofemoral joint compressive forces in forward and backward running. *J Orthop Sports Phys Ther.* 1995;21(5):277-282.
83. Roos PE, Barton N, van Deursen RWM. Patellofemoral joint compression forces in backward and forward running. *J Biomech.* 2012;45(9):1656-1660.
84. Piva SR, Fitzgerald GK, Wisniewski S, Delitto A. Predictors of pain and function outcome after rehabilitation in patients with patellofemoral pain syndrome. *J Rehabil Med.* 2009;41(8):604-612.

85. Halabchi F, Mazaheri R, Mansournia MA, Hamed Z. Additional Effects of an Individualized Risk Factor-Based Approach on Pain and the Function of Patients With Patellofemoral Pain Syndrome: A Randomized Controlled Trial. *Clin J Sport Med*. 2015;25(6):478-486.
86. Zakaria D, Harburn KL, Kramer JF. Preferential activation of the vastus medialis oblique, vastus lateralis, and hip adductor muscles during isometric exercises in females. *J Orthop Sports Phys Ther*. 1997;26(1):23-28.
87. Tepperman PS, Mazliah J, Naumann S, Delmore T. Effect of ankle position on isometric quadriceps strengthening. *Am J Phys Med*. 1986;65(2):69-74.
88. Bolgla L a, Shaffer SW, Malone TR. Vastus medialis activation during knee extension exercises: evidence for exercise prescription. *J Sport Rehabil*. 2008;17(1):1-10.
89. Malone T, Davies G, Walsh WM. Muscular control of the patella. *Clin Sports Med*. 2002;21(3):349-362.
90. Powers CM, Perry J, Hsu a, Hislop HJ. Are patellofemoral pain and quadriceps femoris muscle torque associated with locomotor function? *Phys Ther*. 1997;77(10):1063-1075-1078.
91. Kooiker L, Van De Port IGL, Weir A, Moen MH. Effects of physical therapist-guided quadriceps-strengthening exercises for the treatment of patellofemoral pain syndrome: a systematic review. *J Orthop Sports Phys Ther*. 2014;44(6):391-B1.
92. Gryzlo SM, Patek RM, Pink M, Perry J. Electromyographic analysis of knee rehabilitation exercises. *J Orthop Sports Phys Ther*. 1994;20(1):36-43.
93. Spencer JD, Hayes KC, Alexander IJ. Knee joint effusion and quadriceps reflex inhibition in man. *Arch Phys Med Rehabil*. 1984;65(4):171-177.
94. Palmieri-Smith RM, Kreinbrink J, Ashton-Miller JA, Wojtys EM. Quadriceps inhibition induced by an experimental knee joint effusion affects knee joint mechanics during a single-legged drop landing. *Am J Sports Med*. 2007;35(8):1269-1275.
95. Fahrer H, Rentsch HU, Gerber NJ, Beyeler C, Hess CW, Grünig B. Knee effusion and reflex inhibition of the quadriceps. A bar to effective retraining. *J Bone Joint Surg Br*. 1988;70:635-638.
96. Lepley AS, Bahhur NO, Murray AM, Pietrosimone BG. Quadriceps corticomotor excitability following an experimental knee joint effusion. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2013:1-8.
97. Palmieri RM, Ingersoll CD, Edwards JE, et al. Arthrogenic muscle inhibition is not present in the limb contralateral to a simulated knee joint effusion. *Am J Phys Med Rehabil*. 2003;32(12):910-916.
98. Torry MR, Decker MJ, Viola RW, O'Connor DD, Richard Steadman J. Intra-articular knee joint effusion induces quadriceps avoidance gait patterns. *Clin Biomech*. 2000;15(3):147-159.
99. Hopkins JT, Ingersoll CD, Edwards J, Klootwyk TE. Cryotherapy and transcutaneous electric neuromuscular stimulation decrease arthrogenic muscle inhibition of the vastus medialis after knee joint effusion. *J Athl Train*. 2002;37(1):25-31.
100. Hopkins JT. Knee joint effusion and cryotherapy alter lower chain kinetics and muscle activity. *J Athl Train*. 2006;41(2):177-184.
101. Toutoungi DE, Lu TW, Leardini A, Catani F, O'Connor JJ. Cruciate ligament forces in the human knee during rehabilitation exercises. *Clin Biomech (Bristol, Avon)*. 2000;15(3):176-187.
102. Beynnon B, Howe JG, Pope MH, Johnson RJ, Fleming BC. The measurement of anterior cruciate ligament strain in vivo. *Int Orthop*. 1992;16(1):1-12.
103. Beynnon BD, Fleming BC, Johnson RJ, Nichols CE, Renström P a, Pope MH. Anterior cruciate ligament strain behavior during rehabilitation exercises in vivo. *Am J Sports Med*. 1993;23(1):24-34.
104. Fleming BC, Oksendahl H, Beynnon BD. Open- or closed-kinetic chain exercises after anterior cruciate ligament reconstruction? *Exerc Sport Sci Rev*. 2005;33(3):134-140.
105. Escamilla RF, Fleisig GS, Zheng N, Barrentine SW, Wilk KE, Andrews JR. Biomechanics of the knee during closed kinetic chain and open kinetic chain exercises. *Med Sci Sports Exerc*. 1998;30(4):556-569.
106. Escamilla RF, Macleod TD, Wilk KE, Paulos L, Andrews JR. Anterior cruciate ligament strain and tensile forces for weight-bearing and non-weight-bearing exercises: a guide to exercise selection. *J Orthop Sports Phys Ther*. 2012;42(3):208-220.
107. Witvrouw E, Lysens R, Bellemans J, Peers K, Vanderstraeten G. Open versus closed kinetic chain exercises for patellofemoral pain. A prospective, randomized study. *Am J Sports Med*. 2000;28(5):687-694.
108. Kushion D. EMG Activation of the Vastus Medialis Oblique and Vastus Lateralis During Four Rehabilitative Exercises. *Open Rehabil J*. 2012;5(1):1-7.
109. Lieb FJ, Perry J. Quadriceps function. An anatomical and mechanical study using amputated limbs. *J Bone Joint Surg Am*. 1968;50(8):1535-1548.
110. Peeler J, Cooper J, Porter MM, Thliveris JA, Anderson JE. Structural parameters of the vastus medialis muscle. *Clin Anat*. 2005;18(4):281-289.
111. Hubbard JK, Sampson HW, Elledge JR. Prevalence and morphology of the vastus medialis oblique muscle in human cadavers. *Anat Rec*. 1997;249(1):135-142.
112. Nozic M, Mitchell J, de Klerk D, Nozic M. A comparison of the proximal and distal parts of the vastus medialis muscle. *Aust J Physiother*. 1997;43(4):277-281.
113. Smith TO, Nichols R, Harle D, Donell ST. Do the vastus medialis obliquus and vastus medialis longus really exist? A systematic review. *Clin Anat*. 2009;22(2):183-199.



114. Tenan MS, Peng Y-L, Hackney AC, Griffin L. Menstrual cycle mediates vastus medialis and vastus medialis oblique muscle activity. *Med Sci Sports Exerc.* 2013;45(11):2151-2157.
115. Lieb FJ, Perry J. Quadriceps function. An electromyographic study under isometric conditions. *J Bone Joint Surg Am.* 1971;53(4):749-758.
116. Giles LS, Webster KE, McClelland J a, Cook J. Does quadriceps atrophy exist in individuals with patellofemoral pain? A systematic literature review with meta-analysis. *J Orthop Sports Phys Ther.* 2013;43(11):766-776.
117. Pattyn E, Verdonk P, Steyaert A, et al. Vastus medialis obliquus atrophy: does it exist in patellofemoral pain syndrome? *Am J Sports Med.* 2011;39(7):1450-1455.
118. Callaghan MJ, Oldham J a. Quadriceps atrophy: to what extent does it exist in patellofemoral pain syndrome? *Br J Sports Med.* 2004;38(3):295-299.
119. Gerber C, Hoppeler H, Claassen H, Robotti G, Zehnder R, Jakob RP. The lower-extremity musculature in chronic symptomatic instability of the anterior cruciate ligament. *J Bone Joint Surg Am.* 1985;67(7):1034-1043.
120. Smillie I. *Injuries of the Knee Joint.* 3rd Edition. Baltimore, MD: Williams and Wilkins Publishers; 1962.
121. Janks ES. The role of the quadriceps muscle in knee extension. *Physiotherapy.* 1983;39(1):13-14.
122. Brewerton DA. The function of the vastus medialis muscle. *Rheumatology.* 1955;2(5):164-168. doi:10.1093/rheumatology/2.5.164.
123. Smillie I. *Injuries of the Knee: Importance of the Quadriceps.* New York: Churchill Livingstone; 1970.
124. N P, D F, Soames R. *Anatomy and Human Movement: Structure and Function.* Oxford, UK: Heinemann Medical Books; 1989.
125. MacKinnon P, Morris J. *Textbook of Functional Anatomy. Volume 1: Musculo-Skeletal System.* Oxford, UK: Oxford University Press; 1986.
126. Reider B, Marshall JL, Koslin B, Ring B, Girgis FG. The anterior aspect of the knee joint. *J Bone Joint Surg Am.* 1981;63(3):351-356.
127. Nissen CW, Cullen MC, Hewett TE, Noyes FR. Physical and arthroscopic examination techniques of the patellofemoral joint. *J Orthop Sports Phys Ther.* 1998;28(5):277-285.
128. Grob K, Manestar M, Figueira L, Ackland T, Gilbey H, Kuster MS. New insight in the architecture of the quadriceps tendon Journal of Experimental Orthopaedics. *J Exp Orthop.* 2016;3.
129. Salzman A, Torburn L, Perry J. Contribution of rectus femoris and vasti to knee extension. An electromyographic study. *Clin Orthop Relat Res.* 1993;(290):236-243.
130. Toumi H, Poumarat G, Benjamin M, Best T, F'Guyer S, Fairclough J. New insights into the function of the vastus medialis with clinical implications. *Med Sci Sports Exerc.* 2007;39(7):1153-1159.
131. Grabiner MD, Koh TJ, Von Haefen L. Decreased excitation of vastus medialis oblique and vastus lateralis in patellofemoral pain. *Eur J Exp Musculoskelet Res.* 1993;1:121-124.
132. Lee TQ, Sandusky MD, Adeli A, McMahon PJ. Effects of simulated vastus medialis strength variation on patellofemoral joint biomechanics in human cadaver knees. *J Rehabil Res Dev.* 2002;39(3):429-438.
133. Mirzabeigi E, Jordan C, Gronley JK, Rockowitz NL, Perry J. Isolation of the vastus medialis oblique muscle during exercise. *Am J Sports Med.* 1999;27(1):50-53.
134. Cowan SM, Bennell KL, Hodges PW. Therapeutic patellar taping changes the timing of vasti muscle activation in people with patellofemoral pain syndrome. *Clin J Sport Med.* 2002;12(6):339-347.
135. Cerny K. Vastus medialis oblique/vastus lateralis muscle activity ratios for selected exercises in persons with and without patellofemoral pain syndrome. *Phys Ther.* 1995;75(8):672-683.
136. Jang E-M, Heo H-J, Kim M-H, Yoo W-G. Activation of VMO and VL in Squat Exercises for Women with Different Hip Adduction Loads. *J Phys Ther Sci.* 2013;25(3):257-258.
137. Wong YM, Straub RK, Powers CM. The VMO: VL activation ratio while squatting with hip adduction is influenced by the choice of recording electrode. *J Electromyogr Kinesiol.* 2013;23(2):443-447.
138. Boling M, Padua D, Blackburn J. Hip Adduction Does not Affect VMO EMG Amplitude or VMO: VL Ratios During a Dynamic Squat Exercise. *J Sport Rehabil.* 2006;15(VI):195-205.
139. Willis FB, Burkhardt EJ, Walker JE, Johnson M a, Spears TD. Preferential vastus medialis oblique activation achieved as a treatment for knee disorders. *J Strength Cond Res.* 2005;19(2):286-291.
140. Gregersen CS, Hull ML, Hakansson NA. How Changing the Inversion/Eversion Foot Angle Affects the Nondriving Intersegmental Knee Moments and the Relative Activation of the Vastii Muscles in Cycling. *J Biomech Eng.* 2006;128(3):391.
141. Smith TO, Bowyer D, Dixon J, Stephenson R, Chester R, Donell ST. Can vastus medialis oblique be preferentially activated? A systematic review of electromyographic studies. *Physiother Theory Pract.* 2009;25(2):69-98.
142. Chester R, Smith TO, Sweeting D, Dixon J, Wood S, Song F. The relative timing of VMO and VL in the aetiology of anterior knee pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord.* 2008;9:64.
143. Cesarelli M, Bifulco P, Bracale M. Study of the control strategy of the quadriceps muscles in anterior knee pain. *IEEE Trans Rehabil Eng.* 2000;8(3):330-341.
144. Cowan SM, Bennell KL, Hodges PW, Crossley KM, McConnell J. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Arch Phys Med Rehabil.* 2001;82(2):183-189.

145. Van Tiggelen D, Cowan S, Coorevits P, Duvigneaud N, Witvrouw E. Delayed vastus medialis obliquus to vastus lateralis onset timing contributes to the development of patellofemoral pain in previously healthy men: a prospective study. *Am J Sports Med.* 2009;37(6):1099-1105.
146. Souza DR, Gross MT. Comparison of vastus medialis obliquus: vastus lateralis muscle integrated electromyographic ratios between healthy subjects and patients with patellofemoral pain. *Phys Ther.* 1991;71(4):310-320.
147. Syme G, Rowe P, Martin D, Daly G. Disability in patients with chronic patellofemoral pain syndrome: A randomised controlled trial of VMO selective training versus general quadriceps strengthening. *Man Ther.* 2009;14(3):252-263.
148. Coqueiro KRR, Bevilaqua-Grossi D, B??rzin F, Soares AB, Candolo C, Monteiro-Pedro V. Analysis on the activation of the VMO and VLL muscles during semisquat exercises with and without hip adduction in individuals with patellofemoral pain syndrome. *J Electromyogr Kinesiol.* 2005;15(6):596-603.
149. Earl JE, Schmitz RJ, Arnold BL. Activation of the VMO and VL during dynamic mini-squat exercises with and without isometric hip adduction. *J Electromyogr Kinesiol.* 2001;11(6):381-386.
150. Roush MB, Sevier TL, Wilson JK, et al. Anterior knee pain: a clinical comparison of rehabilitation methods. *Clin J Sport Med.* 2000;10(1):22-28.
151. Cavazzuti L, Merlo A, Orlandi F, Campanini I. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Gait Posture.* 2010;32(3):290-295.
152. Karst GM, Willett GM. Onset timing of electromyographic activity in the vastus medialis oblique and vastus lateralis muscles in subjects with and without patellofemoral pain syndrome. *Phys Ther J.* 1995;75(9):813-823.
153. Powers CM, Landel R, Perry J. Timing and intensity of vastus muscle activity during functional activities in subjects with and without patellofemoral pain. *Phys Ther.* 1996;76(9):946-55-67.
154. Powers CM. Patellar Kinematics, Part I: The Influence of Vastus Muscle Activity in Subjects With and Without Patellofemoral Pain. *Phys Ther.* 2000;80(10):956-964.
155. Steinkamp LA, Dillingham MF, Markel MD, Hill JA, Kaufman KR. Biomechanical considerations in patellofemoral joint rehabilitation. *Am J Sport Med.* 1993;21(3):438-444.
156. Cohen Z a, Roglic H, Grelsamer RP, et al. Patellofemoral stresses during open and closed kinetic chain exercises. An analysis using computer simulation. *Am J Sports Med.* 2001;29(4):480-487.
157. Choi S-A, Cynn H-S, Yoon T-L, Choi W-J, Lee J-H. Effects of Ankle Dorsiflexion on Vastus Medialis Oblique and Vastus Lateralis Muscle Activity During Straight Leg Raise Exercise with Hip External Rotation in Patellofemoral Pain Syndrome. *J Musculoskelet Pain.* 2014;(November 2013):1-8.
158. Bakhtiary a H, Fatemi E. Open versus closed kinetic chain exercises for patellar chondromalacia. *Br J Sports Med.* 2008;42(December):99-102; discussion 102.
159. Thomeé R. A comprehensive treatment approach for patellofemoral pain syndrome in young women. *Phys Ther.* 1997;77(12):1690-1703.
160. Thomee R, Grimby G, Svantesson U, Osterberg U. Quadriceps muscle performance in sitting and standing in young women with patellofemoral pain syndrome and young healthy women. *Scand J Med Sci Sport.* 1996;6(4):233-241.
161. Hart JM, Pietrosimone B, Hertel J, Ingersoll CD. Quadriceps activation following knee injuries: a systematic review. *J Athl Train.* 2010;45(1):87-97.
162. Yosmaoglu HB, Kaya D, Guney H, et al. Is there a relationship between tracking ability, joint position sense, and functional level in patellofemoral pain syndrome? *Knee Surgery, Sport Traumatol Arthrosc.* 2013;21(11):2564-2571.
163. Steinkamp LA, Dillingham MF, Markel MD, Hill JA, Kaufman KR. Biomechanical considerations in patellofemoral joint rehabilitation. *Am J Sports Med.* 1993;21(3):438-444.
164. Stiene HA, Brosky T, Reinking MF, Nyland J, Mason MB. A comparison of closed kinetic chain and isokinetic joint isolation exercise in patients with patellofemoral dysfunction. *J Orthop Sports Phys Ther.* 1996;24(3):136-141.
165. Powers CM, Ho K-Y, Chen Y-J, Souza RB, Farrokhi S. Patellofemoral Joint Stress During Weight Bearing and Non-Weight Bearing Quadriceps Exercises. *J Orthop Sports Phys Ther.* 2014;44(5):320-327.
166. Ebben WP, Feldmann CR, Dayne a, Mitsche D, Alexander P, Knetzger KJ. Muscle activation during lower body resistance training. *Int J Sports Med.* 2009;30(1):1-8.
167. Witvrouw E, Danneels L, Van Tiggelen D, Willems TM, Cambier D. Open versus closed kinetic chain exercises in patellofemoral pain: a 5-year prospective randomized study. *Am J Sports Med.* 2004;32(5):1122-1130.
168. Lutz GE, Palmitier R a, An KN, Chao EY. Comparison of tibiofemoral joint forces during open-kinetic-chain and closed-kinetic-chain exercises. *J Bone Joint Surg Am.* 1993;75(5):732-739.
169. Mikkelsen C, Werner S, Eriksson E. Closed kinetic chain alone compared to combined open and closed kinetic chain exercises for quadriceps strengthening after anterior cruciate ligament reconstruction with respect to return to sports: a prospective matched follow-up study. *Knee Surg Sports Traumatol Arthrosc.* 2000;8(6):337-342
170. Uçar M, Koca I, Eroglu M, et al. Evaluation of open

and closed kinetic chain exercises in rehabilitation following anterior cruciate ligament reconstruction. *J Phys Ther Sci.* 2014;26(12):1875-1878.

171. Doucette SA, Child DD. The effect of open and closed chain exercise and knee joint position on patellar tracking in lateral patellar compression syndrome. *J Orthop Sports Phys Ther.* 1996;23(2):104-110.

172. McConnell J. The management of chondromalacia patellae: a long term solution. *Aust J Physiother.* 1986;32(4):215-223.

173. Campbell DE, Glenn W. Rehabilitation of knee flexor and knee extensor muscle strength in patients with meniscectomies, ligamentous repairs, and chondromalacia. *Phys Ther.* 1982;62(1):10-15.

174. Laprade J, Culham E, Brouwer B. Comparison of five isometric exercises in the recruitment of the vastus medialis oblique in persons with and without patellofemoral pain syndrome. *J Orthop Sport Phys Ther.* 1998;27(3):197-204.

175. Hazneci B, Yildiz Y, Sekir U, Aydin T, Kalyon TA. Efficacy of Isokinetic Exercise on Joint Position Sense and Muscle Strength in Patellofemoral Pain Syndrome. *Am J Phys Med Rehabil.* 2005; 84(7):521-7