

REFERENCES

1. Grävare Silbernagel K, Crossley KM. A Proposed Return-to-Sport Program for Patients With Midportion Achilles Tendinopathy: Rationale and Implementation. *J Orthop Sport Phys Ther.* 2015;45(11):876-886.
2. Kubo K, Kanehisa H, Ito M, Fukunaga T. Effects of isometric training on the elasticity of human tendon structures in vivo. *J Appl Physiol.* 2001;91(1):26-32
3. Arampatzis A, Karamanidis K, Albracht K. Adaptational responses of the human Achilles tendon by modulation of the applied cyclic strain magnitude. *J Exp Biol.* 2007;210(Pt 15):2743-2753.
4. Kongsgaard M, Reitelseder S, Pedersen TG, et al. Region specific patellar tendon hypertrophy in humans following resistance training. *Acta Physiol.* 2007; Oct;191(2):111-21.
5. Arampatzis A, Peper A, Bierbaum S, Albracht K. Plasticity of human Achilles tendon mechanical and morphological properties in response to cyclic strain. *J Biomech.* 2010;43(16):3073-3079.
6. Bohm S, Mersmann F, Tettke M, Kraft M, Arampatzis A. Human Achilles tendon plasticity in response to cyclic strain: Effect of rate and duration. *J Exp Biol.* 2014; 217(Pt 22):4010-7
7. Wiesinger HP, Kösters A, Müller E, Seynnes OR. Effects of Increased Loading on in Vivo Tendon Properties: A Systematic Review. *Med Sci Sports Exerc.* 2015; 47(9):1885-95
8. Kubo K, Miyazaki D, Yamada K, Shimoju S, Tsunoda N. Are the knee and ankle angles at contact related to the tendon properties of lower limbs in long distance runners? *Springerplus.* 2016; 24;5:151
9. Morgan GE, Martin R, Williams L, Pearce O, Morris K. Objective assessment of stiffness in Achilles tendinopathy: A novel approach using the MyotonPRO. *BMJ Open Sport Exerc Med.* 2018;4(1):e000446
10. Bohm S, Mersmann F, Arampatzis A. Human tendon adaptation in response to mechanical loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults. *Sport Med - Open.* 2015; 1(1):7.
11. Epro G, Mierau A, Doerner J, et al. The Achilles tendon is mechanosensitive in older adults: Adaptations following 14 weeks versus 1.5 years of cyclic strain exercise. *J Exp Biol.* 2017; 220(Pt 6):1008-1018
12. O'Neill S, Barry S, Watson P. Plantarflexor strength and endurance deficits associated with mid-portion Achilles tendinopathy: The role of soleus. *Phys Ther Sport.* 2019; 37:69-76.
13. Jorgensen MG, Andersen S, Ryg J, Masud T. Novel use of the Nintendo Wii Board for measuring isometric lower limb strength: A reproducible and valid method in older adults. *PLoS One.* 2015; 10(10):e0138660
14. Marmon AR, Pozzi F, Alnahdi AH, Zeni JA. The validity of plantarflexor strength measures obtained through hand-held dynamometry measurements of force. *Int J Sports Phys Ther.* 2013;8(6):820-7.
15. Hébert-Losier K, Wessman C, Alricsson M, Svantesson U. Updated reliability and normative values for the standing heel-rise test in healthy adults. *Physiother (United Kingdom).* 2017;103(4):446-452.
16. Griffith S, O'Neill S. Talking Tendinopathy & Solving the Soleus Conundrum with Dr Seth O'Neill. *British Journal of Sports Medicine.* <https://blogs.bmj.com/bjbm/2019/04/12/talking-tendinopathy-solving-the-soleus-conundrum-with-dr-seth-oneill/>. Published 2019. Accessed July 17, 2019.
17. Maganaris CN. Tendon conditioning: Artefact or property? *Proc R Soc B Biol Sci.* 2003;270 Suppl 1:S39-42.
18. Nuri L, Obst SJ, Newsham-West R, Barrett RS. The tendinopathic Achilles tendon does not remain iso-volumetric upon repeated loading: Insights from 3D ultrasound. *J Exp Biol.* 2017;220(Pt 17):3053-3061.
19. Stevens M, Tan C-W. Effectiveness of the Alfredson Protocol Compared With a Lower Repetition-Volume Protocol for Midportion Achilles Tendinopathy: A Randomized Controlled Trial. *J Orthop Sport Phys Ther.* 2014;44(2):59-67.
20. O'Neill S, Watson PJ, Barry S. Why are eccentric exercises effective for Achilles tendinopathy? *Int J Sports Phys Ther.* 2015;10(4):552-62.
21. Rees JD, Lichtwark G a, Wolman RL, Wilson a M. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an in vivo study in humans. *Rheumatology (Oxford).* 2008;47(10):1493-1497.
22. Henriksen M, Aaboe J, Bliddal H, Langberg H. Biomechanical characteristics of the eccentric Achilles tendon exercise. *J Biomech.* 2009;42(16):2702-7.
23. Grigg NL, Wearing SC, O'Toole JM, Smeathers JE. The effect of exercise repetition on the frequency characteristics of motor output force: Implications for Achilles tendinopathy rehabilitation. *J Sci Med Sport.* 2014;17(1):13-7
24. Grigg NL, Wearing SC, O'toole JM, Smeathers JE. Achilles tendinopathy modulates force frequency characteristics of eccentric exercise. *Med Sci Sports Exerc.* 2013;45(3):520-6.
25. Rees JD, Lichtwark GA, Wolman RL, Wilson AM. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an in vivo study in humans. *Rheumatology.* 2008;47(10):1493-7.
26. Rubin C, Turner AS, Bain S, Mallinckrodt C, McLeod K. Anabolism. Low mechanical signals strengthen long bones. *Nature.* 2001; 412(6847):603-4

27. O'Neill S, Radia J, Bird K, et al. Acute sensory and motor response to 45-s heavy isometric holds for the plantar flexors in patients with Achilles tendinopathy. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2018; 27(9):2765-2773
28. Rio E, Kidgell D, Purdam C, et al. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med*. 2015;49(19):1277-1283.
29. Rio E, Kidgell D, Purdam C, et al. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med*. 2015;(19):1277-83
30. Coombes BK, Tucker K, Vicenzino B, et al. Achilles and patellar tendinopathy display opposite changes in elastic properties: A shear wave elastography study. *Scand J Med Sci Sport*. 2018;28(3):1201-1208.
31. Edama M, Kubo M, Onishi H, et al. The twisted structure of the human Achilles tendon. *Scand J Med Sci Sport*. 2015;25(5):e497-503.
32. Counsel P, Comin J, Davenport M, Connell D. Pattern of Fascicular Involvement in Midportion Achilles Tendinopathy at Ultrasound. *Sports Health*. 2015;7(5):424-8.
33. Szaro P, Witkowski G, Smigielski R, Krajewski P, Ciszek B. Fascicles of the adult human Achilles tendon - An anatomical study. *Ann Anat*. 2009;191(6):586-93.

REFERENCES

1. Grävare Silbernagel K, Crossley KM. A Proposed Return-to-Sport Program for Patients With Midportion Achilles Tendinopathy: Rationale and Implementation. *J Orthop Sport Phys Ther.* 2015;45(11):876-886.
2. Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with achilles tendinopathy: A randomized controlled study. *Am J Sports Med.* 2007; 35(6):897-906.
3. Magnusson SP, Langberg H, Kjaer M. The pathogenesis of tendinopathy: Balancing the response to loading. *Nat Rev Rheumatol.* 2010; 6(5):262-8.
4. Silbernagel KG, Gustavsson A, Thomeé R, Karlsson J. Evaluation of lower leg function in patients with Achilles tendinopathy. *Knee Surgery, Sport Traumatol Arthrosc.* 2006; 14(11):1207-17.
5. Bohm S, Mersmann F, Arampatzis A. Human tendon adaptation in response to mechanical loading: a systematic review and meta-analysis of exercise intervention studies on healthy adults. *Sport Med - Open.* 2015; Dec;1(1):7.
6. Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. Full symptomatic recovery does not ensure full recovery of muscle-tendon function in patients with Achilles tendinopathy. *Br J Sports Med.* 2007; 41(4):276-80.
7. Kannus P, Józsa L, Natri A, Järvinen M. Effects of training, immobilization and remobilization on tendons. *Scand J Med Sci Sports.* 2007;7(2):67-71.
8. Riley GP. Gene expression and matrix turnover in overused and damaged tendons. In: *Scandinavian Journal of Medicine and Science in Sports.* 2005;15(4):241-51.
9. Sarasa-Renedo A, Chiquet M. Mechanical signals regulating extracellular matrix gene expression in fibroblasts. In: *Scandinavian Journal of Medicine and Science in Sports.* 2005;15(4):223-30.
10. Moerch L, Pingel J, Boesen M, Kjaer M, Langberg H. The effect of acute exercise on collagen turnover in human tendons: Influence of prior immobilization period. *Eur J Appl Physiol.* 2013;113(2):449-455.
11. Christensen B, Dyrberg E, Aagaard P, Kjaer M, Langberg H. Short-term immobilization and recovery affect skeletal muscle but not collagen tissue turnover in humans. *J Appl Physiol.* 2008; 105(6):1845-51.
12. de Boer MD, Selby A, Atherton P, et al. The temporal responses of protein synthesis, gene expression and cell signalling in human quadriceps muscle and patellar tendon to disuse. *J Physiol.* 2007; 585(Pt 1):241-51.
13. Sun Y-L, Thoreson AR, Cha SS, Zhao C, An K-N, Amadio PC. Temporal response of canine flexor tendon to limb suspension. *J Appl Physiol.* 2010; 585(Pt 1):241-51.
14. Azevedo LB, Lambert MI, Vaughan CL, O'Connor CM, Schwellnus MP. Biomechanical variables associated with Achilles tendinopathy in runners. *Br J Sports Med.* 2009; 43(4):288-92.
15. Hoke B, Torrance A. Video Gait Analysis. In: O'Connor F, Wider R, eds. *Textbook of Running Medicine. 1st Edn.* New York: McGraw-Hill; 2001:59-66.
16. Malliaras P, Barton CJ, Reeves ND, Langberg H. Achilles and patellar tendinopathy loading programmes: A systematic review comparing clinical outcomes and identifying potential mechanisms for effectiveness. *Sport Med.* 2013; 43(4):267-86.
17. Silbernagel KG, Crossley KM. A proposed return-to-sport program for patients with midportion achilles tendinopathy: Rationale and implementation. *J Orthop Sports Phys Ther.* 2015; 45(11):876-86.
18. Grävare Silbernagel K, Brorsson A, Lundberg M. The majority of patients with Achilles tendinopathy recover fully when treated with exercise alone: A 5-year follow-up. *Am J Sports Med.* 2011; 39(3):607-13.
19. Van Der Plas A, De Jonge S, De Vos RJ, et al. A 5-year follow-up study of Alfredson's heel-drop exercise programme in chronic midportion Achilles tendinopathy. *Br J Sports Med.* 2012; 46(3):214-8.
20. Paavola M, Kannus P, Paakkala T, Pasanen M, Järvinen M. Long-term prognosis of patients with achilles tendinopathy: An observational 8-year follow-up study. *Am J Sports Med.* 2000; 28(5):634-42.
21. Johannsen F, Jensen S, Wetke E. 10-year follow-up after standardised treatment for Achilles tendinopathy. *BMJ Open Sport Exerc Med.* 2018; 4;4(1):e000415.
22. Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic achilles tendinosis. *Am J Sports Med.* 1998;26(3):360-6.
23. Maffulli N, Walley G, Sayana M, Longo UG, Denaro V. Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil.* 2008;30(20-22):1677-84.
24. Lorimer A V., Hume PA. Stiffness as a Risk Factor for Achilles Tendon Injury in Running Athletes. *Sport Med.* 2016; 46(12):1921-1938.
25. Stevens M, Tan C-W. Effectiveness of the Alfredson Protocol Compared With a Lower Repetition-Volume Protocol for Midportion Achilles Tendinopathy: A Randomized Controlled Trial. *J Orthop Sport Phys Ther.* 2014; 44(2):59-67.
26. Revak A, Diers K, Kernozek TW, Gheidi N, Olbrantz C. Achilles Tendon Loading During Heel-Raising and -Lowering Exercises. *J Athl Train.* 2017;52(2):89-96.
27. Mahieu NN, Witvrouw E, Stevens V, Van Tiggelen D, Roget P. Intrinsic risk factors for the development of Achilles tendon overuse injury: A prospective study. *Am J Sports Med.* 2006; 34(2):226-35.
28. McAuliffe S, Tabuena A, McCreesh K, et al. Altered Strength Profile in Achilles Tendinopathy: A Systematic Review and Meta-Analysis. *J Athl Train.* August 2019;1062-6050-43-18.

29. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surgery, Sport Traumatol Arthrosc.* 2001;9:42-47.
30. Niesen-Vertommen SL, Taunton JE, Clement DB, Mosher RE. The effect of eccentric versus concentric exercise in the management of achilles tendonitis. *Clin J Sport Med.* 2006;2:(2)
31. Roos EM, Engström M, Lagerquist A, Söderberg B. Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy - A randomized trial with 1-year follow-up. *Scand J Med Sci Sports.* 2004;14(5):286-95.
32. Silbernagel KG, Thomeé R, Thomeé P, Karlsson J. Eccentric overload training for patients with chronic Achilles tendon pain--a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sports.* 2001;11(4):197-206.
33. Van Der Vlist AC, Breda SJ, Oei EHG, Verhaar JAN, De Vos RJ. Clinical risk factors for Achilles tendinopathy: a systematic review. *Br J Sports Med.* 2019; pii: bjsports-2018-099991 [Epub ahead of print]
34. Owens BD, Wolf JM, Seelig AD, et al. Risk factors for lower extremity tendinopathies in military personnel. *Orthop J Sport Med.* 2013;11;1(1):2325967113492707
35. Weier AT, Pearce AJ, Kidgell DJ. Strength training reduces intracortical inhibition. *Acta Physiol.* 2012; 206(2):109-19.
36. Goodwill AM, Pearce AJ, Kidgell DJ. Corticomotor plasticity following unilateral strength training. *Muscle and Nerve.* 2012; 46(3):384-93.
37. Rantalainen T, Weier A, Leung M, Brandner C, Spittle M, Kidgell D. Short-interval intracortical inhibition is not affected by varying visual feedback in an isometric task in biceps brachii muscle. *Front Hum Neurosci.* 2013; 8;7:68
38. Bohm S, Mersmann F, Tettke M, Kraft M, Arampatzis A. Human Achilles tendon plasticity in response to cyclic strain: Effect of rate and duration. *J Exp Biol.* 2014; 15;217(Pt 22):4010-7.
39. Ackerley SJ, Stinear CM, Byblow WD. Promoting use-dependent plasticity with externally-paced training. *Clin Neurophysiol.* 2011;122(12):2462-8.
40. Kidgell DJ, Pearce AJ. Corticospinal properties following short-term strength training of an intrinsic hand muscle. *Hum Mov Sci.* 2010; 29(5):631-41..
41. Kidgell DJ, Stokes MA, Castricum TJ, Pearce AJ. Neurophysiological responses after short-term strength training of the biceps brachii muscle. *J Strength Cond Res.* 2010; 24(11):3123-32
42. Leung M, Rantalainen T, Teo WP, Kidgell D. Motor cortex excitability is not differentially modulated following skill and strength training. *Neuroscience.* 2015; 1;305:99-108.
43. Rees JD, Lichtwark GA, Wolman RL, Wilson AM. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an in vivo study in humans. *Rheumatology.* 2008;47(10):1493-7
44. Arya S, Kulig K. Tendinopathy alters mechanical and material properties of the Achilles tendon. *J Appl Physiol.* 2010;108(3):670-5
45. Wren TA, Yerby SA, Beaupre GS, Carter DR. Mechanical properties of the human achilles tendon. *Clin Biomech (Bristol, Avon).* 2001; 16(3):245-51.
46. Weiler JM. Medical modifiers of sports injury. The use of nonsteroidal anti- inflammatory drugs (NSAIDs) in sports soft-tissue injury. *Clin Sport Med.* 1992; 11(3):625-44.
47. Mosca MJ, Rashid MS, Snelling SJ, Kirtley S, Carr AJ, Dakin SG. Trends in the theory that inflammation plays a causal role in tendinopathy: A systematic review and quantitative analysis of published reviews. *BMJ Open Sport Exerc Med.* 2018;26;4(1):e000332.
48. Khan KM, Bonar F, Desmond PM, et al. Patellar tendinosis (jumper's knee): findings at histopathologic examination, US, and MR imaging. Victorian Institute of Sport Tendon Study Group. *Radiology.* 1996; 200(3):821-7.
49. Scott A, Khan KM, Cook JL, Duronio V. What is "inflammation"? Are we ready to move beyond Celsus? *Br J Sports Med.* 2004;38(3):248-249.
50. Rees JD, Stride M, Scott A. Tendons - Time to revisit inflammation. *Br J Sports Med.* 2014; 48(21):1553-7.
51. Kleinman M, Gross AE. Achilles tendon rupture following steroid injection. Report of three cases. *J Bone Jt Surg - Ser A.* 1983; 65(9):1345-7.
52. Ford LT, Debender J. Tendon rupture after local steroid injection. *South Med J.* 1979; 72(7):827-30.
53. Fredberg U, Bolvig L, Pfeiffer-Jensen M, Clemmensen D, Jakobsen BW, Stengaard-Pedersen K. Ultrasonography as a tool for diagnosis, guidance of local steroid injection and, together with pressure algometry, monitoring of the treatment of athletes with chronic jumper's knee and Achilles tendinitis: A randomized, double-blind, placebo-controlled study. *Scand J Rheumatol.* 2004.33(2):94-101.
54. O'Neill S, Watson P, Barry S. 75 Plantarflexor Muscle Power Deficits In Runners With Achilles Tendinopathy. *Br J Sports Med.* 2014;48:A49.
55. Toumi H, Lerguech G, Cherief M, et al. Implications of the calf musculature and Achilles tendon architectures for understanding the site of injury. *J Biomech.* 2016; 49(7):1180-1185.
56. Lieber RL, Ward SR. Skeletal muscle design to meet functional demands. *Philos Trans R Soc B Biol Sci.* 2011; 366(1570):1466-76
57. Dorn TW, Schache AG, Pandy MG. Muscular strategy shift in human running: Dependence of running speed on hip and ankle muscle performance. *J Exp Biol.* 2012; 215(Pt 11):1944-56.

58. Ishikawa M. Muscle-tendon interaction and elastic energy usage in human walking. *J Appl Physiol.* 2005; 99(2):603-8.
59. Szaro P, Witkowski G, Śmigielski R, Krajewski P, Ciszek B. Fascicles of the adult human Achilles tendon - An anatomical study. *Ann Anat.* 2009; (6):586-93.
60. Edama M, Kubo M, Onishi H, et al. The twisted structure of the human Achilles tendon. *Scand J Med Sci Sport.* 2015; 25(5):e497-503.
61. Cook JL, Rio E, Purdam CR, Docking SI. Revisiting the continuum model of tendon pathology: What is its merit in clinical practice and research? *Br J Sports Med.* 2016; 50(19):1187-91
62. Silbernagel KG, Nilsson-Helander K, Thomeé R, Eriksson BI, Karlsson J. A new measurement of heel-rise endurance with the ability to detect functional deficits in patients with Achilles tendon rupture. *Knee Surgery, Sport Traumatol Arthrosc.* 2010; 18(2):258-64.
63. Schache AG, Dorn TW, Williams GP, Brown NAT, Pandy MG. Lower-limb muscular strategies for increasing running speed. *J Orthop Sports Phys Ther.* 2014; 44(10):813-24
64. Farris DJ, Sawicki GS. Human medial gastrocnemius force-velocity behavior shifts with locomotion speed and gait. *Proc Natl Acad Sci U S A.* 2012;109(3):977-82.
65. Hof AL, Van Zandwijk JP, Bobbert MF. Mechanics of human triceps surae muscle in walking, running and jumping. *Acta Physiol Scand.* 2002;174(1):17-30.
66. Lai A, Schache AG, Lin YC, Pandy MG. Tendon elastic strain energy in the human ankle plantar-flexors and its role with increased running speed. *J Exp Biol.* 2014; 217(Pt 17):3159-68.
67. Sasaki K, Neptune RR. Muscle mechanical work and elastic energy utilization during walking and running near the preferred gait transition speed. *Gait Posture.* 2006; 23(3):383-90
68. Kalson NS, Holmes DF, Herchenhan A, Lu Y, Starborg T, Kadler KE. Slow stretching that mimics embryonic growth rate stimulates structural and mechanical development of tendon-like tissue in vitro. *Dev Dyn.* 2011;240(11):2520-2528.
69. Wang JHC. Mechanobiology of tendon. *J Biomech.* 2006;39(9):1563-1582.
70. Arnoczky SP, Lavagnino M, Egerbacher M. The mechanobiological aetiopathogenesis of tendinopathy: Is it the over-stimulation or the under-stimulation of tendon cells? In: *International Journal of Experimental Pathology.* Vol 88. 2007:217-226.
71. Dideriksen K. Muscle and tendon connective tissue adaptation to unloading, exercise and NSAID. *Connect Tissue Res.* 2014;55(2):61-70.
72. Yamamoto E, Hayashi K, Yamamoto N. Mechanical properties of collagen fascicles from stress-shielded patellar tendons in the rabbit. *Clin Biomech.* 1999;14(6):418-425.
73. Couppé C, Svensson RB, Silbernagel KG, Langberg H, Magnusson SP. Eccentric or Concentric Exercises for the Treatment of Tendinopathies? *J Orthop Sport Phys Ther.* 2015;45(11):1-25.
74. Langberg H, Olesen J, Skovgaard D, Kjaer M. Age related blood flow around the Achilles tendon during exercise in humans. *Eur J Appl Physiol.* 2001;84:246-248.
75. Langberg H, Skovgaard D, Petersen LJ, Bulow J, Kjaer M. Type I collagen synthesis and degradation in peritendinous tissue after exercise determined by microdialysis in humans. *J Physiol.* 1999;521 Pt 1:299-306.
76. Couppé C, Kongsgaard M, Aagaard P, et al. Habitual loading results in tendon hypertrophy and increased stiffness of the human patellar tendon. *J Appl Physiol.* 2008;105:805-810.
77. Shalabi A, Kristoffersen-Wilberg M, Svensson L, Aspelin P, Movin T. Eccentric training of the gastrocnemius-soleus complex in chronic Achilles tendinopathy results in decreased tendon volume and intratendinous signal as evaluated by MRI. *Am J Sports Med.* 2004;32(5):1286-1296.
78. Woo SL, Ritter M a, Amiel D, et al. The biomechanical and biochemical properties of swine tendons--long term effects of exercise on the digital extensors. *Connect Tissue Res.* 1980;7(3):177-183.
79. Milgrom Y, Milgrom C, Altaras T, Globus O, Zeltzer E, Finestone AS. Achilles tendons hypertrophy in response to high loading training. *Foot Ankle Int.* 2014; 35(12):1303-8.
80. Chimenti RL, Flemister AS, Ketz J, Bucklin M, Buckley MR, Richards MS. Ultrasound strain mapping of Achilles tendon compressive strain patterns during dorsiflexion. *J Biomech.* 2016; 49(1):39-44
81. Fahlstrom M, Jonsson P, Lorentzon R, Alfredson H. Chronic Achilles tendon pain treated with eccentric calf-muscle training. *Knee Surgery, Sport Traumatol Arthrosc.* 2004; 11(5):327-33.
82. Jonsson P, Alfredson H, Sunding K, Fahlström M, Cook J. New regimen for eccentric calf-muscle training in patients with chronic insertional Achilles tendinopathy: Results of a pilot study. *Br J Sports Med.* 2008; 42(9):746-9
83. Gajhede-Knudsen M, Ekstrand J, Magnusson H, Maffulli N. Recurrence of Achilles tendon injuries in elite male football players is more common after early return to play: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med.* 2013;47(12):763-768.
84. Hagglund M, Walden M, Ekstrand J. Lower Reinjury Rate With a Coach-Controlled Rehabilitation Program in Amateur Male Soccer: A Randomized Controlled Trial. *Am J Sports Med.* 2007;35(9):1433-1442.
85. Mohammadi F. A coach-controlled rehabilitation program reduces the risk of reinjury among amateur soccer players. *Aust J Physiother.* 2008;54(1):74.

86. Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med.* 2007;35(6):897-906.
87. Newman JS, Adler RS, Bude RO, Rubin JM. Detection of soft-tissue hyperemia: Value of power Doppler sonography. *Am J Roentgenol.* 1994;163(2):385-389.
88. Sharma P, Maffulli N. Basic biology of tendon injury and healing. *Surg.* 2005;3(5):309-316.
89. Alfredson H, Ohberg L, Forsgren S. Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc.* 2003;11(5):334-338.
90. Tol JL, Spiezia F, Maffulli N. Neovascularization in Achilles tendinopathy: Have we been chasing a red herring? *Knee Surgery, Sport Traumatol Arthrosc.* 2012;20(10):1891-1894.
91. Alfredson H, Ohberg L. Sclerosing injections to areas of neo-vascularisation reduce pain in chronic Achilles tendinopathy: a double-blind randomised controlled trial. *Knee Surg Sport Traumatol Arthrosc.* 2005;13(4):338-344.
92. Lind B, Öhberg L, Alfredson H. Sclerosing polidocanol injections in mid-portion Achilles tendinosis: Remaining good clinical results and decreased tendon thickness at 2-year follow-up. *Knee Surgery, Sport Traumatol Arthrosc.* 2006;14(12):1327-1332.
93. Ohberg L, Alfredson H. Ultrasound guided sclerosis of neovessels in painful chronic Achilles tendinosis: pilot study of a new treatment. *Br J Sports Med.* 2002;36(3):173-177.
94. Öhberg L, Lorentzon R, Alfredson H. Neovascularisation in Achilles tendons with painful tendinosis but not in normal tendons: An ultrasonographic investigation. *Knee Surgery, Sport Traumatol Arthrosc.* 2001;9(4):233-238.
95. de Vos R-J, Weir A, Cobben LPJ, Tol JL. The value of power Doppler ultrasonography in Achilles tendinopathy: a prospective study. *Am J Sports Med.* 2007;35(10):1696-1701.
96. Peers KHE, Brys PPM, Lysens RJJ. Correlation between power Doppler ultrasonography and clinical severity in Achilles tendinopathy. *Int Orthop.* 2003;27(3):180-183.
97. Reiter M, Ulreich N, Dirisamer A, Tscholakoff D, Bocek RA. Colour and power doppler sonography in symptomatic achilles tendon disease. *Int J Sports Med.* 2004;25(4):301-305.
98. Zanetti M, Metzdorf A, Kundert H-P, et al. Achilles Tendons: Clinical Relevance of Neovascularization Diagnosed with Power Doppler US1. *Radiology.* 2003;227:556-560.
99. Sengkerij PM, de Vos R-J, Weir A, van Weelde BJG, Tol JL. Interobserver reliability of neovascularization score using power Doppler ultrasonography in midportion achilles tendinopathy. *Am J Sports Med.* 2009;37(8):1627-1631.
100. Boesen MI, Boesen A, Koenig MJ, Bliddal H, Torp-Pedersen S. Ultrasonographic investigation of the Achilles tendon in elite badminton players using color Doppler. *Am J Sports Med.* 2006;34(12):2013-2021.
101. De Jonge S, Warnars JLF, De Vos RJ, et al. Relationship between neovascularization and clinical severity in Achilles tendinopathy in 556 paired measurements. *Scand J Med Sci Sport.* 2014;24(5):773-778.
102. Van Snellenberg W, Wiley JP, Brunet G. Achilles tendon pain intensity and level of neovascularization in athletes as determined by color Doppler ultrasound. *Scand J Med Sci Sport.* 2007;17(5):530-534.
103. van Sterkenburg MN, de Jonge MC, Sierevelt IN, van Dijk CN. Less promising results with sclerosing ethoxysclerol injections for midportion achilles tendinopathy: a retrospective study. *Am J Sports Med.* 2010;38(11):2226-2232.
104. Witchalls C. High Cholesterol Linked to Tendinopathy. *Medscape Fam Med.* 2015.
105. B. Tilley, J. Gaida, J. Cook, J. Docking. Is higher serum cholesterol associated with altered Achilles tendon structure or Achilles tendon pain in the general population? *J Sci Med Sport.* 2015; 49:1504-1509.
106. Murtaugh B, Ihm JM. Eccentric training for the treatment of tendinopathies. *Curr Sports Med Rep.* 2013; 12(3):175-82
107. Robinson JM, Cook JL, Purdam C, et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med.* 2001; 35(5):335-41.
108. Rompe JD, Nafe B, Furia JP, Maffulli N. Eccentric loading, shock-wave treatment, or a wait- and-see policy for tendinopathy of the main body of tendo Achillis: A randomized controlled trial. *Am J Sports Med.* 2007; 35(3):374-83.
109. Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med.* 1998;26(3):360-366.
110. Bohm S, Mersmann F, Tettke M, Kraft M, Arampatzis A. Human Achilles tendon plasticity in response to cyclic strain: effect of rate and duration. *J Exp Biol.* 2014;217(Pt 22):4010-4017.
111. Arampatzis A, Peper A, Bierbaum S, Albracht K. Plasticity of human Achilles tendon mechanical and morphological properties in response to cyclic strain. *J Biomech.* 2010;43(16):3073-3079.
112. Rees JD, Lichtwark G a, Wolman RL, Wilson a M. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an in vivo study in humans. *Rheumatology (Oxford).* 2008;47(10):1493-1497.
113. Gärdin A, Movin T, Svensson L, Shalabi A. The long-term clinical and MRI results following eccentric calf muscle training in chronic Achilles tendinosis. *Skeletal Radiol.* 2010; 39(5):435-42

114. Reid D, McNair PJ, Johnson S, Potts G, Witvrouw E, Mahieu N. Electromyographic analysis of an eccentric calf muscle exercise in persons with and without Achilles tendinopathy. *Phys Ther Sport*. 2012;13(3):150-5.
115. Synott A, O'Sullivan K, Purtill H, Mc Creesh K, Casey H, Mc Auliffe S. Beyond the tendon: Experiences and perceptions of people with persistent Achilles tendinopathy. *Musculoskelet Sci Pract*. 2017; 29:108-114.
116. Meyer A, Tumilty S, Baxter GD. Eccentric exercise protocols for chronic non-insertional Achilles tendinopathy: How much is enough? *Scand J Med Sci Sport*. 2009; 19(5):609-15
117. Miles CL, Pincus T, Carnes D, et al. Can we identify how programmes aimed at promoting self-management in musculoskeletal pain work and who benefits? A systematic review of sub-group analysis within RCTs. *Eur J Pain*. 2011; 15(8):775.e1-11.
118. Herrington L, McCulloch R. The role of eccentric training in the management of Achilles tendinopathy: A pilot study. *Phys Ther Sport*. 2007.8(4)191-196
119. Nørregaard J, Larsen CC, Bieler T, Langberg H. Eccentric exercise in treatment of Achilles tendinopathy. *Scand J Med Sci Sport*. 2007.17(2):133-8.
120. Stanish WD, Rubinovich RM, Curwin S. Eccentric exercise in chronic tendinitis. *Clin Orthop Relat Res*. 1986;(208):65-68.
121. Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med*. 1998;26(3):360-366.
122. Dimitrios S, Malliaras P. It is time to abandon the myth that eccentric training is best practice. *Biol Exerc*. 2016;12(1):15-21.
123. Proske U, Morgan DL. Muscle damage from eccentric exercise: Mechanism, mechanical signs, adaptation and clinical applications. *J Physiol*. 2001;537(2):333-345.
124. Morgan DL, Allen DG. Early events in stretch-induced muscle damage. *J Appl Physiol*. 1999;87(6):2007-15.
125. Smith LL. Acute inflammation: The underlying mechanism in delayed onset muscle soreness? *Med Sci Sports Exerc*. 1991; 23(5):542-51.
126. MacIntyre DL, Reid WD, McKenzie DC. Delayed Muscle Soreness: The Inflammatory Response to Muscle Injury and its Clinical Implications. *Sport Med*. 1995.20(1):24-40.
127. Kingma JJ, de Knikker R, Wittink HM, Takken T. Eccentric overload training in patients with chronic Achilles tendinopathy: a systematic review. *Br J Sports Med*. 2006. 41(6):e3. Epub 2006 Oct 11.
128. Wasielewski NJ, Kotsko KM. Does eccentric exercise reduce pain and improve strength in physically active adults with symptomatic lower extremity tendinosis? A systematic review. *J Athl Train*. 2007; 42(3):409-21.
129. Woodley BL, Newsham-West RJ, Baxter GD. Chronic tendinopathy: Effectiveness of eccentric exercise. *Br J Sports Med*. 2007;41(4):188-98
130. Satyendra L, Byl N. Effectiveness of physical therapy for Achilles tendinopathy: An evidence based review of eccentric exercises. *Isokinet Exerc Sci*. 2018;14(1): 71-80.
131. Chaudhry S, Morrissey D, Woledge RC, Bader DL, Screen HRC. Eccentric and concentric loading of the triceps surae: an in vivo study of dynamic muscle and tendon biomechanical parameters. *J Appl Biomech*. 2015;31(2):69-78.
132. Öhberg L, Lorentzon R, Alfredson H. Eccentric training in patients with chronic Achilles tendinosis: Normalised tendon structure and decreased thickness at follow up. *Br J Sports Med*. 2004;38(1):8-11.
133. Drew BT, Smith TO, Littlewood C, Sturrock B. Do structural changes (eg, collagen/matrix) explain the response to therapeutic exercises in tendinopathy: a systematic review. *Br J Sports Med*. 2014;48(12):966-972.
134. Fouré A, Nordez A, Cornu C. Plyometric training effects on Achilles tendon stiffness and dissipative properties. *J Appl Physiol*. 2010.109(3):849-54.
135. Gravare Silbernagel K, Thomee R, Thomee P, Karlsson J. Eccentric overload training for patients with chronic Achilles tendon pain - a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sport*. 2001.
136. Balsalobre-Fernández C, Glaister M, Lockey RA. The validity and reliability of an iPhone app for measuring vertical jump performance. *J Sports Sci*. 2015; 33(15):1574-9.