

## REFERENCES

1. Robertson MC, Devlin N, Gardner MM, Campbell AJ. Effectiveness and economic evaluation of a nurse delivered home exercise programme to prevent falls. 1: Randomised controlled trial. *BMJ*. 2001;322(7288):697-701.
2. Rossiter-Fornoff JE, Wolf SL, Wolfson LI, et al. A cross-sectional validation study of the FICSIT common data base static balance measures. *Journals Gerontol - Ser A Biol Sci Med Sci*. 1995;50A(6):M291-M297.
3. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport*. 1999;70(2):113-119.
4. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991;39(2):142-148.
5. Power V, Clifford AM. Characteristics of optimum falls prevention exercise programmes for community-dwelling older adults using the FITT principle. *Eur Rev Aging Phys Act*. 2013;10(2):95-106.
6. Reinsch S, Mac PR, Lachenbruch PA, Tobis JS. Attempts to Prevent Falls and Injury: A Prospective Community Study. *Gerontologist*. 1992;32(4):450-456.
7. Kyrdalen IL, Moen K, Roysland AS, Helbostad JL. The Otago exercise program performed as group training versus home training in fall-prone older people: A randomized controlled trial. *Physiother Res Int*. 2014;19(2):108-116.
8. Latham NK, Anderson CS, Lee A, et al. A randomized, controlled trial of quadriceps resistance exercise and vitamin D in frail older people: the Frailty Interventions Trial in Elderly Subjects (FITNESS). *J Am Geriatr Soc*. 2003;51(3):291-299.
9. Campbell AJ, Robertson MC, Gardner MM, Norton RN, Tilyard MW, Buchner DM. Randomised controlled trial of a general practice programme of home based exercise to prevent falls in elderly women. *BMJ*. 1997;315(7115):1065-1069.
10. Campbell A, Robertson M. *Otago Exercise Programme. To Prevent Falls in Older Adults*. Otago Medical School, University of Otago, Otago, New Zealand March 2003.
11. Sherrington C, Whitney JC, Lord SR, Herbert RD, Cumming RG, Close JCT. Effective exercise for the prevention of falls: A systematic review and meta-analysis. *J Am Geriatr Soc*. 2008;56(12):2234-2243.
12. Sherrington C, Lord SR, Close JCT. *Best-Practice Recommendations for Physical Activity to Prevent Falls in Older Adults*. The George Institute for International Health, Prince of Wales Medical Research Institute, Prince of Wales Hospital, The Sax Institute Sydney; November 2008. pp 1-36
13. Benavent-Caballer V, Rosado-Calatayud P, Segura-Ortí E, Amer-Cuenca JJ, Lisón JF. The effectiveness of a video-supported group-based Otago exercise programme on physical performance in community-dwelling older adults: a preliminary study. *Physiotherapy*. 2015. [Epub ahead of print]
14. Son N-K, Ryu YU, Jeong H-W, Jang Y-H, Kim H-D. Comparison of 2 Different Exercise Approaches: Tai Chi Versus Otago, in Community-Dwelling Older Women. *J Geriatr Phys Ther*. 2016;39(2):51-7
15. Thomas S, Mackintosh S, Halbert J. Does the "Otago exercise programme" reduce mortality and falls in older adults?: A systematic review and meta-analysis. *Age Ageing*. 2010;39(6):681-687.
16. Gillespie LD, Robertson MC, Gillespie WJ, et al. Interventions for preventing falls in older people living in the community. *Cochrane database Syst Rev*. 2012;9(9):CD007146.
17. American Geriatrics Society, Society BG, Prevention AA of OSP on F. Guideline for the prevention of falls in older persons. *J Am Geriatr Soc*. 2001;49:664-672.
18. Deandrea S, Lucenteforte E, Bravi F, Foschi R, La Vecchia C, Negri E. Risk Factors for Falls in Community-dwelling Older People. *Epidemiology*. 2010;21(5):658-668.
19. Cauley JA, Cawthon PM, Peters KE, et al. Risk Factors for Hip Fracture in Older Men: The Osteoporotic Fractures in Men Study (MrOS). *J Bone Miner Res*. 2016. [Epub ahead of print]
20. Stubbs B, Binnekade T, Eggermont L, Sepehry AA, Patchay S, Schofield P. Pain and the risk for falls in community-dwelling older adults: Systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2014;95(1).
21. Spink MJ, Menz HB, Fotoohabadi MR, et al. Effectiveness of a multifaceted podiatry intervention to prevent falls in community dwelling older people with disabling foot pain: randomised controlled trial. *BMJ*. 2011;342(26):d3411.
22. Menz HB, Morris ME, Lord SR. Foot and Ankle Risk Factors for Falls in Older People: A Prospective Study. *Journals Gerontol Ser A Biol Sci Med Sci*. 2006;61(8):866-870.
23. Menz HB, Morris ME, Lord SR. Foot and ankle characteristics associated with impaired balance and functional ability in older people. *J Gerontol A Biol Sci Med Sci*. 2005;60(12):1546-1552.
24. Crews JE, Chou C-F, Stevens JA, Saaddine JB. Falls Among Persons Aged ≥65 Years With and Without Severe Vision Impairment - United States, 2014. *MMWR Morb Mortal Wkly Rep*. 2016;65(17):433-437.
25. Foss AJE, Harwood RH, Osborn F, Gregson RM, Zaman A, Masud T. Falls and health status in elderly women following second eye cataract surgery: A randomised controlled trial. *Age Ageing*. 2006;35(1):66-71.
26. Pahwa AK, Andy UU, Newman DK, Stambakio H, Schmitz KH, Arya LA. Nocturnal Enuresis as a Risk Factor for Falls in Older Community Dwelling Women with Urinary Incontinence. *J Urol*. 2016.195(5):1512-6.

27. Ekvall Hansson E, Magnusson M. Vestibular asymmetry predicts falls among elderly patients with multi-sensory dizziness. *BMC Geriatr.* 2013;13:77.
28. Campbell AJ, Robertson MC, Gardner MM, Norton RN, Buchner DM. Psychotropic medication withdrawal and a home-based exercise program to prevent falls: a randomized, controlled trial. *J Am Geriatr Soc.* 1999;47(7):850-853.
29. Marcum ZA, Perera S, Thorpe JM, et al. Antidepressant Use and Recurrent Falls in Community-Dwelling Older Adults: Findings From the Health ABC Study. *Ann Pharmacother.* 2016;50(7):525-533.
30. Hanlon JT, Boudreau RM, Roumani YF, et al. Number and dosage of central nervous system medications on recurrent falls in community elders: The health, aging and body composition study. *Journals Gerontol - Ser A Biol Sci Med Sci.* 2009;64(4):492-498.
31. Callisaya ML, Sharman JE, Close J, Lord SR, Srikanth VK. Greater daily defined dose of antihypertensive medication increases the risk of falls in older people--a population-based study. *J Am Geriatr Soc.* 2014;62(8):1527-1533.
32. Marcum ZA, Perera S, Newman AB, et al. Antihypertensive use and recurrent falls in community-dwelling older adults: Findings from the health ABC study. *Journals Gerontol - Ser A Biol Sci Med Sci.* 2015;70(12):1562-1568.
33. Leipzig RM, Cumming RG, Tinetti ME. Drugs and falls in older people: a systematic review and meta-analysis: II. Cardiac and analgesic drugs. *J Am Geriatr Soc.* 1999;47(1):40-50.
34. Bischoff-Ferrari HA, Orav EJ, Dawson-Hughes B. Effect of cholecalciferol plus calcium on falling in ambulatory older men and women: a 3-year randomized controlled trial. *Arch Intern Med.* 2006;166(4):424-430.
35. Harwood RH, Sahota O, Gaynor K, Masud T, Hosking DJ. A randomised, controlled comparison of different calcium and vitamin D supplementation regimens in elderly women after hip fracture: The Nottingham Neck of Femur (NoNOF) study. *Age Ageing.* 2004;33(1):45-51.
36. Kvelde T, McVeigh C, Toson B, et al. Depressive symptomatology as a risk factor for falls in older people: Systematic review and meta-analysis. *J Am Geriatr Soc.* 2013;61(5):694-706.
37. Haran MJ, Cameron ID, Ivers RQ, et al. Effect on falls of providing single lens distance vision glasses to multifocal glasses wearers: VISIBLE randomised controlled trial. *BMJ.* 2010;340:c2265.
38. Kenny R a, Richardson D a, Steen N, Bexton RS, Shaw FE, Bond J. Carotid sinus syndrome: a modifiable risk factor for nonaccidental falls in older adults (SAFE PACE). *J Am Coll Cardiol.* 2001;38(5):1491-1496.
39. Menant JC, Wong AKW, Trollor JN, Close JCT, Lord SR. Depressive Symptoms and Orthostatic Hypotension Are Risk Factors for Unexplained Falls in Community-Living Older People. *J Am Geriatr Soc.* 2016;64(5):1073-1078.
40. Gangavati A, Hajjar I, Quach L, et al. Hypertension, orthostatic hypotension, and the risk of falls in a community-dwelling elderly population: The maintenance of balance, independent living, intellect, and zest in the elderly of Boston study. *J Am Geriatr Soc.* 2011;59(3):383-389.
41. Kuhlman KA. Cervical range of motion in the elderly. *Arch Phys Med Rehabil.* 1993;74(10):1071-1079.
42. Fiebert IM, Armstrong T, Caliendo F, Herstick M, Roach K. Cervical Range of Motion in Geriatric Individuals. *Phys Occup Ther Geriatr.* 2009;12(3):65-83.
43. Liu B, Wu B, Van Hoof T, Okito J-PK, Liu Z, Zeng Z. Are the standard parameters of cervical spine alignment and range of motion related to age, sex, and cervical disc degeneration? *J Neurosurg Spine.* 2015;23(3):274-279.
44. Dasgupta S, Das S, Chaudhuri A. Carotid sinus hypersensitivity: Entity warrants a caution in the critical care unit. *J Sci Soc.* 2016;43(2):82.
45. Poole E, Treleaven J, Jull G. The influence of neck pain on balance and gait parameters in community-dwelling elders. *Man Ther.* 2008;13(4):317-324.
46. Uthairkhum S, Jull G, Sungkarat S, Treleaven J. The influence of neck pain on sensorimotor function in the elderly. *Arch Gerontol Geriatr.* 2012;55(3):667-672.
47. Quek JM, Pua YH, Bryant AL, Clark RA. The influence of cervical spine flexion-rotation range-of-motion asymmetry on postural stability in older adults. *Spine (Phila Pa 1976).* 2013;38(19):1648-1655.
48. Karlberg M, Magnusson M, Johansson R. Effects of restrained cervical mobility on voluntary eye movements and postural control. *Acta Otolaryngol.* 1991;111(4):664-670.
49. Kelders WP a, Kleinrensink GJ, van der Geest JN, et al. The cervico-ocular reflex is increased in whiplash injury patients. *J Neurotrauma.* 2005;22(1):133-137.
50. Akeson WH, Amiel D, Woo SL. Immobility effects on synovial joints the pathomechanics of joint contracture. *Biorheology.* 1980;17(1-2):95-110.
51. Donatelli R, Owens-Burkhart H. Effects of Immobilization on the Extensibility of Periarticular Connective Tissue. *J Orthop Sport Phys Ther.* 1981;3(2):67-72.
52. Gross J. Collagen. *Sci Am.* 1961;204:121-130.
53. Kelders WP a, Kleinrensink GJ, van der Geest JN, Feenstra L, de Zeeuw CI, Frens M a. Compensatory increase of the cervico-ocular reflex with age in healthy humans. *J Physiol.* 2003;553(Pt 1):311-317.
54. Paige GD. Senescence of human visual-vestibular interactions: smooth pursuit, optokinetic, and vestibular control of eye movements with aging. *Exp Brain Res.* 1994;98(2):355-372.
55. Huygen PL, Verhagen WI, Nicolaisen MG. Cervico-ocular reflex enhancement in labyrinthine-defective and normal subjects. *Exp brain Res.* 1991;87(2):457-464.

56. De-la-Llave-Rincón AI, Fernández-de-las-Peñas C, Palacios-Ceña D, Cleland J a. Increased forward head posture and restricted cervical range of motion in patients with carpal tunnel syndrome. *J Orthop Sports Phys Ther.* 2009;39(9):658-664.
57. Kang JH, Park RY, Lee SJ, Kim JY, Yoon SR, Jung KI. The effect of the forward head posture on postural balance in long time computer based worker. *Ann Rehabil Med.* 2012;36(1):98-104.
58. Silva AG, Johnson MI. Does forward head posture affect postural control in human healthy volunteers? *Gait Posture.* 2013;38(2):352-353.
59. Asmita K, Cresida DS, Anuprita T, Twinkle D, Unnati P, Sujata Y. Effects of Forward Head Posture on Balance in Asymptomatic Young Adults. *Indian J Public Heal Res Dev.* 2015;6(3):123-126.
60. Ohmure H, Miyawaki S, Nagata J, Ikeda K, Yamasaki K, Al-Kalaly A. Influence of forward head posture on condylar position. *J Oral Rehabil.* 2008;35(11):795-800.
61. Weon JH, Oh JS, Cynn HS, Kim YW, Kwon OY, Yi CH. Influence of forward head posture on scapular upward rotators during isometric shoulder flexion. *J Bodyw Mov Ther.* 2010;14(4):367-374.
62. Fon GJ, Pitt MJ, Thies AC. Thoracic kyphosis: Range in normal subjects. *Am J Roentgenol.* 1980;134(5):979-983.
63. Granito RN, Aveiro MC, Rennó ACM, Oishi J, Driusso P. Degree of thoracic kyphosis and peak torque of trunk flexors and extensors among healthy women. *Rev Bras Ortop (English Ed.* 2014;49(3):286-291.
64. Lau KT, Cheung KY, Chan kwok B, Chan MH, Lo KY, Wing Chiu TT. Relationships between sagittal postures of thoracic and cervical spine, presence of neck pain, neck pain severity and disability. *Man Ther.* 2010;15(5):457-462.
65. Quek J, Pua YH, Clark RA, Bryant AL. Effects of thoracic kyphosis and forward head posture on cervical range of motion in older adults. *Man Ther.* 2013;18:65-71.
66. Raine S, Twomey LT. Head and shoulder posture variations in 160 asymptomatic women and men. *Arch Phys Med Rehabil.* 1997;78(11):1215-1223.
67. Caneiro JP, O'Sullivan P, Burnett A, et al. The influence of different sitting postures on head/neck posture and muscle activity. *Man Ther.* 2010;15(1):54-60.
68. Nejati P, Lotfian S, Moezy A, Nejati M. The relationship of forward head posture and rounded shoulders with neck pain in Iranian office workers. *Med J Islam Repub Iran.* 2014;28:26.
69. Yip CHT, Chiu TTW, Poon ATK. The relationship between head posture and severity and disability of patients with neck pain. *Man Ther.* 2008;13(2):148-154.
70. De Groot MH, van der Jagt-Willems HC, van Campen JPCM, Lems WF, Beijnen JH, Lamoth CJC. A flexed posture in elderly patients is associated with impairments in postural control during walking. *Gait Posture.* 2014;39(2):767-772.
71. McLean L. The effect of postural correction on muscle activation amplitudes recorded from the cervicobrachial region. *J Electromyogr Kinesiol.* 2005;15(6):527-535.
72. Enwemeka CS, Bonet IM, Ingle J a, Prudhithumrong S, Ogbahon FE, Gbenedio N a. Postural Correction in Persons with Neck Pain (II. Integrated Electromyography of the Upper Trapezius in Three Simulated Neck Positions). *J Orthop Sports Phys Ther.* 1986;8(5):240-242.
73. Pearson ND, Walmsley RP. Trial into the effects of repeated neck retractions in normal subjects. *Spine (Phila Pa 1976).* 1995;20(11):1245-1251.
74. Abdulwahab SS, Sabbahi M. Neck retractions, cervical root decompression, and radicular pain. *J Orthop Sports Phys Ther.* 2000;30(1):4-9; discussion 10-12.
75. Silva AG, Punt TD, Sharples P, Vilas-boas JP, Johnson MI. Head posture assessment for patients with neck pain: Is it useful? *Int J Ther Rehabil.* 2009;16(1):43-53.
76. Canales JZ, Cordás TA, Fiquer JT, Cavalcante AF, Moreno RA. Posture and body image in individuals with major depressive disorder: A controlled study. *Rev Bras Psiquiatr.* 2010;32(4):375-380.
77. Tonosu J, Matsudaira K, Oka H, et al. A population approach to analyze the effectiveness of a back extension exercise "One Stretch" in patients with low back pain: A replication study. *J Orthop Sci.* 2016;21(4):414-8.
78. Matsudaira K, Hiroe M, Kikkawa M, et al. Can standing back extension exercise improve or prevent low back pain in Japanese care workers? *J Man Manip Ther.* 2015;23(4):205-209.
79. Dettori JR, Bullock SH, Sutlive TG, Franklin RJ, Patience T. The effects of spinal flexion and extension exercises and their associated postures in patients with acute low back pain. *Spine (Phila Pa 1976).* 1995;20(21):2303-2312.
80. Elnaggar IM, Nordin M, Sheikhzadeh A, Parnianpour M, Kahanovitz N. Effects of spinal flexion and extension exercises on low-back pain and spinal mobility in chronic mechanical low-back pain patients. *Spine (Phila Pa 1976).* 1991;16(8):967-972.
81. Stankovic R, Johnell O. Conservative Treatment of Acute Low-Back Pain: A Prospective Randomized Trial: McKenzie Method of Treatment Versus Patient Education in "Mini Back School." *Spine (Phila Pa 1976).* 1990;15(2):120-123.
82. Halliday MH, Pappas E, Hancock MJ, et al. A Randomized Controlled Trial Comparing the McKenzie Method to Motor Control Exercises in People With Chronic Low Back Pain and a Directional Preference. *J Orthop Sports Phys Ther.* 2016:1-30.
83. Machado LA, de Souza M, Ferreira PH, Ferreira ML. The McKenzie method for low back pain: a systematic review of the literature with a meta-analysis approach. *Spine (Phila Pa 1976).* 2006;31(9):E254-E262.
84. Machado LA, Maher CG, Herbert RD, Clare H, McAuley JH. The effectiveness of the McKenzie method in addition to first-line care for acute low back pain: a randomized controlled trial. *BMC Med.* 2010;8:10.

85. Gondhalekar GA, Kumar SP, Eapen C, Mahale A. Reliability and Validity of Standing Back Extension Test for Detecting Motor Control Impairment in Subjects with Low Back Pain. *J Clin Diagn Res.* 2016;10(1):KC07-KC11.
86. Murphy DR, Byfield D, McCarthy P, Humphreys K, Gregory AA, Rochon R. Interexaminer Reliability of the Hip Extension Test for Suspected Impaired Motor Control of the Lumbar Spine. *J Manipulative Physiol Ther.* 2006;29(5):374-377.
87. Edmondston SJ, Aggerholm M, Elfving S, et al. Influence of Posture on the Range of Axial Rotation and Coupled Lateral Flexion of the Thoracic Spine. *J Manipulative Physiol Ther.* 2007;30(3):193-199.
88. McGill SM, Yingling VR, Peach JP. Three-dimensional kinematics and trunk muscle myoelectric activity in the elderly spine - a database compared to young people. *Clin Biomech (Bristol, Avon).* 1999;14(6):389-395.
89. Simoneau M, Guillaud É, Blouin J. Effects of underestimating the kinematics of trunk rotation on simultaneous reaching movements: predictions of a biomechanical model. *J Neuroeng Rehabil.* 2013;10:54.
90. Nagai K, Tateuchi H, Takashima S, et al. Effects of trunk rotation on scapular kinematics and muscle activity during humeral elevation. *J Electromyogr Kinesiol.* 2013;23(3):679-687.
91. Marras WS, Lavender SA, Leurgans SE, et al. The role of dynamic three-dimensional trunk motion in occupationally-related low back disorders. The effects of workplace factors, trunk position, and trunk motion characteristics on risk of injury. *Spine (Phila Pa 1976).* 1993;18(5):617-628.
92. Shan X, Ning X, Chen Z, Ding M, Shi W, Yang S. Low back pain development response to sustained trunk axial twisting. *Eur Spine J.* 2013;22(9):1972-1978.
93. Spink MJ, Fotoohabadi MR, Wee E, Hill KD, Lord SR, Menz HB. Foot and ankle strength, range of motion, posture, and deformity are associated with balance and functional ability in older adults. *Arch Phys Med Rehabil.* 2011;92(1):68-75.
94. Mecagni C, Smith JP, Roberts KE, O'Sullivan SB. Balance and ankle range of motion in community-dwelling women aged 64 to 87 years: a correlational study. *Phys Ther.* 2000;80(1):1004-1011.
95. Gehlsen GM, Whaley MH. Falls in the elderly: Part II, Balance, strength, and flexibility. *Arch Phys Med Rehabil.* 1990;71(10):739-741.
96. Nitz JC, Choy NL. The relationship between ankle dorsiflexion range, falls and activity level in women aged 40 to 80 years. *NZ J Physiother.* 2004;121-125.
97. Maki BE, Perry SD, Norrie RG, McIlroy WE. Effect of facilitation of sensation from plantar foot-surface boundaries on postural stabilization in young and older adults. *J Gerontol.* 1999;54(6):M281-M287.
98. Nurse MA, Nigg BM. Quantifying a relationship between tactile and vibration sensitivity of the human foot with plantar pressure distributions during gait. *Clin Biomech.* 1999;14(9):667-672.
99. Perry SD, McIlroy WE, Maki BE. The role of plantar cutaneous mechanoreceptors in the control of compensatory stepping reactions evoked by unpredictable, multi-directional perturbation. *Brain Res.* 2000;877(2):401-406.
100. Perry SD, Santos LC, Patla AE. Contribution of vision and cutaneous sensation to the control of centre of mass (COM) during gait termination. *Brain Res.* 2001;913(1):27-34.
101. Kavounoudias A, Roll R, Roll JP. Foot sole and ankle muscle inputs contribute jointly to human erect posture regulation. *J Physiol.* 2001;532(3):869-878.
102. Palluel E, Nougier V, Olivier I. Do spike insoles enhance postural stability and plantar-surface cutaneous sensitivity in the elderly? *Age (Omaha).* 2008;30(1):53-61.
103. Eils E, Nolte S, Tewes M, Thorwesten L, Völker K, Rosenbaum D. Modified pressure distribution patterns in walking following reduction of plantar sensation. *J Biomech.* 2002;35(10):1307-1313.
104. Eils E, Behrens S, Mers O, Thorwesten L, Völker K, Rosenbaum D. Reduced plantar sensation causes a cautious walking pattern. *Gait Posture.* 2004;20(1):54-60.
105. Do MC, Bussel B, Breniere Y. Influence of plantar cutaneous afferents on early compensatory reactions to forward fall. *Exp Brain Res.* 1990;79(2):319-324.
106. Horak FB, Nashner LM, Diener HC. Postural strategies associated with somatosensory and vestibular loss. *Exp Brain Res.* 1990;82(1):167-177.
107. Meyer PF, Oddsson LIE, De Luca CJ. The role of plantar cutaneous sensation in unperturbed stance. *Exp Brain Res.* 2004;156(4):505-512.
108. Meyer PF, Oddsson LIE, De Luca CJ. Reduced plantar sensitivity alters postural responses to lateral perturbations of balance. *Exp Brain Res.* 2004;157(4):526-536.
109. Diener HC, Dichgans J, Guschlbauer B, Mau H. The significance of proprioception on postural stabilization as assessed by ischemia. *Brain Res.* 1984;296(1):103-109.
110. Perry SD. Evaluation of age-related plantar-surface insensitivity and onset age of advanced insensitivity in older adults using vibratory and touch sensation tests. *Neurosci Lett.* 2006;392(1-2):62-67.
111. Dhruv NT, Niemi JB, Harry JD, Lipsitz LA, Collins JJ. Enhancing tactile sensation in older adults with electrical noise stimulation. *Neuroreport.* 2002;13(5):597-600.
112. Priplata AA, Niemi JB, Harry JD, Lipsitz LA, Collins JJ. Vibrating insoles and balance control in elderly people. *Lancet.* 2003;362(9390):1123-1124.
113. Priplata AA, Patriitti BL, Niemi JB, et al. Noise-enhanced balance control in patients with diabetes and patients with stroke. *Ann Neurol.* 2006;59(1):4-12.

114. Bernard-Demanze L, Burdet C, Berger L, Rougier P. Recalibration of somesthetic plantar information in the control of undisturbed upright stance maintenance. *J Integr Neurosci*. 2004;3(4):433-451.
115. Maurer C, Mergner T, Bolha B, Hlavacka F. Human balance control during cutaneous stimulation of the plantar soles. *Neurosci Lett*. 2001;302(1):45-48.
116. Ikeda ER, Borg A, Brown D, Malouf J, Showers KM, Li S. The valsalva maneuver revisited: the influence of voluntary breathing on isometric muscle strength. *J Strength Cond Res*. 2009;23(1):127-132.
117. Kaya D, Doral MN, Callaghan M. How can we strengthen the quadriceps femoris in patients with patellofemoral pain syndrome? *Muscles Ligaments Tendons J*. 2012;2(19):25-32.
118. Steinkamp LA, Dillingham MF, Markel MD, Hill JA, Kaufman KR. Biomechanical considerations in patellofemoral joint rehabilitation. *Am J Sports Med*. 1993;21(3):438-444.
119. Stiene HA, Brosky T, Reinking MF, Nyland J, Mason MB. A comparison of closed kinetic chain and isokinetic joint isolation exercise in patients with patellofemoral dysfunction. *J Orthop Sports Phys Ther*. 1996;24(3):136-141.
120. Rubenstein LZ, Josephson KR. The epidemiology of falls and syncope. *Clin Geriatr Med*. 2002;18(2):141-158.
121. Overend TJ, Cunningham DA, Kramer JF, Lefcoe MS, Paterson DH. Knee extensor and knee flexor strength: cross-sectional area ratios in young and elderly men. *J Gerontol*. 1992;47(6):M204-M210 OD - 1992/11/01.
122. Kubo K, Ishida Y, Komuro T, Tsunoda N, Kanehisa H, Fukunaga T. Age-related differences in the force generation capabilities and tendon extensibilities of knee extensors and plantar flexors in men. *J Gerontol A Biol Sci Med Sci*. 2007;62(11):1252-1258.
123. Lynch N a, Metter EJ, Lindle RS, et al. Muscle quality. I. Age-associated differences between arm and leg muscle groups. *J Appl Physiol*. 1999;86(10):188-194.
124. Watanabe K, Holobar A, Kouzaki M, Ogawa M, Akima H, Moritani T. Age-related changes in motor unit firing pattern of vastus lateralis muscle during low-moderate contraction. *Age (Dordr)*. 2016;38(3):48.
125. Abe T, Kawakami Y, Bemben M, Fukunaga T. Comparison of Age-Related, Site-Specific Muscle Loss Between Young and Old Active and Inactive Japanese Women. *J Geriatr Phys Ther*. 2011;34(4):168-173.
126. Abe T, Loenneke JP, Thiebaud RS, Fukunaga T. Age-related site-specific muscle wasting of upper and lower extremities and trunk in Japanese men and women. *Age (Omaha)*. 2014;36(2):813-821.
127. Zion AS, De Meersman R, Diamond BE, Bloomfield DM. A home-based resistance-training program using elastic bands for elderly patients with orthostatic hypotension. *Clin Auton Res*. 2003;13(4):286-292.
128. Cho S-I, An D-H. Effects of a Fall Prevention Exercise Program on Muscle Strength and Balance of the Old-old Elderly. *J Phys Ther Sci*. 2014;26(11):1771-1774.
129. Kim S, Lockhart T. Effects of 8 weeks of balance or weight training for the independently living elderly on the outcomes of induced slips. *Int J Rehabil Res*. 2010;33(1):49-55.
130. Witvrouw E, Danneels L, Van Tiggelen D, Willems TM, Cambier D. Open versus closed kinetic chain exercises in patellofemoral pain: a 5-year prospective randomized study. *Am J Sports Med*. 2004;32(5):1122-1130.
131. Campbell DE, Glenn W. Rehabilitation of knee flexor and knee extensor muscle strength in patients with meniscectomies, ligamentous repairs, and chondromalacia. *Phys Ther*. 1982;62(1):10-15.
132. Patten C, Kamen G, Rowland DM. Adaptations in maximal motor unit discharge rate to strength training in young and older adults. *Muscle Nerve*. 2001;24(4):542-550.
133. Bickel CS, Cross JM, Bamman MM. Exercise dosing to retain resistance training adaptations in young and older adults. *Med Sci Sports Exerc*. 2011;43(7):1177-1187.
134. Taaffe DR, Henwood TR, Nalls MA, Walker DG, Lang TF, Harris TB. Alterations in muscle attenuation following detraining and retraining in resistance-trained older adults. *Gerontology*. 2009;55(2):217-223.
135. Ivey FM, Tracy BL, Lemmer JT, et al. Effects of strength training and detraining on muscle quality: age and gender comparisons. *J Gerontol A Biol Sci Med Sci*. 2000;55(3):B152-B157; discussion B158-B159.
136. Yau DTY, Chung RCK, Pang MYC. Knee muscle strength and visual acuity are the most important modifiable predictors of falls in patients after hip fracture surgery: a prospective study. *Calcif Tissue Int*. 2013;92(3):287-295.
137. Cebolla EC, Rodacki ALF, Bento PCB. Balance, gait, functionality and strength: Comparison between elderly fallers and non-fallers. *Brazilian J Phys Ther*. 2015;19(2):146-151.
138. Bento PCB, Pereira G, Ugrinowitsch C, Rodacki ALF. Peak torque and rate of torque development in elderly with and without fall history. *Clin Biomech*. 2010;25(5):450-454.
139. Tiedemann AC, Sherrington C, Lord SR. Physical and psychological factors associated with stair negotiation performance in older people. *J Gerontol A Biol Sci Med Sci*. 2007;62(11):1259-1265.
140. Keskin D, Borman P, Ersöz M, Kurtaran A, Bodur H, Akyüz M. The Risk Factors Related to Falling in Elderly Females. *Geriatr Nurs (Minneap)*. 2008;29(1):58-63.
141. Morcelli MH, LaRoche DP, Crozara LF, et al. Neuromuscular performance in the hip joint of elderly fallers and non-fallers. *Aging Clinical and Experimental Research*. 2015. 28(3):443-50
142. Fielding RA, LeBrasseur NK, Cuoco A, Bean J, Mizer K, Fiatarone Singh MA. High-velocity resistance training increases skeletal muscle peak power in older women. *J Am Geriatr Soc*. 2002;50(4):655-662.

143. Reid KF, Callahan DM, Carabello RJ, Phillips EM, Frontera WR, Fielding R a. Lower extremity power training in elderly subjects with mobility limitations: a randomized controlled trial. *Aging Clin Exp Res*. 2008;20(4):337-343.
144. Vincent KR, Vincent KR, Braith RW, et al. Resistance exercise and physical performance in adults aged 60 to 83. *J Am Geriatr Soc*. 2002;50(6):1100-1107.
145. Chang SJ. *Improving lateral stability in older adults at risk of falls*. Thesis, School of Medicine, University of North Carolina, Chapel Hill 2007;2-4
146. Van Cutsem M, Duchateau J, Hainaut K. Changes in single motor unit behaviour contribute to the increase in contraction speed after dynamic training in humans. *J Physiol*. 1998;513 ( Pt 1):295-305.
147. Aagaard P. Training-induced changes in neural function. *Exerc Sport Sci Rev*. 2003;31(2):61-67.
148. Semmler JG. Motor unit synchronization and neuromuscular performance. *Exerc Sport Sci Rev*. 2002;30(1):8-14.
149. Miszko T a, Cress ME, Slade JM, Covey CJ, Agrawal SK, Doerr CE. Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci*. 2003;58(2):171-175.
150. Sayers SP, Bean J, Cuoco A, LeBrasseur NK, Jette A, Fielding R a. Changes in function and disability after resistance training: does velocity matter?: a pilot study. *Am J Phys Med Rehabil*. 2003;82(8):605-613.
151. Henwood TR, Taaffe DR. Improved physical performance in older adults undertaking a short-term programme of high-velocity resistance training. *Gerontology*. 2005;51(2):108-115.
152. Bolgla LA, Uhl TL. Electromyographic Analysis of Hip Rehabilitation Exercises in a Group of Healthy Subjects. *J Orthop Sports Phys Ther*. 2005;35(8):487-94
153. Neumann D a. Hip abductor muscle activity as subjects with hip prostheses walk with different methods of using a cane. *Phys Ther*. 1998;78(5):490-501.
154. Lord SR, Rogers MW, Howland a, Fitzpatrick R. Lateral stability, sensorimotor function and falls in older people. *J Am Geriatr Soc*. 1999;47(9):1077-1081.
155. Maki BE, Holliday PJ, Topper a K. A prospective study of postural balance and risk of falling in an ambulatory and independent elderly population. *J Gerontol*. 1994;49(2):M72-M84.
156. Rogers MW, Hedman LD, Johnson ME, Cain TD, Hanke T a. Lateral stability during forward-induced stepping for dynamic balance recovery in young and older adults. *J Gerontol A Biol Sci Med Sci*. 2001;56(9):M589-M594.
157. Johnson ME, Mille ML, Martinez KM, Crombie G, Rogers MW. Age-related changes in hip abductor and adductor joint torques. *Arch Phys Med Rehabil*. 2004;85(4):593-597.
158. Foldvari M, Clark M, Laviolette LC, et al. Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A Biol Sci Med Sci*. 2000;55(4):M192-M199.
159. Izquierdo M, Ibanez J, Gorostiaga E, et al. Maximal strength and power characteristics in isometric and dynamic actions of the upper and lower extremities in middle-aged and older men. *Acta Physiol Scand*. 1999;167(1):57-68.
160. Chang SHJ, Mercer VS, Giuliani CA, Sloane PD. Relationship between hip abductor rate of force development and mediolateral stability in older adults. *Arch Phys Med Rehabil*. 2005;86(9):1843-1850.
161. Hakkinen K, Kraemer WJ, Newton RU, Alen M. Changes in electromyographic activity, muscle fibre and force production characteristics during heavy resistance/power strength training in middle-aged and older men and women. *Acta Physiol Scand*. 2001;171(1):51-62.
162. Suetta C, Aagaard P, Rosted A, et al. Training-induced changes in muscle CSA, muscle strength, EMG, and rate of force development in elderly subjects after long-term unilateral disuse. *J Appl Physiol*. 2004;97(5):1954-1961.
163. Barry BK, Warman GE, Carson RG. Age-related differences in rapid muscle activation after rate of force development training of the elbow flexors. *Exp Brain Res*. 2005;162(1):122-132.
164. Maganaris CN, Narici M V, Reeves ND. In vivo human tendon mechanical properties: effect of resistance training in old age. *J Musculoskelet Neuronal Interact*. 2004;4(2):204-208.
165. Häkkinen K, Newton RU, Gordon SE, et al. Changes in muscle morphology, electromyographic activity, and force production characteristics during progressive strength training in young and older men. *J Gerontol A Biol Sci Med Sci*. 1998;53(6):B415-B423.
166. Chang SJ, Mercer VS, Guiliani CA, Morey MC, Sloane CPD, Williams S. Hip abductor exercise and lateral stability in older adults at risk of falls. *J Geriatr Phys Ther*. 2007;30(3):146-147.
167. Pasco JA, Holloway KL, Brennan-Olsen SL, Moloney DJ, Kotowicz MA. Muscle strength and areal bone mineral density at the hip in women: a cross-sectional study. *BMC Musculoskelet Disord*. 2015;16:124.
168. Bayramoglu M, Sözüy S, Karatas M, Kuilinc S. Relationships between muscle strength and bone mineral density of three body regions in sedentary postmenopausal women. *Rheumatol Int*. 2005;25(7):513-517.168.
169. Nelson ME, Fiatarone M a, Morganti CM, Trice I, Greenberg R a, Evans WJ. Effects of high-intensity strength training on multiple risk factors for osteoporotic fractures. A randomized controlled trial. *JAMA*. 1994;272:1909-1914.
170. Maki BE, Mclroy WE. Control of rapid limb movements for balance recovery: Age-related changes and implications for fall prevention. In: *Age and Ageing*. Vol 2006;35 Suppl 2:ii12-ii18. Review.

171. McIlroy WE, Maki BE. Age-related changes in compensatory stepping in response to unpredictable perturbations. *J Gerontol A Biol Sci Med Sci*. 1996;51(6):M289-M296.
172. Roden-Reynolds DC, Walker MH, Wasserman CR, Dean JC. Hip proprioceptive feedback influences the control of mediolateral stability during human walking. *J Neurophysiol*. 2015;114(4):2220-2229.
173. Bok SK, Lee TH, Lee SS. The effects of changes of ankle strength and range of motion according to aging on balance. *Ann Rehabil Med*. 2013;37(1):10-16.
174. Vandervoort a a, McComas a J. Contractile changes in opposing muscles of the human ankle joint with aging. *J Appl Physiol*. 1986;61(1):361-367.
175. Onambele GL, Narici M V, Maganaris CN. Calf muscle-tendon properties and postural balance in old age. *J Appl Physiol*. 2006;100(6):2048-2056.
176. Simoneau E, Martin A, Porter MM, Van Hoecke J. Strength training in old age: Adaptation of antagonist muscles at the ankle joint. *Muscle and Nerve*. 2006;33(4):546-555.
177. Simoneau E, Martin A, Van Hoecke J. Adaptations to long-term strength training of ankle joint muscles in old age. *Eur J Appl Physiol*. 2007;100(5):507-514.
178. Padberg FTJ, Johnston M V, Sisto SA. Structured exercise improves calf muscle pump function in chronic venous insufficiency: a randomized trial. *J Vasc Surg*. 2004;39(1):79-87.
179. O'Brien JA, Edwards HE, Finlayson KJ, Kerr GK. Understanding the relationships between the calf muscle pump, ankle range of motion and healing for adults with venous leg ulcers: a review of the literature. *Fac Heal Inst Heal Biomed Innov Sch Exerc Nutr Sci Sch Nurs*. 2012.20(2)80-85.
180. Orsted HL, Radke L, Gorst R. The impact of musculoskeletal changes on the dynamics of the calf muscle pump. *Ostomy Wound Manage*. 2001;47(10):18-24.
181. Recek C. Impact of the calf perforators on the venous hemodynamics in primary varicose veins. *J Cardiovasc Surg (Torino)*. 2006;47(6):629-635.
182. Gross EA, Wood CR, Lazarus GS, Margolis DJ. Venous leg ulcers: an analysis of underlying venous disease. *Br J Dermatol*. 1993;129(3):270-274.
183. Meissner MH, Moneta G, Burnand K, et al. The hemodynamics and diagnosis of venous disease. *J Vasc Surg*. 2007;46(6 SUPPL.).
184. Gardner AM, Fox RH. The venous footpump: influence on tissue perfusion and prevention of venous thrombosis. *Ann Rheum Dis*. 1992;51(10):1173-1178.
185. Styf J. The venous pump of the human foot. *Clin Physiol*. 1990;10(1):77-84.
186. Broderick BJ, Dessus S, Grace PA, O'Laughlin G. Technique for the computation of lower leg muscle bulk from magnetic resonance images. *Med Eng Phys*. 2010;32(8):926-933.
187. Kawakami R, Murakami H, Sanada K, et al. Calf circumference as a surrogate marker of muscle mass for diagnosing sarcopenia in Japanese men and women. *Geriatr Gerontol Int*. 2015;15(8):969-976.
188. Rolland Y, Lauwers-Cances V, Cournot M, et al. Sarcopenia, calf circumference, and physical function of elderly women: A cross-sectional study. *J Am Geriatr Soc*. 2003;51(8):1120-1124.
189. Velazquez-Alva MC, Irigoyen Camacho ME, Lazarevich I, Delgadillo Velazquez J, Acosta Dominguez P, Zepeda Zepeda MA. Comparison of the prevalence of sarcopenia using skeletal muscle mass index and calf circumference applying the European consensus definition in elderly Mexican women. *Geriatr Gerontol Int*. 2015;[Epub ahead of print]
190. Kuyumcu ME, Halil M, Kara O et al. Ultrasonographic evaluation of the calf muscle mass and architecture in elderly patients with and without sarcopenia. *Arch Gerontol Geriatr*. 2016;65:218-224.
191. Rolland Y, Lauwers-Cances V, Cournot M, et al. Sarcopenia, calf circumference, and physical function of elderly women: A cross-sectional study. *J Am Geriatr Soc*. 2003;51(8):1120-1124.
192. Diaz Villegas GM, Runzer Colmenares F. Calf circumference and its association with gait speed in elderly participants at Peruvian Naval Medical Center. *Revista Espanola de Geriatria y Gerontologia*. 2015. 50(1):22-5.
193. Pérez-Zepeda MU, Gutiérrez-Robledo LM. Calf circumference predicts mobility disability: A secondary analysis of the Mexican health and ageing study. *Eur Geriatr Med*. 2016;7(3):262-266.
194. Rice CL, Cunningham D a, Paterson DH, Lefcoe MS. Arm and leg composition determined by computed tomography in young and elderly men. *Clin Physiol*. 1989;9(3):207-220.
195. Whipple RH, Wolfson LI, Amerman PM. The relationship of knee and ankle weakness to falls in nursing home residents: An isokinetic study. *J Am Geriatr Soc*. 1987;35(1):13-20.
196. MacRae PG, Lacourse M, Moldavon R. Physical performance measures that predict faller status in community-dwelling older adults. *J Orthop Sports Phys Ther*. 1992;16(3):123-128.
197. Daubney ME, Culham EG. Lower-extremity muscle force and balance performance in adults aged 65 years and older. *Phys Ther*. 1999;79(12):1177-1185.
198. Wolfson L, Judge J, Whipple R, King M. Strength is a major factor in balance, gait, and the occurrence of falls. *J Gerontol A Biol Sci Med Sci*. 1995;50 Spec No:64-67.

199. Abd El- Kader SM. Ankle Dorsiflexors Strength Improves Balance Performance in Elderly: A Correlational Study. *Eur J Gen Med*. 2014;11(2):60-65.
200. Hess JA, Woollacott M. Effect of high-intensity strength-training on functional measures of balance ability in balance-impaired older adults. *J Manipulative Physiol Ther*. 2005;28(8):582-590.
201. Wolfson L, Whipple R, Derby C, et al. Balance and strength training in older adults: intervention gains and Tai Chi maintenance. *J Am Geriatr Soc*. 1996;44(5):498-506.
202. Rubenstein LZ, Josephson KR, Trueblood PR, et al. Effects of a group exercise program on strength, mobility, and falls among fall-prone elderly men. *J Gerontol A Biol Sci Med Sci*. 2000;55(6):M317-M321.
203. Tinetti ME, Franklin Williams T, Mayewski R. Fall risk index for elderly patients based on number of chronic disabilities. *Am J Med*. 1986;80(3):429-434.
204. Crilly RG, Willems DA, Trenholm KJ, Hayes KC, Delaquerriere-Richardson LFO. Effect of exercise on postural sway in the elderly. *Gerontology*. 1989;35(2-3):137-143.
205. Hornbrook MC, Stevens VJ, Wingfield DJ, Hollis JF, Greenlick MR, Ory MG. Preventing falls among community-dwelling older persons: results from a randomized trial. *Gerontologist*. 1994;34(1):16-23.
206. Fiatarone MA, Evans WJ. The etiology and reversibility of muscle dysfunction in the aged. *J Gerontol*. 1993;48 Spec No:77-83.
207. Hu MH, Woollacott MH. Multisensory training of standing balance in older adults: II. Kinematic and electromyographic postural responses. *J Gerontol*. 1994;49(2):M62-M71.
208. Woollacott MH, Shumway-Cook A, Nashner LM. Aging and posture control: changes in sensory organization and muscular coordination. *Int J Aging Hum Dev*. 1986;23(2):97-114.
209. Manchester D, Woollacott M, Zederbauer-Hylton N, Marin O. Visual, Vestibular and Somatosensory Contributions to Balance Control in the Older Adult. *J Gerontol*. 1989;44(4):M118-M127.
210. Epley B. Pounding chart. In: *Boyd Epley Workout*. Lincoln, NE: Body Enterprises; 1985:86.
211. Wood TM, Maddalozzo GF, Harter R a. Accuracy of Seven Equations for Predicting 1-RM Performance of Apparently Healthy, Sedentary Older Adults. *Meas Phys Educ Exerc Sci*. 2002;6(2):67-94.
212. Thiele RM, Conchola EC, Palmer TB, DeFreitas JM, Thompson BJ. The effects of a high-intensity free-weight back-squat exercise protocol on postural stability in resistance-trained males. *J Sports Sci*. 2014;33(2):211-218.
213. Fernie GR, Gryfe CI, Holliday PJ, Llewellyn a. The relationship of postural sway in standing to the incidence of falls in geriatric subjects. *Age Ageing*. 1982;11:11-16.
214. Fujita E, Takeshima N, Kato Y, et al. Effects of Body-weight Squat Training on Muscular Size, Strength and Balance Ability in Physically Frail Older Adults. *Int J Sport Heal Sci*. 2016. [Epub ahead of print]
215. Flanagan S, Salem GJ, Wang MY, Sanker SE, Greendale GA. Squatting exercises in older adults: Kinematic and kinetic comparisons. *Med Sci Sports Exerc*. 2003;35(4):635-643.
216. Osugi T, Iwamoto J, Yamazaki M, Takakuwa M. Effect of a combination of whole body vibration exercise and squat training on body balance, muscle power, and walking ability in the elderly. *Ther Clin Risk Manag*. 2014;10(1):131-138.
217. Runge M, Rehfeld G, Resnicek E. Balance training and exercise in geriatric patients. *J Musculoskelet Neuronal Interact*. 2000;1(1):61-65.
218. Runge M, Hunger G. Determinants of musculoskeletal frailty and the risk of falls in old age. *J Musculoskelet Neuronal Interact*. 2006;6(2):167-173.
219. Frost HM. Defining osteopenias and osteoporoses: Another view (with insights from a new paradigm). *Bone*. 1997;20(5):385-391.
220. Schiessl H, Frost HM, Jee WSS. Estrogen and bone-muscle strength and mass relationships. *Bone*. 1998;22(1):1-6.
221. Laufer Y. Effect of age on characteristics of forward and backward gait at preferred and accelerated walking speed. *J Gerontol A Biol Sci Med Sci*. 2005;60(5):627-632.
222. Fritz NE, Worstell AM, Kloos AD, Siles AB, White SE, Kegelmeyer DA. Backward walking measures are sensitive to age-related changes in mobility and balance. *Gait Posture*. 2013;37(4):593-597.
223. Agmon M, Kodesh E, Kizony R. The Effect of Different Types of Walking on Dual-Task Performance and Task Prioritization among Community-Dwelling Older Adults. *Sci World J*. 2014;2014:1-6.
224. Hooper TL, Dunn DM, Erick Props J, et al. The Effects of Graded Forward and Backward Walking on Heart Rate and Oxygen Consumption. *J Orthop Sports Phys Ther*. 2004 34(2):65-71
225. Lee M, Kim J, Son J, Kim Y. Kinematic and kinetic analysis during forward and backward walking. *Gait Posture*. 2013;38(4):674-678.
226. Godde B, Voelcker-Rehage C. More automation and less cognitive control of imagined walking movements in high- versus low-fit older adults. *Front Aging Neurosci*. 2010;2(September):1-13.
227. Hoogkamer W, Meyns P, Duysens J. Steps forward in understanding backward gait: From basic circuits to rehabilitation. *Exerc Sport Sci Rev*. 2014;42(1):23-29.
228. Handford ML, Srinivasan M. Sideways walking: preferred is slow, slow is optimal, and optimal is expensive. *Biol Lett*. 2014;10(January):20131006.



229. Melzer I, Benjuya N, Kaplanski J. Postural stability in the elderly: A comparison between fallers and non-fallers. *Age Ageing*. 2004;33(6):602-607.
230. Bryant MS, Workman CD, Hou J-GG, Henson HK, York MK. Acute and Long-Term Effects of Multidirectional Treadmill Training on Gait and Balance in Parkinson Disease. *PM R*. 2016. [Epub ahead of print]
231. Long LL, Srinivasan M. Walking, running, and resting under time, distance, and average speed constraints: optimality of walk-run-rest mixtures. *J R Soc Interface*. 2013;10:20120980.
232. Glaister BC, Bernatz GC, Klute GK, Orendurff MS. Video task analysis of turning during activities of daily living. *Gait Posture*. 2007;25(2):289-294.
233. Segal AD, Orendurff MS, Czerniecki JM, Shofer JB, Klute GK. Local dynamic stability in turning and straight-line gait. *J Biomech*. 2008;41(7):1486-1493.
234. Thigpen MT, Light KE, Creel GL, Flynn SM. Turning difficulty characteristics of adults aged 65 years or older. *Phys Ther*. 2000;80(12):1174-1187.
235. Dite W, Temple VA. Development of a Clinical Measure of Turning for Older Adults. *Am J Phys Med Rehabil*. 2002;81(11):857-866.
236. Asmidawati A, Hamid TA, Hussain RM, Hill KD. Home based exercise to improve turning and mobility performance among community dwelling older adults: protocol for a randomized controlled trial. *BMC Geriatr*. 2014;14(1):100.
237. Cumming RG, Klineberg RJ. Fall frequency and characteristics and the risk of hip fractures. *J Am Geriatr Soc*. 1994;42:774-778.
238. Nagano H, Sparrow W a, Begg RK. Biomechanical characteristics of slipping during unconstrained walking, turning, gait initiation and termination. *Ergonomics*. 2013;56(6):1038-1048.
239. de Dreu MJ, van der Wilk a. SD, Poppe E, Kwakkel G, van Wegen EEH. Rehabilitation, exercise therapy and music in patients with Parkinson's disease: a meta-analysis of the effects of music-based movement therapy on walking ability, balance and quality of life. *Parkinsonism Relat Disord*. 2012;18:S114-S119.
240. Schaffert N, Thaut MH, Mattes K. Rhythm-based regulation/modification of movements in high-performance rowing and neurologic rehabilitaton. In: *Proceedings of ISON, 4th Interactive Sonification Workshop*. Erlangen, Germany: Fraunhofer IIS; 2013:67-72.
241. Thaut MH. Rhythm, Music and the Brain: Scientific Foundations and Clinical Applications. New York: Routledge; 2005.
242. Thaut M, Abiru M. Rhythmic Auditory Stimulation in Rehabilitation of Movement Disorders: A Review Of Current Research. *Music Percept*. 2010;27(4):263-269.
243. Bengtsson SL, Ullen F, Henrik Ehrsson H, et al. Listening to rhythms activates motor and premotor cortices. *Cortex*. 2009;45(1):62-71.
244. Peretz I, Zatorre RJ. Brain organization for music processing. *Annu Rev Psychol*. 2005;56:89-114.
245. Kenyon GP, Thaut MH. Rhythmic-driven Optimization of Motor Control. In: *Rhythm, Music and the Brain: Scientific Foundations and Clinical Applications*. New York: Routledge Champman & Hall; 2005:82-112.
246. Thaut MH, Miller R a, Schauer LM. Multiple synchronization strategies in rhythmic sensorimotor tasks: phase vs period correction. *Biol Cybern*. 1998;79:241-250.
247. Winter DA, Patla AE, Prince F, Ishac M, Gielo-Perczak K. Stiffness Control of Balance in Quiet Standing. *J Neurophysiol*. 1998;80(3):1211-1221.
248. Sozzi S, Honeine JL, Do MC, Schieppati M. Leg muscle activity during tandem stance and the control of body balance in the frontal plane. *Clin Neurophysiol*. 2013;124(6):1175-1186.
249. Jonsson E, Seiger Å, Hirschfeld H. Postural steadiness and weight distribution during tandem stance in healthy young and elderly adults. *Clin Biomech*. 2005;20(2):202-208.
250. Kavounoudias A, Roll R, Roll JP. The plantar sole is a "dynamometric map" for human balance control. *Neuroreport*. 1998;9(14):3247-3252.
251. Henry SM, Fung J, Horak FB. Effect of stance width on multidirectional postural responses. *J Neurophysiol*. 2001;85(2):559-570.
252. Roll JP, Vedel JP, Roll R. Eye, head and skeletal muscle spindle feedback in the elaboration of body references. *Prog Brain Res*. 1989;80:113-123; discussion 57-60.
253. Di Giulio I, Maganaris CN, Baltzopoulos V, Loram ID. The proprioceptive and agonist roles of gastrocnemius, soleus and tibialis anterior muscles in maintaining human upright posture. *J Physiol*. 2009;587(Pt 10):2399-2416.
254. Quoniam C, Hay L, Roll JP, Harlay F. Age effects on reflex and postural responses to propriomuscular inputs generated by tendon vibration. *J Gerontol A Biol Sci Med Sci*. 1995;50(3):B155-B165.
255. Benjuya N, Melzer I, Kaplanski J. Aging-induced shifts from a reliance on sensory input to muscle cocontraction during balanced standing. *Journals Gerontol A, Biol Sci Med Sci*. 2004;59(2):166-171.
256. Slobounov S, Hallett M, Stanhope S, Shibasaki H. Role of cerebral cortex in human postural control: An EEG study. *Clin Neurophysiol*. 2005;116(2):315-323.
257. Bove M, Trompetto C, Abbruzzese G, Schieppati M. The posture-related interaction between Ia-afferent and descending input on the spinal reflex excitability in humans. *Neurosci Lett*. 2006;397(3):301-306.
258. Fitzpatrick R, Burke D, Gandevia SC. Loop gain of reflexes controlling human standing measured with the use of postural and vestibular disturbances. *J Neurophysiol*. 1996;76(6):3994-4008.

259. Solopova IA, Kazennikov O V., Deniskina NB, Levik YS, Ivanenko YP. Postural instability enhances motor responses to transcranial magnetic stimulation in humans. *Neurosci Lett.* 2003;337(1):25-28.
260. Reynolds RF. The ability to voluntarily control sway reflects the difficulty of the standing task. *Gait Posture.* 2010;31(1):78-81.
261. Remaud A, Boyas S, Caron GAR, Bilodeau M. Attentional Demands Associated With Postural Control Depend on Task Difficulty and Visual Condition. *J Mot Behav.* 2012;44(5):329-340.
262. Teasdale N, Bard C, Larue J, Fleury M. On the cognitive penetrability of posture control. *Exp Aging Res.* 1993;19(1):1-13.
263. Sozzi S, Do MC, Monti A, Schieppati M. Sensorimotor integration during stance: Processing time of active or passive addition or withdrawal of visual or haptic information. *Neuroscience.* 2012;212:59-76.
264. Chiu a YY, Au-Yeung SSY, Lo SK. A comparison of four functional tests in discriminating fallers from non-fallers in older people. *Disabil Rehabil.* 2003;25(1):45-50.
265. Stel VS, Smit JH, Pluijm SMF, Lips P. Balance and mobility performance as treatable risk factors for recurrent falling in older persons. *J Clin Epidemiol.* 2003;56(7):659-668.
266. Maughan KK. *Does balance training improve balance in physically active older adults?* Thesis, Department of Science, Iowa State University 2008;p9
267. Hiroyuki S, Uchiyama Y, Kakurai S. Specific effects of balance and gait exercises on physical function among the frail elderly. *Clin Rehabil.* 2003;17(5):472-479.
268. Islam MM, Nasu E, Rogers ME, Koizumi D, Rogers NL, Takeshima N. Effects of combined sensory and muscular training on balance in Japanese older adults. *Prev Med (Baltim).* 2004;39(6):1148-1155.
269. Hile ES, Brach JS, Perera S, Wert DM, VanSwearingen JM, Studenski S a. Interpreting the need for initial support to perform tandem stance tests of balance. *Phys Ther.* 2012;92(10):1316-1328.
270. Morris R, Harwood RH, Baker R, Sahota O, Armstrong S, Masud T. A comparison of different balance tests in the prediction of falls in older women with vertebral fractures: a cohort study. *Age Ageing.* 2007;36(1):78-83.
271. Nevitt MC, Cummings SR, Kidd S, Black D. Risk factors for recurrent nonsyncopal falls. A prospective study. *JAMA.* 1989;261(18):2663-2668.
272. Dargent-Molina P, Favier F, Grandjean H, et al. Fall-related factors and risk of hip fracture: The EPIDOS prospective study. *Lancet.* 1996;348(9021):145-149.
273. Cummings SR, Nevitt MC, Browner WS, et al. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. *N Engl J Med.* 1995;332(12):767-773.
274. Lark SD, Pasupuleti S. Validity of a Functional Dynamic Walking Test for the Elderly. *Arch Phys Med Rehabil.* 2009;90(3):470-474.
275. Chantanachai T, Pichaiyongwongdee S, Jalayondeja C. Fall prediction in thai elderly with timed up and go and tandem walk test: a cross-sectional study. *J Med Assoc Thai.* 2014;97 Suppl 7:S21-S25.
276. Mihara M, Miyai I, Hatakenaka M, Kubota K, Sakoda S. Role of the prefrontal cortex in human balance control. *Neuroimage.* 2008;43(2):329-336.
277. Salazar-Gonzalez BC, Cruz-Quevedo JE, Gallegos-Cabrales EC, et al. A Physical-Cognitive Intervention to Enhance Gait Speed in Older Mexican Adults. *Am J Health Promot.* 2015;30(2):77-84.
278. Bloem BR, Grimbergen YAM, van Dijk JG, Munneke M. The "posture second" strategy: A review of wrong priorities in Parkinson's disease. *J Neurol Sci.* 2006;248(1-2):196-204.
279. Madureira MM, Takayama L, Gallinaro AL, Caparbo VF, Costa RA, Pereira RMR. Balance training program is highly effective in improving functional status and reducing the risk of falls in elderly women with osteoporosis: a randomized controlled trial. *Osteoporos Int.* 2007;18:419-425.
280. Mansfield A, Peters AL, Liu BA, Maki BE. A perturbation-based balance training program for older adults: study protocol for a randomised controlled trial. *BMC Geriatr.* 2007;7:12.
281. Cyarto E V., Brown WJ, Marshall AL, Trost SG. Comparative effects of home- and group-based exercise on balance confidence and balance ability in older adults: Cluster randomized trial. *Gerontology.* 2008;54:272-280.
282. Addison O, Young P, Inacio M, et al. Hip but Not Thigh Intramuscular Adipose Tissue is Associated with Poor Balance and Increased Temporal Gait Variability in Older Adults. *Curr Aging Sci.* 2014.7(2):137-43.
283. Inacio M, Ryan AS, Bair W-N, Prettyman M, Beamer B a, Rogers MW. Gluteal muscle composition differentiates fallers from non-fallers in community dwelling older adults. *BMC Geriatr.* 2014;14:37.
284. Leavey VJ, Sandrey M a, Dahmer G. Comparative effects of 6-week balance, gluteus medius strength, and combined programs on dynamic postural control. *J Sport Rehabil.* 2010;19(3):268-287.
285. Jonsson E, Seiger Å, Hirschfeld H. One-leg stance in healthy young and elderly adults: A measure of postural steadiness? *Clin Biomech.* 2004;19(7):688-694.
286. Hertel J, Gay MR, Denegar CR. Differences in postural control during single-leg stance among healthy individuals with different foot types. *J Athl Train.* 2002;37(2):129-132.
287. Beurskens R, Gollhofer A, Muehlbauer T, Cardinale M, Granacher U. Effects of heavy-resistance strength and balance training on unilateral and bilateral leg strength performance in old adults. *PLoS One.* 2015;10(2).

288. Goble DJ, Coxon JP, Van Impe a., et al. Brain Activity during Ankle Proprioceptive Stimulation Predicts Balance Performance in Young and Older Adults. *J Neurosci*. 2011;31(45):16344-16352.
289. Inglis JT, Horak FB, Shupert CL, Jones-Rycewicz C. The importance of somatosensory information in triggering and scaling automatic postural responses in humans. *Exp Brain Res*. 1994;101(1):159-164.
290. Lajoie Y, Teasdale N, Cole JD, et al. Gait of a deafferented subject without large myelinated sensory fibers below the neck. *Neurology*. 1996;47:109-115.
291. Horak FB, Dickstein R, Peterka RJ. Diabetic neuropathy and surface sway-referencing disrupt somatosensory information for postural stability in stance. *Somatosens Mot Res*. 2002;19(4):316-326.
292. Speers RA, Kuo AD, Horak FB. Contributions of altered sensation and feedback responses to changes in coordination of postural control due to aging. *Gait Posture*. 2002;16(1):20-30.
293. Lord SR, Clark RD, Webster IW. Postural stability and associated physiological factors in a population of aged persons. *J Gerontol*. 1991;46(3):M69-M76.
294. Simoneau GG, Leibowitz HW, Ulbrecht JS, Tyrrell R a, Cavanagh PR. The effects of visual factors and head orientation on postural steadiness in women 55 to 70 years of age. *J Gerontol*. 1992;47(5):M151-M158.
295. Goble DJ, Coxon JP, Van Impe A, et al. The neural basis of central proprioceptive processing in older versus younger adults: an important sensory role for right putamen. *Hum Brain Mapp*. 2012;33(4):895-908.
296. Tinetti ME, Richman D, Powell L. Falls efficacy as a measure of fear of falling. *J Gerontol*. 1990;45(6):P239-P243.
297. Maki BE, Holliday PJ, Topper a K. Fear of falling and postural performance in the elderly. *J Gerontol*. 1991;46(4):M123-M131.
298. Wilson M-MG, Miller DK, Andresen EM, Malmstrom TK, Miller JP, Wolinsky FD. Fear of falling and related activity restriction among middle-aged African Americans. *J Gerontol A Biol Sci Med Sci*. 2005;60(3):355-360.
299. Curcio C-L, Gomez F, Reyes-Ortiz C a. Activity restriction related to fear of falling among older people in the Colombian Andes mountains: are functional or psychosocial risk factors more important? *J Aging Health*. 2009;21(3):460-479.
300. Fletcher PC, Hirdes JP. Restriction in activity associated with fear of falling among community-based seniors using home care services. *Age Ageing*. 2004;33(3):273-279.
301. Delbaere K, Crombez G, Vanderstraeten G, Willems T, Cambier D. Fear-related avoidance of activities, falls and physical frailty. A prospective community-based cohort study. *Age Ageing*. 2004;33(4):368-373.
302. Gomez F, Curcio CL. The development of a fear of falling interdisciplinary intervention program. *Clin Interv Aging*. 2007;2(4):661-667.
303. Young WR, Mark Williams A. How fear of falling can increase fall-risk in older adults: Applying psychological theory to practical observations. *Gait Posture*. 2015;41(1):7-12.
304. Adkin AL, Frank JS, Carpenter MG, Peysar GW. Postural control is scaled to level of postural threat. *Gait Posture*. 2000;12(2):87-93.
305. Brown LA, Gage WH, Polych MA, Sleik RJ, Winder TR. Central set influences on gait: Age-dependent effects of postural threat. *Exp Brain Res*. 2002;145(3):286-296.
306. Hiller CE, Refshauge KM, Herbert RD, Kilbreath SL. Balance and recovery from a perturbation are impaired in people with functional ankle instability. *Clin J Sport Med*. 2007;17(4):269-275.
307. Perry J, Burnfield JM, Gronley JK, Mulroy SJ. Toe walking: Muscular demands at the ankle and knee. *Arch Phys Med Rehabil*. 2003;84(1):7-16.
308. Cunningham CB, Schilling N, Anders C, Carrier DR, Carrier DR. The influence of foot posture on the cost of transport in humans. *J Exp Biol*. 2010;213(5):790-797.
309. Barbour KE, Lui L-Y, McCulloch CE, et al. Trajectories of Lower Extremity Physical Performance: Effects on Fractures and Mortality in Older Women. *J Gerontol A Biol Sci Med Sci*. 2016. [Epub ahead of print]
310. Najafi DA, Dahlberg LE, Hansson EE. A combination of clinical balance measures and FRAX® to improve identification of high-risk fallers. *BMC Geriatr*. 2016;16(1):94.
311. Kim JW, Koh JM, Park JH, Chang JS. Validation of FRAX without BMD: An age-related analysis of the Fifth Korean National Health and Nutrition Examination Survey (KNHANES V-1, 2010). *Bone*. 2015;75:27-31.
312. Rosie J, Taylor D. Sit-to-stand as home exercise for mobility-limited adults over 80 years of age - GrandStand System may keep you standing? *Age Ageing*. 2007;36(5):555-562.
313. Lord SR, Murray SM, Chapman K, Munro B, Tiedemann A. Sit-to-stand performance depends on sensation, speed, balance, and psychological status in addition to strength in older people. *J Gerontol A Biol Sci Med Sci*. 2002;57(8):M539-M543.
314. Rosenberg DE, Gell NM, Jones SMW, et al. The Feasibility of Reducing Sitting Time in Overweight and Obese Older Adults. *Health Educ Behav*. 2015;42(5):669-76.
315. Gardiner PA, Eakin EG, Healy GN, Owen N. Feasibility of reducing older adults' sedentary time. *Am J Prev Med*. 2011;41(2):174-177.
316. Miller WR, Rose GS. Toward a theory of motivational interviewing. *Am Psychol*. 2009;64(6):527-537.
317. Pauls JL. Review of stair-safety research with an emphasis on Canadian studies. *Ergonomics*. 1985;28(7):999-1010.
318. Cohen HH, Templer J, Archea J. An analysis of occupational stair accident patterns. *J Safety Res*. 1985;16(4):171-181.

319. Nagata H. Analysis of fatal falls on the same level or on stairs/steps. *Saf Sci*. 1991;14(3-4):213-222.
320. Templer J, Archea J, Chen HH. Study of factors associated with risk of work-related stairway falls. *J Safety Res*. 1985;16(4):183-196.
321. Jackson PL, Cohen HH. An in-depth investigation of 40 stairway accidents and the stair safety literature. *J Safety Res*. 1995;26(3):151-159.
322. Kim BJ. Prevention of falls during stairway descent in older adults. *Appl Ergon*. 2009;40(3):348-352.
323. Ozanne-Smith J, Guy J, Kelly M, Clapperton A. *The Relationship between Slips, Trips and Falls and the Design and Construction of Buildings*. Monash University Accident Research Centre, Clayton; 2008.pp14 - 230
324. Maki BE, Perry SD, Mclroy WE. Efficacy of handrails in preventing stairway falls: A new experimental approach. *Saf Sci*. 1998;28(3):189-206.
325. Loo-Morrey M, Hallas K, Thorpe S. Minimum going requirements and friction demands during stair descent. In: *Contemporary Ergonomics*. CRC Press; Boca Raton London New York Washington, D.C 2004,pp8-12.
326. Kreisfeld R, Newson R. *Hip Fracture Injuries*. NISU briefing, Flinders University, Adelaide; 2006.
327. Cayless SM. Slip, trip and fall accidents: Relationship to building features and use of coroners' reports in ascribing cause. In: *Applied Ergonomics*.Vol 32.; 2001:155-162.
328. Ragg M, Hwang S, Steinhart B. Analysis of serious injuries caused by stairway falls. *Emerg Med Australas*. 2000;12(1):45-49.
329. Startzell JK, Owens D a, Mulfinger LM, Cavanagh PR. Stair negotiation in older people: a review. *J Am Geriatr Soc*. 2000;48(5):567-580.
330. Hortobágyi T, Mizelle C, Beam S, DeVita P. Old adults perform activities of daily living near their maximal capabilities. *J Gerontol A Biol Sci Med Sci*. 2003;58(5):M453-M460.
331. Ploutz-Snyder LL, Manini T, Ploutz-Snyder RJ, Wolf D a. Functionally relevant thresholds of quadriceps femoris strength. *J Gerontol A Biol Sci Med Sci*. 2002;57(4):B144-B152.
332. Salem GJ, Wang MY, Young JT, Marion M, Greendale G a. Knee strength and lower- and higher-intensity functional performance in older adults. *Med Sci Sport Exerc*. 2000;32(10):1679-1684.
333. Simoneau GG, Cavanagh PR, Ulbrecht JS, Leibowitz HW, Tyrrell RA. The influence of visual factors on fall-related kinematic variables during stair descent by older women. *J Gerontol*. 1991;46(6):M188-M195.
334. Tikkanen O, Sipilä S, Kuula A-S, Pesola A, Haakana P, Finni T. Muscle activity during daily life in the older people. *Aging Clin Exp Res*. 2015.28(4):713-20
335. Young WR, Wing AM, Hollands MA. Influences of state anxiety on gaze behavior and stepping accuracy in older adults during adaptive locomotion. *Journals Gerontol - Ser B Psychol Sci Soc Sci*. 2012;67 B(1):43-51.
336. Young WR, Hollands MA. Can telling older adults where to look reduce falls? Evidence for a causal link between inappropriate visual sampling and suboptimal stepping performance. *Exp Brain Res*. 2010;204(1):103-113.
337. Deshpande N, Metter EJ, Lauretani F, Bandinelli S, Ferrucci L. Interpreting fear of falling in the elderly: what do we need to consider? *J Geriatr Phys Ther*. 2009;32(3):91-96.
338. Rikli RE, Jones CJ. Development and validation of criterion-referenced clinically relevant fitness standards for maintaining physical independence in later years. *Gerontologist*. 2013;53(2):255-267.
339. Kuo YL. The influence of chair seat height on the performance of community-dwelling older adults' 30-second chair stand test. *Aging Clin Exp Res*. 2013;25(3):305-309.
340. Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. *Phys Ther*. 2000;80:896-903.
341. Barry E, Galvin R, Keogh C, Horgan F, Fahey T. Is the Timed Up and Go test a useful predictor of risk of falls in community dwelling older adults: a systematic review and meta-analysis. *BMC Geriatr*. 2014;14(1):14.