

REFERENCES

1. Knobloch K, Vogt PM. [Nordic pole walking injuries--nordic walking thumb as novel injury entity]. *Sportverletz Sportschaden*. 2006;20(3):137-142.
2. Lejczak A, Josiak K, Wegrzynowska-Teodorczyk K, et al. Nordic Walking May Safely Increase the Intensity of Exercise Training in Healthy Subjects and in Patients with Chronic Heart Failure. *Adv Clin Exp Med*. 25(1):145-149.
3. Piotrowicz E, Buchner T, Piotrowski W, Piotrowicz R. Influence of home-based telemonitored Nordic walking training on autonomic nervous system balance in heart failure patients. *Arch Med Sci*. 2015;11(6):1205-1212.
4. Piotrowicz E, Zielinski T, Bodalski R, et al. Home-based telemonitored Nordic walking training is well accepted, safe, effective and has high adherence among heart failure patients, including those with cardiovascular implantable electronic devices: a randomised controlled study. *Eur J Prev Cardiol*. 2015;22(11):1368-1377.
5. Keast M-L, Sloviniec D'Angelo ME, Nelson CRM, et al. Randomized trial of Nordic walking in patients with moderate to severe heart failure. *Can J Cardiol*. 2013;29(11):1470-1476.
6. Knobloch K. No difference in the hemodynamic response to Nordic pole walking vs. conventional brisk walking - A randomized exercise field test using the ultrasonic cardiac output monitor (USCOM). *Int J Cardiol*. 2009;132(1):133-135.
7. Salim M, Brodin E, Spaals-Abrahamsson Y, Berntorp E, Zetterberg E. The effect of Nordic Walking on joint status, quality of life, physical ability, exercise capacity and pain in adult persons with haemophilia. *Blood Coagul Fibrinolysis*. 2016;27(4):467-472.
8. Buliska K, Kropielnicka K, Jasiski T, et al. Nordic pole walking improves walking capacity in patients with intermittent claudication: a randomized controlled trial. *Disabil Rehabil*. 2016;38(13):1318-24
9. Spafford C, Oakley C, Beard JD. Randomized clinical trial comparing Nordic pole walking and a standard home exercise programme in patients with intermittent claudication. *Br J Surg*. 2014;101(7):760-767.
10. Oakley C, Zwierska I, Tew G, Beard JD, Saxton JM. Nordic Poles Immediately Improve Walking Distance in Patients with Intermittent Claudication. *Eur J Vasc Endovasc Surg*. 2008;36(6):689-694.
11. Barberan-Garcia A, Arbillaga-Etxarri A, Gimeno-Santos E, et al. Nordic walking enhances oxygen uptake without increasing the rate of perceived exertion in patients with chronic obstructive pulmonary disease. *Respiration*. 2015;89(3):221-225.
12. Breyer M-K, Breyer-Kohansal R, Funk G-C, et al. Nordic walking improves daily physical activities in COPD: a randomised controlled trial. *Respir Res*. 2010;11(1):112.
13. Bieler T, Siersma V, Magnusson SP, Kjaer M, Christensen HE, Beyer N. In hip osteoarthritis, Nordic Walking is superior to strength training and home-based exercise for improving function. *Scand J Med Sci Sports*. 2016.
14. Monteiro EP, Franzoni LT, Cubillos DM, et al. Effects of Nordic walking training on functional parameters in Parkinson's disease: a randomized controlled clinical trial. *Scand J Med Sci Sports*. 2016. [Epub ahead of print]
15. Cugusi L, Solla P, Serpe R, et al. Effects of a Nordic Walking program on motor and non-motor symptoms, functional performance and body composition in patients with Parkinson's disease. *NeuroRehabilitation*. 2015;37(2):245-254.
16. van Eijkeren FJM, Reijmers RSJ, Kleinveld MJ, Minten A, ter Brugge JP, Bloem BR. Nordic walking improves mobility in Parkinson's disease. *Mov Disord*. 2008;23(15):2239-2243.
17. Skórkowska-Telichowska K, Kropielnicka K, Buliska K, et al. Nordic walking in the second half of life. *Aging Clin Exp Res*. 2016. [Epub ahead of print]
18. Alves da Rocha P, McClelland J, Morris M. Complementary physical therapies for movement disorders in parkinson's disease: a systematic review. *Eur J Phys Rehabil Med*. 2015;51(6):693-704
19. Fritz B, Rombach S, Godau J, Berg D, Horstmann T, Grau S. The influence of Nordic Walking training on sit-to-stand transfer in Parkinson patients. *Gait Posture*. 2011;34(2):234-238.
20. Reuter I, Mehnert S, Leone P, Kaps M, Oechsner M, Engelhardt M. Effects of a flexibility and relaxation programme, walking, and nordic walking on Parkinson's disease. *J Aging Res*. 2011;2011:232473.
21. Hagner-Derengowska M, Kaluzny K, Kochanski B, et al. Effects of Nordic Walking and Pilates exercise programs on blood glucose and lipid profile in overweight and obese postmenopausal women in an experimental, nonrandomized, open-label, prospective controlled trial. *Menopause*. 2015;22(11):1215-1223.
22. Hagner-Derengowska M, Kaluzny K, Hagner W, et al. The influence of a ten-week Nordic walking training-rehabilitation program on the level of lipids in blood in overweight and obese postmenopausal women. *J Phys Ther Sci*. 2015;27(10):3039-3044.
23. Figard-Fabre H, Fabre N, Leonardi A, Schena F. Efficacy of nordic walking in obesity management. *Int J Sports Med*. 2011;32(6):407-414.
24. Fritz T, Caidahl K, Osler M, Ostenson CG, Zierath JR, Wändell P. Effects of Nordic walking on health-related quality of life in overweight individuals with type 2 diabetes mellitus, impaired or normal glucose tolerance. *Diabet Med*. 2011;28(11):1362-1372.
25. Gram B, Christensen R, Christiansen C, Gram J. Effects of nordic walking and exercise in type 2 diabetes mellitus: A randomized controlled trial. *Clin J Sport Med*. 2010;20(5):355-361.
26. Park SD, Yu SH. The effects of Nordic and general walking on depression disorder patients' depression, sleep, and body composition. *J Phys Ther Sci*. 2015;27(8):2481-2485.

27. Lee HS, Park JH. Effects of Nordic walking on physical functions and depression in frail people aged 70 years and above. *J Phys Ther Sci*. 2015;27(8):2453-2456.
28. Suija K, Pechter U, Kalda R, Tähepõld H, Maaros J, Maaros H-I. Physical activity of depressed patients and their motivation to exercise: Nordic Walking in family practice. *Int J Rehabil Res*. 2009;32(2):132-138.
29. Hartvigsen J, Morsø L, Bendix T, Manniche C. Supervised and non-supervised Nordic walking in the treatment of chronic low back pain: A single blind randomized clinical trial. *BMC Musculoskelet Disord*. 2010;11.
30. Park H-S, Lee S-N, Sung D-H, Choi H-S, Kwon TD, Park GD. The Effect of Power Nordic Walking on Spine Deformation and Visual Analog Pain Scale in Elderly Women with Low Back Pain. *J Phys Ther Sci*. 2014;26(11):1809-1812.
31. Jones KD. Nordic walking in fibromyalgia: a means of promoting fitness that is easy for busy clinicians to recommend. *Arthritis Res Ther*. 2011;13(1):103.
32. Mannerkorpi K, Nordeman L, Cider A, Jonsson G. Does moderate-to-high intensity Nordic walking improve functional capacity and pain in fibromyalgia? A prospective randomized controlled trial. *Arthritis Res Ther*. 2010;12(5):R189.
33. Herfurth M, Godau J, Kattner B, et al. Gait velocity and step length at baseline predict outcome of Nordic walking training in patients with Parkinson's disease. *Parkinsonism Relat Disord*. 2015;21(4):413-416.
34. Mikalacki M, Cokorilo N, Katic R. Effect of nordic walking on functional ability and blood pressure in elderly women. *Coll Antropol*. 2011;35(3):889-894.
35. Jasiski R, Socha M, Sitko L, Kubicka K, Woniewski M, Sobiech KA. Effect of nordic walking and water aerobics training on body composition and the blood flow in lower extremities in elderly women. *J Hum Kinet*. 2015;45(1):113-122.
36. Latosik E, Zubrzycki IZ, Ossowski Z, et al. Physiological Responses Associated with Nordic-walking training in Systolic Hypertensive Postmenopausal Women. *J Hum Kinet*. 2014;43(November):185-190.
37. Hagner-Derengowska M, Kałuzny K, Hagner W, et al. The effect of a 10-week Nordic walking training program on the level of GH and LH in elderly women. *Climacteric*. 2015;18(6):835-840.
38. O'Donovan R, Kennedy N. "Four legs instead of two"--perspectives on a Nordic walking-based walking programme among people with arthritis. *Disabil Rehabil*. 2015;37(18):1635-1642.
39. Song MS, Yoo YK, Choi CH, Kim NC. Effects of nordic walking on body composition, muscle strength, and lipid profile in elderly women. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2013;7(1):1-7.
40. Korsten-Reck U. [Physical activity in pregnancy and in breast-feeding period in obese mothers]. *Z Geburtshilfe Neonatol*. 2010;214(3):95-102.
41. Kocur P, Deskur-Smielecka E, Wilk M, Dylewicz P. Effects of Nordic Walking training on exercise capacity and fitness in men participating in early, short-term inpatient cardiac rehabilitation after an acute coronary syndrome -- a controlled trial. *Clin Rehabil*. 2009;23(11):995-1004.
42. Kang T-W, Lee J-H, Cynn H-S. Six-Week Nordic Treadmill Training Compared with Treadmill Training on Balance, Gait, and Activities of Daily Living for Stroke Patients: A Randomized Controlled Trial. *J Stroke Cerebrovasc Dis*. 2016;25(4):848-856.
43. Jönsson C, Johansson K. Pole walking for patients with breast cancer-related arm lymphedema. *Physiother Theory Pract*. 2009;25(3):165-173.
44. Rybicki JR, Leszczynska-Bolewska BM, Grochulska WE, et al. Oxygen uptake during Nordic walking training in patients rehabilitated after coronary events. *Kardiol Pol*. 2015;73(1):17-23.
45. Dominelli PB, Sheel AW, Foster GE. Effect of carrying a weighted backpack on lung mechanics during treadmill walking in healthy men. *Eur J Appl Physiol*. 2012;112(6):2001-2012.
46. Li JX, Hong Y, Robinson PD. The effect of load carriage on movement kinematics and respiratory parameters in children during walking. *Eur J Appl Physiol*. 2003;90(1-2):35-43.
47. Chow DHK, Leung KTY, Holmes a D. Changes in spinal curvature and proprioception of schoolboys carrying different weights of backpack. *Ergonomics*. 2007;50(12):2148-2156.
48. Devroey C, Jonkers I, de Becker A, Lenaerts G, Spaepen A. Evaluation of the effect of backpack load and position during standing and walking using biomechanical, physiological and subjective measures. *Ergonomics*. 2007;50(5):728-742.
49. Stuempfle KJ, Drury DG, Wilson AL. Effect of load position on physiological and perceptual responses during load carriage with an internal frame backpack. *Ergonomics*. 2004;47(7):784-789.
50. Golriz S, Hebert JJ, Foreman KB, Walker BF. The effect of hip belt use and load placement in a backpack on postural stability and perceived exertion: a within-subjects trial. *Ergonomics*. 2015;58(1):140-147.
51. Sharpe SR, Holt KG, Saltzman E, Wagenaar RC. Effects of a hip belt on transverse plane trunk coordination and stability during load carriage. *J Biomech*. 2008;41(5):968-976.
52. Legg SJ, Cruz CO. Effect of single and double strap backpacks on lung function. *Ergonomics*. 2004;47(3):318-323.
53. Schiffer T, Knicker A, Dannöhl R, Strüder HK. Energy cost and pole forces during nordic walking under different surface conditions. *Med Sci Sports Exerc*. 2009;41(3):663-668.
54. Hansen EA, Smith G. Energy expenditure and comfort during nordic walking with different pole lengths. *J Strength Cond Res*. 2009;23(4):1187-1194.

55. Schiffer T, Knicker A, Montanarella M, Strüder HK. Mechanical and physiological effects of varying pole weights during Nordic walking compared to walking. *Eur J Appl Physiol.* 2011;111(6):1121-1126.
56. Hagen M, Hennig EM, Stieldorf P. Lower and upper extremity loading in nordic walking in comparison with walking and running. *J Appl Biomech.* 2011;27(1):22-31.
57. Wulf G, Prinz W. Directing attention to movement effects enhances learning: a review. *Psychon Bull Rev.* 2001;8(4):648-660.
58. Wulf G, Höß M, Prinz W. Instructions for motor learning: differential effects of internal versus external focus of attention. *J Mot Behav.* 1998;30(October 2012):169-179.
59. Wulf G, McConnel N, Gartner M, Schwarz A. Enhancing the learning of sports skills through external-focus feedback. *J Mot Behav.* 2002;34(2):171-182.
60. Nakakubo S, Doi T, Sawa R, Misu S, Tsutsumimoto K, Ono R. Does arm swing emphasized deliberately increase the trunk stability during walking in the elderly adults? *Gait Posture.* 2014;40(4):516-520.
61. Bruijn SM, Meijer OG, Beek PJ, van Dieën JH. The effects of arm swing on human gait stability. *J Exp Biol.* 2010;213(Pt 23):3945-3952.
62. Ortega JD, Fehlman LA, Farley CT. Effects of aging and arm swing on the metabolic cost of stability in human walking. *J Biomech.* 2008;41(16):3303-3308.
63. Foissac MJ, Berthollet R, Seux J, Belli A, Millet GY. Effects of hiking pole inertia on energy and muscular costs during uphill walking. *Med Sci Sports Exerc.* 2008;40(6):1117-1125.
64. Pellegrini B, Peyré-Tartaruga LA, Zoppirolli C, et al. Exploring muscle activation during nordic walking: A comparison between conventional and uphill walking. *PLoS One.* 2015;10(9).
65. Meyns P, Bruijn SM, Duysens J. The how and why of arm swing during human walking. *Gait Posture.* 2013;38(4):555-562.
66. Li Y, Wang W, Crompton RH, Gunther MM. Free vertical moments and transverse forces in human walking and their role in relation to arm-swing. *J Exp Biol.* 2001;204(Pt 1):47-58.
67. Hof AL. The equations of motion for a standing human reveal three mechanisms for balance. *J Biomech.* 2007;40(2):451-457.
68. Umberger BR. Effects of suppressing arm swing on kinematics, kinetics, and energetics of human walking. *J Biomech.* 2008;41(11):2575-2580.
69. Yizhar Z, Boulos S, Inbar O, Carmeli E. The effect of restricted arm swing on energy expenditure in healthy men. *Int J Rehabil Res.* 2009;32(2):115-123.
70. Dietz V. Do human bipeds use quadrupedal coordination? *Trends Neurosci.* 2002;25(9):462-467.
71. Forssberg H, Grillner S, Halbertsma J, Rossignol S. The locomotion of the low spinal cat. II. Interlimb coordination. *Acta Physiol Scand.* 1980;108(3):283-295.
72. Balter JE, Zehr EP. Neural coupling between the arms and legs during rhythmic locomotor-like cycling movement. *J Neurophysiol.* 2007;97(2):1809-1818.
73. Dietz V, Fouad K, Bastiaanse CM. Neuronal coordination of arm and leg movements during human locomotion. *Eur J Neurosci.* 2001;14(11):1906-1914.
74. Zehr EP, Duysens J. Regulation of arm and leg movement during human locomotion. *Neuroscientist.* 2004;10(4):347-361.
75. Massaad F, Levin O, Meyns P, Drijkoningen D, Swinnen SP, Duysens J. Arm sway holds sway: locomotor-like modulation of leg reflexes when arms swing in alternation. *Neuroscience.* 2014;258:34-46.
76. Hundza SR, de Ruiter GC, Klimstra M, Zehr EP. Effect of afferent feedback and central motor commands on Soleus H-reflex suppression during arm cycling. *J Neurophysiol.* 2012;(September 2012):3049-3058.
77. Miaszek JE. The H-reflex as a tool in neurophysiology: Its limitations and uses in understanding nervous system function. *Muscle and Nerve.* 2003;28(2):144-160.
78. Hennig EM, Rosenbaum D, Milani TL. Transfer of tennis racket vibrations onto the human forearm. *Med Sci Sports Exerc.* 1992;24:1134-1140.
79. Hagen M, Verhülsdonk S, Hennig EM. In G.P. Brüggemann & A. Niehoff (Eds.), Deutscher Kongress für Biomechanik 2007 - Programm und Tagungsband. In: *Aus- Wirkungen Verschiedener Stockkonstruktionen Auf Die Belastung Der Oberen Extremitäten Und Die Vortrieb-Swirkung Beim Nordic Walking.* 2007:149.
80. Hansen L, Henriksen M, Larsen P, Alkjaer T. Nordic Walking does not reduce the loading of the knee joint. *Scand J Med Sci Sport.* 2008;18(4):436-441.
81. Stief F, Kleindienst FI, Wiemeyer J, Wedel F, Campe S, Krabbe B. Inverse dynamic analysis of the lower extremities during nordic walking, walking, and running. *J Appl Biomech.* 2008;24(4):351-359.
82. Dziuba AK, Zurek G, Garrard I, Wierzbicka-Damska I. Biomechanical parameters in lower limbs during natural walking and Nordic walking at different speeds. *Acta Bioeng Biomech / Wroclaw Univ Technol.* 2015;17(1):95-101.
83. Jensen SB, Henriksen M, Aaboe J, Hansen L, Simonsen EB, Alkjaer T. Is it possible to reduce the knee joint compression force during level walking with hiking poles? *Scand J Med Sci Sport.* 2011;21(6).
84. Willson J, Torry MR, Decker MJ, Kernozek T, Steadman JR. Effects of walking poles on lower extremity gait mechanics. *Med Sci Sports Exerc.* 2001;33(1):142-147.
85. Encarnación-Martínez A, Pérez-Soriano P, Llana-Belloch S. Differences in ground reaction forces and shock impacts between nordic walking and walking. *Res Q Exerc Sport.* 2015;86(1):94-99.
86. Kleindienst FI, Michel KJ, Schwarz J, Krabbe B. [Comparison of kinematic and kinetic parameters between the locomotion patterns in nordic walking, walking and running]. *Sportverletz Sportschaden.* 2006;20(1):25-30.

87. Collins EG, O'connell S, McBurney C, et al. Comparison of walking with poles and traditional walking for peripheral arterial disease rehabilitation. *J Cardiopulm Rehabil Prev.* 32(4):210-218.
88. Zoffoli L, Lucertini F, Federici A, Ditroilo M. Trunk muscles activation during pole walking vs. walking performed at different speeds and grades. *Gait Posture.* 2016;46:57-62.
89. Shim J-M, Kwon H-Y, Kim H-R, Kim B-I, Jung J-H. Comparison of the Effects of Walking with and without Nordic Pole on Upper Extremity and Lower Extremity Muscle Activation. *J Phys Ther Sci.* 2013;25(12):1553-1556.
90. Schiffer T, Knicker A, Hoffman U, Harwig B, Hollmann W, Strüder HK. Physiological responses to nordic walking, walking and jogging. *Eur J Appl Physiol.* 2006;98(1):56-61.
91. Sugiyama K, Kawamura M, Tomita H, Katamoto S. Oxygen uptake, heart rate, perceived exertion, and integrated electromyogram of the lower and upper extremities during level and Nordic walking on a treadmill. *J Physiol Anthropol.* 2013;32(1):2.
92. Fritschi JO, van Uffelen JGZ, Brown WJ. Pole walking down-under: profile of pole walking leaders, walkers and programs in Australia and factors relating to participation. *Health Promot J Austr.* 2014;25(3):215-221.
93. Saarela M V, Hari R. Listening to humans walking together activates the social brain circuitry. *Soc Neurosci.* 2008;3(3-4):401-409.
94. Denoth J. Load on the locomotor system and modelling. In: Nigg BM, ed. Biomechanics of Running Shoes. *Human Kine.* Champaign, IL; 1986:63-116.
95. Kocur P, Wiernicka M, Wilski M, et al. Does Nordic walking improves the postural control and gait parameters of women between the age 65 and 74: a randomized trial. *J Phys Ther Sci.* 2015;27(12):3733-3737.
96. Julius LM, Brach JS, Wert DM, VanSwearingen JM. Perceived Effort of Walking: Relationship With Gait, Physical Function and Activity, Fear of Falling, and Confidence in Walking in Older Adults With Mobility Limitations. *Phys Ther.* 2012;92(10):1268-1277.
97. Summary of the Updated American Geriatrics Society/ British Geriatrics Society clinical practice guideline for prevention of falls in older persons. *J Am Geriatr Soc.* 2011;59(1):148-157.
98. Parkatti T, Perttunen J, Wacker P. Improvements in functional capacity from Nordic walking: a randomized-controlled trial among elderly people. *J Aging Phys Act.* 2012;20(1):93-105.
99. Figueiredo S, Finch L, Mai J, Ahmed S, Huang A, Mayo NE. Nordic walking for geriatric rehabilitation: a randomized pilot trial. *Disabil Rehabil.* 2012;35(July 2012):1-8.
100. Allison DB, Williams MS, Hand GA, Jakicic JM, Fontaine KR. Conclusion of "Nordic walking for geriatric rehabilitation: a randomized pilot trial" is based on faulty statistical analysis and is inaccurate. *Disabil Rehabil.* 2015;37(18):1692-1693.
101. Dalton C, Nantel J. Nordic Walking Improves Postural Alignment and Leads to a More Normal Gait Pattern Following 8 Weeks of Training: A Pilot Study. *J Aging Phys Act.* 2016.[Epub ahead of print]
102. Virág A, Karóczy CK, Jakab A, Vass Z, Kovács E, Gondos T. Short-term and long-term effects of Nordic Walking training on balance, functional mobility, muscle strength and aerobic endurance among Hungarian community-living older people: a feasibility study. *J Sports Med Phys Fitness.* 2014.55(11):1285-92
103. Church TS, Earnest CP, Morss GM. Field testing of physiological responses associated with Nordic Walking. *Res Q Exerc Sport.* 2002;73(3):296-300.
104. Porcari JP, Hendrickson TL, Walter PR, Terry L, Walsko G. The physiological responses to walking awith and without Power Poles on treadmill exercise. *Res Q Exerc Sport.* 1997;68(2):161-166.
105. Figard-Fabre H, Fabre N, Leonardi A, Schena F. Physiological and perceptual responses to Nordic walking in obese middle-aged women in comparison with the normal walk. *Eur J Appl Physiol.* 2010;108(6):1141-1151.
106. Perrey S, Fabre N. Exertion during uphill, level and downhill walking with and without hiking poles. *J Sport Sci Med.* 2008;7(1):32-38.
107. Fritz T, Caidahl K, Krook A, et al. Effects of Nordic walking on cardiovascular risk factors in overweight individuals with type 2 diabetes, impaired or normal glucose tolerance. *Diabetes Metab Res Rev.* 2013;29(1):25-32.
108. Erickson KI, Voss MW, Prakash RS, et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci U S A.* 2011;108(7):3017-3022.
109. Inoue K, Hanaoka Y, Nishijima T, et al. Long-term mild exercise training enhances hippocampus-dependent memory in rats. *Int J Sports Med.* 2015;36(4):280-285.
110. Erickson KI, Raji CA, Lopez OL, et al. Physical activity predicts gray matter volume in late adulthood: The Cardiovascular Health Study. *Neurology.* 2010;75(16):1415-1422.
111. Erickson KI, Prakash RS, Voss MW, et al. Aerobic fitness is associated with hippocampal volume in elderly humans. *Hippocampus.* 2009;19(10):1030-1039.
112. Cotman CW, Berchtold NC. Physical activity and the maintenance of cognition: Learning from animal models. *Alzheimer's Dement.* 2007;3(2 SUPPL.).
113. Van Praag H. Neurogenesis and exercise: Past and future directions. *NeuroMolecular Med.* 2008;10(2):128-140.
114. Varma VR, Chuang YF, Harris GC, Tan EJ, Carlson MC. Low-intensity daily walking activity is associated with hippocampal volume in older adults. *Hippocampus.* 2015;25(5):605-615.
115. van Praag H, Kempermann G, Gage FH. Neural consequences of environmental enrichment. *Nat Rev Neurosci.* 2000;1(December):191-198.