

REFERENCES

1. Falla D, Bilenkij G, Jull G. Patients with chronic neck pain demonstrate altered patterns of muscle activation during performance of a functional upper limb task. *Spine* (Phila Pa 1976). 2004;29:1436-1440.
2. Gogia PP, Sabbahi MA. Electromyographic analysis of neck muscle fatigue in patients with osteoarthritis of the cervical spine. *Spine* (Phila Pa 1976). 1994;19:502-506.
3. Rosenfeld M, Gunnarsson R, Borenstein P. Early Intervention in Whiplash-Associated Disorders: A Comparison of Two Treatment Protocols. *Spine* (Phila Pa 1976). 2000;14:1782-7.
4. Jull G, Trott P, Potter H, et al. A Randomized Controlled Trial of Exercise and Manipulative Therapy for Cervicogenic Headache. *Spine* (Phila Pa 1976). 2002;27:1835-43.
5. Watson DH, Trott PH. Cervical headache: An investigation of natural head posture and upper cervical flexor muscle performance. *Cephalalgia*. 1993;13:272-284.
6. Jull G, Barrett C, Magee R, Ho P. Further clinical clarification of the muscle dysfunction in cervical headache. *Cephalalgia*. 1999;19:179-185.
7. Jull GA. Deep cervical flexor muscle dysfunction in whiplash. *J Musculoskelet Pain*. 2000;8:143-154.
8. Falla DL, Jull GA, Hodges PW. Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. *Spine* (Phila Pa 1976). 2004;29(19):2108-2114.
9. Rosenfeld M, Seferiadis A, Carlsson J, Gunnarsson R. Active Intervention in Patients with Whiplash-Associated Disorders Improves Long-Term Prognosis: A Randomized Controlled Clinical Trial. *Spine* (Phila Pa 1976) 2003;28:2491-8
10. Mealy K, Brennan H, Fenelon GC. Early Mobilization of Acute Whiplash Injuries. *Br Med J (Clin Res Ed)*. 1986;292:656-7
11. Crawford JR, Khan RJ, Varley GW. Early management and outcome following soft tissue injuries of the neck—a randomised controlled trial. *Injury*. 2004;35:891-895.
12. McKinney LA. Early Mobilisation and Outcome in Acute Sprains of the Neck. *BMJ* 1989;299:1006-8
13. Salter RB. The biologic concept of continuous passive motion of synovial joints. The first 18 years of basic research and its clinical application. *Clin Orthop Relat Res*. 1989;12-25.
14. Van Oosterwijck J, Nijs J, Meeus M, et al. Pain neurophysiology education improves cognitions, pain thresholds, and movement performance in people with chronic whiplash: a pilot study. *J Rehabil Res Dev*. 2011;48(1):43-58.
15. Wang Y, Wang JY, Luo F. Why self-induced pain feels less painful than externally generated pain: Distinct brain activation patterns in self- and externally generated pain. *PLoS One*. 2011;6.
16. Ploghaus A, Becerra L, Borras C, Borsook D. Neural circuitry underlying pain modulation: Expectation, hypnosis, placebo. *Trends Cogn Sci*. 2003;7:197-200.
17. Jull GA, Falla D, Vicenzino B, Hodges PW. The effect of therapeutic exercise on activation of the deep cervical flexor muscles in people with chronic neck pain. *Man Ther*. 2009;14:696-701.
18. Jordan A, Mehlsen J, Ostergaard K. A comparison of physical characteristics between patients seeking treatment for neck pain and age-matched healthy people. *J Manipulative Physiol Ther*. 1997;20:468-475.
19. Silverman JL, Rodriguez AA, Agre JC. Quantitative cervical flexor strength in healthy subjects and in subjects with mechanical neck pain. *Arch Phys Med Rehabil*. 1991;72:679-681.
20. Barton PM, Hayes KC. Neck flexor muscle strength, efficiency, and relaxation times in normal subjects and subjects with unilateral neck pain and headache. *Arch Phys Med Rehabil*. 1996;77:680-687.
21. Jull G, Falla D, Treleaven J, Hodges P, Vicenzino B. Retraining cervical joint position sense: The effect of two exercise regimes. *J Orthop Res*. 2007;25:404-412.
22. Lluch E, Schomacher J, Gizzi L, Petzke F, Seegar D, Falla D. Immediate effects of active cranio-cervical flexion exercise versus passive mobilisation of the upper cervical spine on pain and performance on the cranio-cervical flexion test. *Man Ther*. 2014;19:25-31.
23. O'Leary S, Falla D, Hodges PW, Jull G, Vicenzino B. Specific Therapeutic Exercise of the Neck Induces Immediate Local Hypoalgesia. *J Pain*. 2007;8:832-839.
24. Wenngren BI, Pedersen J, Sjölander P, Bergenheim M, Johansson H. Bradykinin and muscle stretch alter contralateral cat neck muscle spindle output. *Neurosci Res*. 1998;32:119-129.
25. Loudon JK, Ruhl M, Field E. Ability to reproduce head position after whiplash injury. *Spine* (Phila Pa 1976). 1997;22:865-868.
26. Heikkilä H V., Wenngren BI. Cervicocephalic kinesthetic sensibility, active range of cervical motion, and oculomotor function in patients with whiplash injury. *Arch Phys Med Rehabil*. 1998;79:1089-1094.
27. Szeto GPY, Straker LM, O'Sullivan PB. A comparison of symptomatic and asymptomatic office workers performing monotonous keyboard work - 2: Neck and shoulder kinematics. *Man Ther*. 2005;10:281-291.

28. Hiengkaew V, Panichaporn W, Thanungkul S. Postural balance, visual verticality perception, and its association in individuals with and without neck pain. *J Med Assoc Thai*. 2014;97 Suppl 7:S70-S74.
29. Knox JJ, Beilstein DJ, Charles SD, et al. Changes in head and neck position have a greater effect on elbow joint position sense in people with whiplash-associated disorders. *Clin J Pain*. 22(6):512-518.
30. Sandlund J, Djupsjöbacka M, Ryhed B, Hamberg J, Björklund M. Predictive and discriminative value of shoulder proprioception tests for patients with whiplash-associated disorders. *J Rehabil Med*. 2006;38:44-49.
31. Cools AMJ, Struyf F, De Mey K, Maenhout A, Castelein B, Cagnie B. Rehabilitation of scapular dyskinesis: from the office worker to the elite overhead athlete. *Br J Sports Med*. 2014;48(8):692-697.
32. Helgadóttir H, Kristjánsson E, Mottram S, Karduna A, Jonsson H. Altered alignment of the shoulder girdle and cervical spine in patients with insidious onset neck pain and whiplash-associated disorder. *J Appl Biomech*. 2011;27(3):181-191.
33. Helgadóttir H, Kristjánsson E, Mottram S, Karduna AR, Jonsson H. Altered scapular orientation during arm elevation in patients with insidious onset neck pain and whiplash-associated disorder. *J Orthop Sports Phys Ther*. 2010;40(12):784-791.
34. Helgadóttir H, Kristjánsson E, Einarsson E, Karduna A, Jonsson H. Altered activity of the serratus anterior during unilateral arm elevation in patients with cervical disorders. *J Electromyogr Kinesiol*. 2011;21(6):947-953.
35. Zakharova-Luneva E, Jull G, Johnston V, O'Leary S. Altered trapezius muscle behavior in individuals with neck pain and clinical signs of scapular dysfunction. *J Manipulative Physiol Ther*. 2012;35(5):346-353.
36. Szeto GPY, Straker L, Raine S. A field comparison of neck and shoulder postures in symptomatic and asymptomatic office workers. *Appl Ergon*. 2002;33(1):75-84.
37. Wegner S, Jull G, O'Leary S, Johnston V. The effect of a scapular postural correction strategy on trapezius activity in patients with neck pain. *Man Ther*. 2010;15(6):562-566.
38. Weon JH, Oh JS, Cynn HS, Kim YW, Kwon OY, Yi CH. Influence of forward head posture on scapular upward rotators during isometric shoulder flexion. *J Bodyw Mov Ther*. 2010;14(4):367-374.
39. Hägg GM, Åström A. Load pattern and pressure pain threshold in the upper trapezius muscle and psychosocial factors in medical secretaries with and without shoulder/neck disorders. *Int Arch Occup Environ Health*. 1997;69(6):423-432.
40. Veiersted KB, Westgaard RH, Andersen P. Electromyographic evaluation of muscular work pattern as a predictor of trapezius myalgia. *Scand J Work Environ Heal*. 1993;19(4):284-290.
41. Larsen MK, Samani A, Madeleine P, Olsen HB, Søgaard K, Holtermann A. Short-term effects of implemented high intensity shoulder elevation during computer work. *BMC Musculoskelet Disord*. 2009;10:101.
42. Sjörs A, Larsson B, Dahlman J, Falkmer T, Gerdle B. Physiological responses to low-force work and psychosocial stress in women with chronic trapezius myalgia. *BMC Musculoskelet Disord*. 2009;10:63.
43. Staal JB, de Bie RA, Hendriks EJM. Aetiology and management of work-related upper extremity disorders. *Best Pract Res Clin Rheumatol*. 2007;21(1):123-133.
44. Visser B, Van Dieën JH. Pathophysiology of upper extremity muscle disorders. *J Electromyogr Kinesiol*. 2006;16(1):1-16.

REFERENCES

1. Falla D, Bilenkij G, Jull G. Patients with chronic neck pain demonstrate altered patterns of muscle activation during performance of a functional upper limb task. *Spine (Phila Pa 1976)*. 2004;29:1436-1440.
2. Gogia PP, Sabbahi MA. Electromyographic analysis of neck muscle fatigue in patients with osteoarthritis of the cervical spine. *Spine (Phila Pa 1976)*. 1994;19:502-506.
3. Rosenfeld M, Gunnarsson R, Borenstein P. Early Intervention in Whiplash-Associated Disorders: A Comparison of Two Treatment Protocols. *Spine (Phila Pa 1976)*. 2000 Jul 15;25(14):1782-7.
4. Jull G, Trott P, Potter H, et al. A Randomized Controlled Trial of Exercise and Manipulative Therapy for Cervicogenic Headache. *Spine (Phila Pa 1976)*. 2002 Sep 1;27(17):1835-43.:2002.
5. Watson DH, Trott PH. Cervical headache: An investigation of natural head posture and upper cervical flexor muscle performance. *Cephalalgia*. 1993;13:272-284.
6. Jull G, Barrett C, Magee R, Ho P. Further clinical clarification of the muscle dysfunction in cervical headache. *Cephalalgia*. 1999;19:179-185.
7. Jull GA. Deep cervical flexor muscle dysfunction in whiplash. *J Musculoskelet Pain*. 2000;8:143-154.
8. Falla DL, Jull GA, Hodges PW. Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. *Spine (Phila Pa 1976)*. 2004;29(19):2108-2114.
9. Falla D, Jull G, Hodges P, Vicenzino B. An endurance-strength training regime is effective in reducing myoelectric manifestations of cervical flexor muscle fatigue in females with chronic neck pain. *Clin Neurophysiol*. 2006;117:828-837.
10. Biondi DM. Cervicogenic headache: mechanisms, evaluation, and treatment strategies. *J Am Osteopath Assoc*. 2000;100:S7-S14.
11. Elliott J, Sterling M, Noteboom JT, Darnell R, Galloway G, Jull G. Fatty infiltrate in the cervical extensor muscles is not a feature of chronic, insidious-onset neck pain. *Clin Radiol*. 2008;63(6):681-687
12. Elliott J, Jull G, Noteboom JT, Darnell R, Galloway G, Gibbon WW. Fatty infiltration in the cervical extensor muscles in persistent whiplash-associated disorders: a magnetic resonance imaging analysis. *Spine (Phila Pa 1976)*. 2006;31(22):E847-E855.
13. Elliott J, Jull G, Noteboom J, Darnell R, Galloway G, Gibbon W. Fatty infiltration in the cervical extensor muscles in persistent whiplash-associated disorders: a magnetic resonance imaging analysis. *Spine (Phila Pa 1976)*. 2006;31(22):E847-E855.
14. Elliott J, Jull G, Noteboom JT, Galloway G. MRI study of the cross-sectional area for the cervical extensor musculature in patients with persistent whiplash associated disorders (WAD). *Man Ther*. 2008;13(3):258-265.
15. O'Leary S, Falla D, Elliott JM, Jull G. Muscle dysfunction in cervical spine pain: implications for assessment and management. *J Orthop Sports Phys Ther*. 2009;39(5):324-333.
16. Ylinen JJ, Savolainen S, Airaksinen O, Kautiainen H, Salo P, Häkkinen A. Decreased strength and mobility in patients after anterior cervical discectomy compared with healthy subjects. *Arch Phys Med Rehabil*. 2003;84(7):1043-1047.
17. Pettorossi VE, Schieppati M. Neck proprioception shapes body orientation and perception of motion. *Front Hum Neurosci*. 2014;8:895.
18. Jouffroy F. Evolution of the dorsal muscles of the spine in light of their adaptation to gravity effects. In: Berthoz A, Vidal P.P. GW, ed. *Head-Neck Sensory Motor System*. New York: Oxford University Press; 1992:22-35.
19. Ylinen J, Takala EP, Kautiainen H, et al. Association of neck pain, disability and neck pain during maximal effort with neck muscle strength and range of movement in women with chronic non-specific neck pain. *Eur J Pain*. 2004;8:473-478.
20. Curkovi B, Puksec M, Peri P, Babi-Nagli D. The effect of isometric exercises for cervical spine on blood pressure. *Reumatizam*. 1998;46:17-19.
21. Garcés GL, Medina D, Milutinovic L, Garavote P, Guerado E. Normative database of isometric cervical strength in a healthy population. *Med Sci Sports Exerc*. 2002;34:464-470.
22. Jordan A, Mehlsen J, Bülow PM, Ostergaard K, Danneskiold-Samsøe B. Maximal isometric strength of the cervical musculature in 100 healthy volunteers. *Spine (Phila Pa 1976)*. 1999;24:1343-1348.
23. Barton PM, Hayes KC. Neck flexor muscle strength, efficiency, and relaxation times in normal subjects and subjects with unilateral neck pain and headache. *Arch Phys Med Rehabil*. 1996;77:680-687.
24. Chiu TTW, Sing K Lo. Evaluation of cervical range of motion and isometric neck muscle strength: reliability and validity. *Clin Rehabil*. 2002;16:851-858.
25. Ylinen J, Salo P, Nykänen M, Kautiainen H, Häkkinen A. Decreased isometric neck strength in women with chronic neck pain and the repeatability of neck strength measurements. *Arch Phys Med Rehabil*. 2004;85:1303-1308.
26. Chiu TTW, Lam T-H, Hedley AJ. A randomized controlled trial on the efficacy of exercise for patients with chronic neck pain. *Spine (Phila Pa 1976)*. 2005;30(1):
27. Hutton RS, Atwater SW. Acute and chronic adaptations of muscle proprioceptors in response to increased use. *Sports Med*. 1992;14:406-421.