

REFERENCES

1. Retchford T, Crossley KM, Grimaldi A, Kemp JL, Cowan SM. Can local muscles augment stability in the hip? A narrative literature review. *J. Musculoskelet. Neuronal Interact.* 2013;13:1-12.
2. Philippon MJ, Decker MJ, Giphart JE, Torry MR, Wahoff MS, LaPrade RF. Rehabilitation exercise progression for the gluteus medius muscle with consideration for iliopsoas tendinitis: an in vivo electromyography study. *Am. J. Sports Med.* 2011;39:1777-1785.
3. Fredericson M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreicher N, Sahrmann SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin. J. Sport Med.* 2000;10:169-175.
4. MassoudArab A, RezaNourbakhsh M, Mohammadifar A. The relationship between hamstring length and gluteal muscle strength in individuals with sacroiliac joint dysfunction. *J. Man. Manip. Ther.* 2011;19:5-10.
5. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J. Orthop. Sports Phys. Ther.* 2010;40:42-51.
6. Powers CM. The influence of altered lower-extremity kinematics on patellofemoral joint dysfunction: a theoretical perspective. *J. Orthop. Sports Phys. Ther.* 2003;33:639-646.
7. Madureira MM, Takayama L, Gallinaro AL, Caparbo VF, Costa RA, Pereira RMR. Balance training program is highly effective in improving functional status and reducing the risk of falls in elderly women with osteoporosis: a randomized controlled trial. *Osteoporos. Int.* 2007;18:419-425.
8. Mohammad WS, Abdelraouf OR, Elhafez SM, Abdel-Aziem AA, Nassif NS. Isokinetic imbalance of hip muscles in soccer players with osteitis pubis. *J. Sports Sci.* 2014;32:934-9.
9. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J. Orthop. Sports Phys. Ther.* 2013;43(2):54-64.
10. Van Wingerden JP, Vleeming A, Buyruk HM, Raissadat K. Stabilization of the sacroiliac joint in vivo: Verification of muscular contribution to force closure of the pelvis. *Eur. Spine J.* 2004;13:199-205.
11. Hungerford B, Gilleard W, Hodges P. Evidence of altered lumbopelvic muscle recruitment in the presence of sacroiliac joint pain. *Spine (Phila. Pa. 1976).* 2003;28:1593-1600.
12. Okubo Y, Kaneoka K, Imai A, et al. Electromyographic analysis of transversus abdominis and lumbar multifidus using wire electrodes during lumbar stabilization exercises. *J. Orthop. Sports Phys. Ther.* 2010;40(11):743-50.
13. Blackburn SE, Portney LG. Electromyographic activity of back musculature during Williams' flexion exercises. *Phys. Ther.* 1981;61:878-885.
14. Richardson C, Jull G, Toppenburg R CM. Techniques for active lumbar stabilisation for spinal protection: A pilot study. *Aust. J. Physiother.* 1992;38:105-112.
15. Franz JR, Kram R. How does age affect leg muscle activity/coactivity during uphill and downhill walking? *Gait Posture* 2013;37(3):378-84.
16. McBeth JM, Earl-Boehm JE, Cobb SC, Huddleston WE. Hip muscle activity during 3 side-lying hip-strengthening exercises in distance runners. *J. Athl. Train.* 47(1):15-23.
17. Lee J, Cynn H, Choi S, Yoon T, Jeong H. Effects of different hip rotations on gluteus medius and tensor fasciae latae muscle activity during isometric side-lying hip abduction. *J. Sport Rehabil.* 2013;22:301-7.
18. Bolgla LA, Uhl TL. Electromyographic Analysis of Hip Rehabilitation Exercises in a Group of Healthy Subjects. *J Orthop Sports Phys Ther.* 2005; Aug; (8):487-494.
19. Arokoski MH, Arokoski JPA, Haara M, et al. Hip muscle strength and muscle cross sectional area in men with and without hip osteoarthritis. *J. Rheumatol.* 2002;29:2185-2195.
20. Rasch A, Byström AH, Dalen N, Berg HE. Reduced muscle radiological density, cross-sectional area, and strength of major hip and knee muscles in 22 patients with hip osteoarthritis. *Acta Orthop.* 2007;78:505-510.
21. Van der Krogt MM, Delp SL, Schwartz MH. How robust is human gait to muscle weakness? *Gait Posture* 2012;36:113-119.
22. Dwyer MK, Stafford K, Mattacola CG, Uhl TL, Giordani M. Comparison of gluteus medius muscle activity during functional tasks in individuals with and without osteoarthritis of the hip joint. *Clin. Biomech.* 2013;28:757-761.
23. Finley JM, Dhaher YY, Perreault EJ. Contributions of feed-forward and feedback strategies at the human ankle during control of unstable loads. *Exp. Brain Res.* 2012;217:53-66.
24. Dostal WF, Soderberg GL, Andrews JG. Actions of hip muscles. *Phys. Ther.* 1986;66:351-361.
25. Delp SL, Hess WE, Hungerford DS, Jones LC. Variation of rotation moment arms with hip flexion. *J. Biomech.* 1999;32:493-501.
26. Distefano LJ, Blackburn JT, Marshall SW, Padua DA. Gluteal muscle activation during common therapeutic exercises. *J. Orthop. Sports Phys. Ther.* 2009;39:532-540.
27. Roy RR, Kim JA, Monti RJ, Zhong H, Edgerton VR. Architectural and histochemical properties of cat hip "cuff" muscles. *Acta Anat. (Basel).* 1997;159:136-146.
28. Addison O, Young P, Inacio M, et al. Hip but Not Thigh Intramuscular Adipose Tissue is Associated with Poor Balance and Increased Temporal Gait Variability in Older Adults. *Curr. Aging Sci.* 2014.

29. Inacio M, Ryan AS, Bair W-N, Prettyman M, Beamer B a, Rogers MW. Gluteal muscle composition differentiates fallers from non-fallers in community dwelling older adults. *BMC Geriatr.* 2014;14:37.
30. Mansfield A, Peters AL, Liu BA, Maki BE. A perturbation-based balance training program for older adults: study protocol for a randomised controlled trial. *BMC Geriatr.* 2007;7:12.
31. Cyarto E V., Brown WJ, Marshall AL, Trost SG. Comparative effects of home- and group-based exercise on balance confidence and balance ability in older adults: Cluster randomized trial. *Gerontology* 2008;54:272-280.
32. Emery CA, Cassidy JD, Klassen TP, Rosychuk RJ, Rowe BH. Effectiveness of a home-based balance-training program in reducing sports-related injuries among healthy adolescents: A cluster randomized controlled trial. *CMAJ* 2005;172:749-754.
33. Emery CA, Rose MS, McAllister JR, Meeuwisse WH. A prevention strategy to reduce the incidence of injury in high school basketball: a cluster randomized controlled trial. *Clin J Sport Med* 2007;17:17-24.
34. Bolton DAE, Misiaszek JE. Effects of Ankle Extensor Muscle Afferent Inputs on Hip Abductor and Adductor Activity in the Decerebrate Walking Cat. *J. Neurophysiol.* 2012.
35. Rodacki ALF, Souza RM, Ugrinowitsch C, Cristopoliski F, Fowler NE. Transient effects of stretching exercises on gait parameters of elderly women. *Man. Ther.* 2009;14:167-172.
36. Cristopoliski F, Barela JA, Leite N, Fowler NE, Rodacki ALF. Stretching exercise program improves gait in the elderly. *Gerontology* 2009;55:614-620.
37. Kerrigan DC, Xenopoulos-Oddsson A, Sullivan MJ, Lelas JJ, Riley PO. Effect of a Hip Flexor-Stretching Program on Gait in the Elderly. *PM R* 2003; 3 (4):330-5
38. Watt JR, Jackson K, Franz JR, Dicharry J, Evans J, Kerrigan DC. Effect of a supervised hip flexor stretching program on gait in elderly individuals. *PM R* 2011;3:324-329.

REFERENCES

1. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther.* 2013;43(2):54-64.
2. Bass CJ, Connell D a. Sonographic findings of tensor fascia lata tendinopathy: another cause of anterior groin pain. *Skeletal Radiol.* 2002;31(3):143-148.
3. Webster KA, Gribble PA. A comparison of electromyography of gluteus medius and maximus in subjects with and without chronic ankle instability during two functional exercises. *Phys Ther Sport.* 2013;14(1):17-22.
4. Kwak SD, Ahmad CS, Gardner TR, et al. Hamstrings and iliotibial band forces affect knee kinematics and contact pattern. *J Orthop Res.* 2000;18(1):101-108.
5. Merican a M, Amis a a. Anatomy of the lateral retinaculum of the knee. *J Bone Joint Surg Br.* 2008;90(4):527-534.
6. Merican AM, Amis A a. Iliotibial band tension affects patellofemoral and tibiofemoral kinematics. *J Biomech.* 2009;42(10):1539-1546.
7. Sutter R, Kalberer F, Binkert CA, Graf N, Pfirrmann CWA, Gutzzeit A. Abductor tendon tears are associated with hypertrophy of the tensor fasciae latae muscle. *Skeletal Radiol.* 2013;42(5):627-633.
8. Kingzett-Taylor A, Tirman PFJ, Feller J, et al. Tendinosis and tears of gluteus medius and minimus muscles as a cause of hip pain: MR imaging findings. *Am J Roentgenol.* 1999;173(4):1123-1126.
9. Grimaldi A, Richardson C, Durbridge G, Donnelly W, Darnell R, Hides J. The association between degenerative hip joint pathology and size of the gluteus maximus and tensor fascia lata muscles. *Man Ther.* 2009;14(6):611-617.
10. Dwyer MK, Stafford K, Mattacola CG, Uhl TL, Giordani M. Comparison of gluteus medius muscle activity during functional tasks in individuals with and without osteoarthritis of the hip joint. *Clin Biomech.* 2013;28(7):757-761.
11. Grimaldi A, Richardson C, Stanton W, Durbridge G, Donnelly W, Hides J. The association between degenerative hip joint pathology and size of the gluteus medius, gluteus minimus and piriformis muscles. *Man Ther.* 2009;14(6):605-610.
12. Cooper NA, Scavo KM, Strickland KJ, et al. Prevalence of gluteus medius weakness in people with chronic low back pain compared to healthy controls. *Eur Spine J.* 2015. 25(4):1258-65.
13. Simons DG, Travell JG. Myofascial origins of low back pain. 3. Pelvic and lower extremity muscles. *Postgr Med.* 1983;73(2):99-105, 108.
14. Njoo KH, Van der Does E. The occurrence and inter-rater reliability of myofascial trigger points in the quadratus lumborum and gluteus medius: a prospective study in non-specific low back pain patients and controls in general practice. *Pain.* 1994;58(3):317-323.
15. Bewyer KJ, Bewyer DC, Messenger D, Kennedy CM. Pilot data: association between gluteus medius weakness and low back pain during pregnancy. *Iowa Orthop J.* 2009;29:97-99.
16. Cooper NA, Scavo KM, Strickland KJ, et al. Prevalence of gluteus medius weakness in people with chronic low back pain compared to healthy controls. *Eur Spine J.* 2015:1-8.
17. Embaby E, Abdallah A. Trunk and Gluteus-Medius Muscles' Fatigability During Occupational Standing in Clinical Instructors with Low Back Pain. *Br J Sports Med.* 2013;47(10):e3-e3.
18. Kankaanpää M, Taimela S, Laaksonen D, Hänninen O, Airaksinen O. Back and hip extensor fatigability in chronic low back pain patients and controls. *Arch Phys Med Rehabil.* 1998;79(4):412-417.
19. Leinonen V, Kankaanpää M, Airaksinen O, Hänninen O. Back and hip extensor activities during trunk flexion/extension: Effects of low back pain and rehabilitation. *Arch Phys Med Rehabil.* 2000;81(1):32-37.
20. Lee SW, Kim SY, L S, K SY. Effects of hip exercises for chronic low-back pain patients with lumbar instability. *J Phys Ther Sci.* 2015;27(2):345-348.
21. Vleeming A, Stoeckart R, Volkens AC, Snijders CJ. Relation between form and function in the sacroiliac joint. Part I: Clinical anatomical aspects. *Spine (Phila Pa 1976).* 1990;15(2):130-132.
22. van Wingerden JP, Vleeming A, Buyruk HM, Raissadat K. Stabilization of the sacroiliac joint in vivo: verification of muscular contribution to force closure of the pelvis. *Eur Spine J.* 2004;13(3):199-205.
23. Vleeming A, Volkens AC, Snijders CJ, Stoeckart R. Relation between form and function in the sacroiliac joint. Part II: Biomechanical aspects. *Spine (Phila Pa 1976).* 1990;15(2):133-136.
24. Snijders CJ, Vleeming A, Stoeckart R. Transfer of lumbosacral load to iliac bones and legs Part 1: Biomechanics of self-bracing of the sacroiliac joints and its significance for treatment and exercise. *Clin Biomech (Bristol, Avon).* 1993;8(6):285-294.
25. Vleeming a., Van Wingerden JP, Snijders CJ, Stoeckart R, Stijnen T. Load application to the sacrotuberous ligament; influences on sacroiliac joint mechanics. *Clin Biomech.* 1989;4(4):204-209.
26. Vleeming A, Stoeckart R, Snijders CJ. The sacrotuberous ligament: a conceptual approach to its dynamic role in stabilizing the sacroiliac joint. *Clin Biomech.* 1989;4(4):201-203.

27. Van Wingerden JP, Vleeming A, Snijders CJ, Stoeckart R, Wingerden J, Vleeming A. A functional-anatomical approach to the spine-pelvis mechanism: interaction between the biceps femoris muscle and the sacrotuberous ligament. *Eur Spine J*. 1993;2(3):140-144.
28. Mooney V, Pozos R, Vleeming a, Gulick J, Swenski D. Exercise treatment for sacroiliac pain. *Orthopedics*. 2001;24(1):29-32.
29. Vleeming A, Pool-Goudzwaard AL, Hammudoghlu D, Stoeckart R, Snijders CJ, Mens JM. The function of the long dorsal sacroiliac ligament: its implication for understanding low back pain. *Spine (Phila Pa 1976)*. 1996;21(5):556-562.
30. Fortin JD, Aprill CN, Ponthieux B, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part II: Clinical evaluation. *Spine (Phila Pa 1976)*. 1994;19(13):1483-1489.
31. Fortin JD, Dwyer AP, West S, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part I: Asymptomatic volunteers. *Spine (Phila Pa 1976)*. 1994;19(13):1475-1482.
32. Fortin JD, Kissling RO, O'Connor BL, Vilensky JA. Sacroiliac joint innervation and pain. *Am J Orthop (Belle Mead NJ)*. 1999;28(12):687-690.
33. Schwarzer a C, Aprill CN, Bogduk N. The sacroiliac joint in chronic low back pain. *Spine (Phila Pa 1976)*. 1995;20(1):31-37.
34. Vleeming A, de Vries HJ, Mens JM a, van Wingerden J-P. Possible role of the long dorsal sacroiliac ligament in women with peripartum pelvic pain. *Acta Obstet Gynecol Scand*. 2002;81(5):430-436.
35. Ostgaard HC, Zetherstrom G, Roos-Hansson E. The posterior pelvic pain provocation test in pregnant women. *Eur Spine J*. 1994;3(5):258-260.
36. Mens JM a, Vleeming A, Snijders CJ, Stam HJ, Ginai AZ. The active straight leg raising test and mobility of the pelvic joints. *Eur Spine J*. 1999;8(6):468-474.
37. Jan M. A. Mens, Andry Vleeming CJS, Bart W. Koes, and Henk J. Stam M. Reliability and Validity of the Active Straight Leg Raise Test in Posterior Pelvic Pain Since Pregnancy. *Spine (Phila Pa 1976)*. 2001;26(10):1167-1171.
38. Mens JM a, Vleeming A, Snijders CJ, Koes BW, Stam HJ. Validity of the active straight leg raise test for measuring disease severity in patients with posterior pelvic pain after pregnancy. *Spine (Phila Pa 1976)*. 2002;27(2):196-200.
39. Hungerford B, Gilleard W, Hodges P. Evidence of altered lumbopelvic muscle recruitment in the presence of sacroiliac joint pain. *Spine (Phila Pa 1976)*. 2003;28(14):1593-1600.
40. Reiman MP, Bolgla L a, Loudon JK. A literature review of studies evaluating gluteus maximus and gluteus medius activation during rehabilitation exercises. *Physiother Theory Pract*. 2012;28(4):257-268.
41. O'Sullivan K, Smith SM, Sainsbury D. Electromyographic analysis of the three subdivisions of gluteus medius during weight-bearing exercises. *Sports Med Arthrosc Rehabil Ther Technol*. 2010;2:17.
42. Robinson RL, Nee RJ. Analysis of hip strength in females seeking physical therapy treatment for unilateral patellofemoral pain syndrome. *J Orthop Sport Phys Ther*. 2007;37(5):232-238.
43. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther*. 2010;40(2):42-51.
44. Powers CM, Ward SR, Fredericson M, Guillet M, Shellock FG. Patellofemoral kinematics during weight-bearing and non-weight-bearing knee extension in persons with lateral subluxation of the patella: a preliminary study. *J Orthop Sports Phys Ther*. 2003;33(11):677-685.
45. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther*. 2009;39(1):12-19.
46. Reiman MP, Bolgla L a, Lorenz D. Hip functions influence on knee dysfunction: a proximal link to a distal problem. *J Sport Rehabil*. 2009;18(1):33-46.
47. Mascal CL, Landel R, Powers C. Management of patellofemoral pain targeting hip, pelvis, and trunk muscle function: 2 case reports. *J Orthop Sports Phys Ther*. 2003;33(11):647-660.
48. Bolgla L a, Malone TR, Umberger BR, Uhl TL. Hip strength and hip and knee kinematics during stair descent in females with and without patellofemoral pain syndrome. *J Orthop Sports Phys Ther*. 2008;38(1):12-18.
49. Dolak KL, Silkman C, Medina McKeon J, Hosey RG, Lattermann C, Uhl TL. Hip strengthening prior to functional exercises reduces pain sooner than quadriceps strengthening in females with patellofemoral pain syndrome: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2011;41(8):560-570.
50. Nakagawa TH, Muniz TB, Baldon RDM, Dias Maciel C, de Menezes Reiff RB, Serrão FV. The effect of additional strengthening of hip abductor and lateral rotator muscles in patellofemoral pain syndrome: a randomized controlled pilot study. *Clin Rehabil*. 2008;22(12):1051-1060.
51. Ismail MM, Gamaleldein MH, Hassa KA. Closed Kinetic Chain exercises with or without additional hip strengthening exercises in management of Patellofemoral pain syndrome: A randomized controlled trial. *Eur J Phys Rehabil Med*. 2013;49(5):687-698.
52. Fukuda TY. Hip Posterolateral Musculature Strengthening in Sedentary Women With Patellofemoral Pain Syndrome: A Randomized Controlled Clinical Trial With 1-Year Follow-up. *J Orthop Sports Phys Ther*. 2012;42(10):823-830.
53. Ford KR, Nguyen A-D, Dischiavi SL, Hegedus EJ, Zuk EF, Taylor JB. An evidence-based review of hip-focused neuromuscular exercise interventions to address dynamic lower extremity valgus. *Open access J Sport Med*. 2015;6:291-303.

54. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33(4):492-501.
55. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: Part 1, mechanisms and risk factors. *Am J Sports Med.* 2006;34(2):299-311.
56. Myer GD, Ford KR, Di Stasi SL, Foss KDB, Micheli LJ, Hewett TE. High knee abduction moments are common risk factors for patellofemoral pain (PFP) and anterior cruciate ligament (ACL) injury in girls: is PFP itself a predictor for subsequent ACL injury? *Br J Sports Med.* 2015;49(2):118-122.
57. Ford KR, Myer GD, Hewett TE. Valgus knee motion during landing in high school female and male basketball players. *Med Sci Sports Exerc.* 2003;35(10):1745-1750. doi:10.1249/01.MSS.0000089346.85744.D9.
58. Ford KR, Myer GD, Toms HE, Hewett TE. Gender differences in the kinematics of unanticipated cutting in young athletes. *Med Sci Sports Exerc.* 2005;37(1):124-129.
59. Ford KR, Myer GD, Smith RL, Vianello RM, Seiwert SL, Hewett TE. A comparison of dynamic coronal plane excursion between matched male and female athletes when performing single leg landings. *Clin Biomech.* 2006;21(1):33-40.
60. Ford KR, Shapiro R, Myer GD, Van Den Bogert AJ, Hewett TE. Longitudinal sex differences during landing in knee abduction in young athletes. *Med Sci Sports Exerc.* 2010;42(10):1923-1931.
61. Malinzak RA, Colby SM, Kirkendall DT, Yu B, Garrett WE. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech.* 2001;16(5):438-445.
62. Hewett TE, Myer GD, Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Joint Surg Am.* 2004;86-A(8):1601-1608.
63. Chappell JD, Yu B, Kirkendall DT, Garrett WE. A comparison of knee kinetics between male and female recreational athletes in stop-jump tasks. *Am J Sports Med.* 2002;30(2):261-267.
64. McLean SG, Huang X, Su A, Van Den Bogert AJ. Sagittal plane biomechanics cannot injure the ACL during sidestep cutting. *Clin Biomech (Bristol, Avon).* 2004;19(8):828-838.
65. Kernozek TW, Torry MR, Van Hoof H, Cowley H, Tanner S. Gender differences in frontal and sagittal plane biomechanics during drop landings. *Med Sci Sports Exerc.* 2005;37(6):1003-1012.
66. Zeller BL, McCrory JL, Kibler W Ben, Uhl TL. Differences in kinematics and electromyographic activity between men and women during the single-legged squat. *Am J Sports Med.* 31(3):449-456.
67. Pappas E, Hagins M, Sheikhzadeh A, Nordin M, Rose D. Biomechanical differences between unilateral and bilateral landings from a jump: gender differences. *Clin J Sport Med.* 2007;17(4):263-268.
68. Hewett TE, Ford KR, Myer GD, Wanstrath K, Schepers M. Gender differences in hip adduction motion and torque during a single-leg agility maneuver. *J Orthop Res.* 2006;24(3):416-421.
69. Carson DW, Ford KR. Sex Differences in Knee Abduction During Landing: A Systematic Review. *Sport Heal A Multidiscip Approach.* 2011;3(4):373-382.
70. Noehren B, Davis I, Hamill J. ASB Clinical Biomechanics Award Winner 2006. Prospective study of the biomechanical factors associated with iliotibial band syndrome. *Clin Biomech.* 2007;22(9):951-956.
71. Earl JE, Hertel JN, Denegar CR. Patterns of Dynamic Malalignment, Muscle Activation, Joint Motion, and Patellofemoral-Pain Syndrome. *J Sport Rehabil.* 2005;14:216-234.
72. Ferber R, Noehren B, Hamill J, Davis IS. Competitive female runners with a history of iliotibial band syndrome demonstrate atypical hip and knee kinematics. *J Orthop Sports Phys Ther.* 2010;40(2):52-58.
73. Fredericson M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreich N, Sahrmann SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med.* 2000;10:169-175.
74. Fredericson M, Weir A. Practical management of iliotibial band friction syndrome in runners. *Clin J Sport Med.* 2006;16(3):261-268.
75. Fredericson M, Wolf C. Iliotibial band syndrome in runners: Innovations in treatment. *Sport Med.* 2005;35:451-459.
76. Beckman SM, Buchanan TS. Ankle inversion injury and hypermobility: Effect on hip and ankle muscle electromyography onset latency. *Arch Phys Med Rehabil.* 1995;76(12):1138-1143.
77. Presswood L, Cronin J, Keogh JWL, Whatman C. Gluteus Medius: Applied Anatomy, Dysfunction, Assessment, and Progressive Strengthening. *Strength Cond J.* 2008;30:41-53.
78. Lee D. Instability of the Sacroiliac Joint and the Consequences to Gait. *J Man Manip Ther.* 1996;4(1):22-29.
79. Presswood L, Cronin J, Keogh JWL, Whatman C. Gluteus Medius: Applied Anatomy, Dysfunction, Assessment, and Progressive Strengthening. *Strength Cond J.* 2008;30(5):41-53.
80. Blazeovich AJ. Optimizing Hip Musculature For Greater Sprint Running Speed. *Strength Cond J.* 2000;22(2):22.
81. Schache AG, Blanch PD, Dorn TW, Brown N a T, Rosemond D, Pandy MG. Effect of running speed on lower limb joint kinetics. *Med Sci Sports Exerc.* 2011;43(7):1260-1271.

82. Philippon MJ, Decker MJ, Giphart JE, Torry MR, Wahoff MS, LaPrade RF. Rehabilitation exercise progression for the gluteus medius muscle with consideration for iliopsoas tendinitis: an in vivo electromyography study. *Am J Sports Med.* 2011;39(8):1777-1785.
83. Robertson DGE, Wilson JMJ, St. Pierre T a. Lower extremity muscle functions during full squats. *J Appl Biomech.* 2008;24(4):333-339.
84. Hollman JH, Ginos BE, Kozuchowski J, Vaughn AS, Krause D a, Youdas JW. Relationships between knee valgus, hip-muscle strength, and hip-muscle recruitment during a single-limb step-down. *J Sport Rehabil.* 2009;18(1):104-117.
85. Hamstra-Wright KL, Huxel Bliven K. Effective exercises for targeting the gluteus medius. *J Sport Rehabil.* 2012;21(3):296-300.
86. Simenz CJ, Garceau LR, Lutsch BN, Suchomel TJ, Ebben WP. Electromyographical analysis of lower extremity muscle activation during variations of the loaded step-up exercise. *J Strength Cond Res.* 2012;26(12):3398-3405.
87. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther.* 2013;43:54-64.
88. McBeth JM, Earl-Boehm JE, Cobb SC, Huddleston WE. Hip muscle activity during 3 side-lying hip-strengthening exercises in distance runners. *J Athl Train.* 47(1):15-23.
89. Sidorkewicz N, Cambridge EDJ, McGill SM. Examining the effects of altering hip orientation on gluteus medius and tensor fasciae latae interplay during common non-weight-bearing hip rehabilitation exercises. *Clin Biomech (Bristol, Avon).* 2014;29(9):971-976.
90. Dostal WF, Soderberg GL, Andrews JG. Actions of hip muscles. *Phys Ther.* 1986;66:351-361.
91. Delp SL, Hess WE, Hungerford DS, Jones LC. Variation of rotation moment arms with hip flexion. *J Biomech.* 1999;32:493-501.
92. Retchford T, Crossley KM, Grimaldi A, Kemp JL, Cowan SM. Can local muscles augment stability in the hip? A narrative literature review. *J Musculoskelet Neuronal Interact.* 2013;13:1-12.
93. Distefano LJ, Blackburn JT, Marshall SW, Padua DA. Gluteal muscle activation during common therapeutic exercises. *J Orthop Sports Phys Ther.* 2009;39:532-540.
94. Gupta a, Fernihough B, Bailey G, Bombeck P, Clarke a, Hopper D. An evaluation of differences in hip external rotation strength and range of motion between female dancers and non-dancers. *Br J Sports Med.* 2004;38(6):778-783.
95. Bennell KL, Khan KM, Matthews BL, Singleton C. Changes in hip and ankle range of motion and hip muscle strength in 8–11 year old novice female ballet dancers and controls: a 12 month follow up study. *Br J Sport Med.* 2001;35:54-59.
96. Hamilton WG, Hamilton LH, Marshall P, Molnar M. A profile of the musculoskeletal characteristics of elite professional ballet dancers. *Am J Sports Med.* 1992;20(3):267-273.
97. Judge LW, Moreau C, Burke JR. Neural adaptations with sport-specific resistance training in highly skilled athletes. *J Sports Sci.* 2003;21(5):419-427.
98. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. Electromyographic analysis of gluteus medius and gluteus maximus during rehabilitation exercises. *Int J Sports Phys Ther.* 2011;6(3):206-223.
99. Ekstrom RA, Donatelli RA, Carp KC. Electromyographic analysis of core trunk, hip, and thigh muscles during 9 rehabilitation exercises. *J Orthop Sports Phys Ther.* 2007;37(12):754-762.
100. Rozzi SL, Lephart SM, Gear WS, Fu FH. Knee joint laxity and neuromuscular characteristics of male and female soccer and basketball players. *Am J Sports Med.* 1999;27(3):312-319.
101. Lewis CL, Sahrman SA. Muscle activation and movement patterns during prone hip extension exercise in women. *J Athl Train.* 2009;44(3):238-248.
102. Anders M. Glutes to the max. *ACE Fitness Matters.* 2006:7-9.
103. Bishop B, Greenstein J. EMG Activation of Gluteal Musculature During Exercises With and Without Resistance. *13th Annu TRAC Meet.* 2011
104. Lewis CL, Sahrman SA, Moran DW. Anterior hip joint force increases with hip extension, decreased gluteal force, or decreased iliopsoas force. *J Biomech.* 2007;40(16):3725-3731.
105. Shindle MK, Ranawat AS, Kelly BT. Diagnosis and Management of Traumatic and Atraumatic Hip Instability in the Athletic Patient. *Clin Sports Med.* 2006;25(2):309-326.
106. Janda V. On the concept of postural muscles and posture in man. *Aust J Physiother.* 1983;29(3):83-84.
107. Murphy DR, Byfield D, McCarthy P, Humphreys K, Gregory AA, Rochon R. Interexaminer Reliability of the Hip Extension Test for Suspected Impaired Motor Control of the Lumbar Spine. *J Manipulative Physiol Ther.* 2006;29(5):374-377.
108. Lehman GJ, Lennon D, Tresidder B, Rayfield B, Poschar M. Muscle recruitment patterns during the prone leg extension. *BMC Musculoskelet Disord.* 2004;5:3.
109. Lehman GJ. Trunk and hip muscle recruitment patterns during the prone leg extension following a lateral ankle sprain: a prospective case study pre and post injury. *Chiropr Osteopat.* 2006;14:4.
110. Prince F, Winter D, Stergiou P, Walt S. Anticipatory control of upper body balance during human locomotion. *Gait Posture.* 1994;2(1):19-25.
111. Berry JW, Lee TS, Foley HD, Lewis CL. Resisted Side Stepping: The Effect of Posture on Hip Abductor Muscle Activation. *J Orthop Sports Phys Ther.* 2015;45(9):675-682.

112. Cambridge EDJ, Sidorkewicz N, Ikeda DM, McGill SM. Progressive hip rehabilitation: The effects of resistance band placement on gluteal activation during two common exercises. *Clin Biomech*. 2012;27(7):719-724.
113. Youdas JW, Foley BM, Kruger BL, et al. Electromyographic analysis of trunk and hip muscles during resisted lateral band walking. *Physiother Theory Pr*. 2013;29(2):113-123.
114. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther*. 2010;40:42-51.
115. Toutoungi DE, Lu TW, Leardini A, Catani F, O'Connor JJ. Cruciate ligament forces in the human knee during rehabilitation exercises. *Clin Biomech (Bristol, Avon)*. 2000;15(3):176-187.
116. Begalle RL, DiStefano LJ, Blackburn T, Padua DA. Quadriceps and hamstrings coactivation during common therapeutic exercises. *J Athl Train*. 2012;47(4):396-405.
117. Winter DA. Human balance and posture control during standing and walking. *Gait Posture*. 1995;3(4):193-214.

REFERENCES

1. Willy RW, Scholz JP, Davis IS. Mirror gait retraining for the treatment of patellofemoral pain in female runners. *Clin Biomech* (Bristol, Avon). 2012;27(10):1045-1051.
2. Proske U, Morgan DL. Muscle damage from eccentric exercise: Mechanism, mechanical signs, adaptation and clinical applications. *J Physiol*. 2001;537(2):333-345.
3. McHugh MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand J Med Sci Sports*. 2003;13(2):88-97.
4. Macadam P, Cronin J, Contreras B. An examination of the gluteal muscle activity associated with dynamic hip abduction and hip external rotation exercise: a systematic review. *Int J Sports Phys Ther*. 2015;10(5):573-591.
5. Simenz CJ, Garceau LR, Lutsch BN, Suchomel TJ, Ebben WP. Electromyographical analysis of lower extremity muscle activation during variations of the loaded step-up exercise. *J Strength Cond Res*. 2012;26(12):3398-3405.
6. Reiman MP, Bolgla L a, Loudon JK. A literature review of studies evaluating gluteus maximus and gluteus medius activation during rehabilitation exercises. *Physiother Theory Pract*. 2012;28(4):257-268.
7. Ayotte NW, Stetts DM, Keenan G, Greenway EH. Electromyographical analysis of selected lower extremity muscles during 5 unilateral weight-bearing exercises. *J Orthop Sports Phys Ther*. 2007;37:48-55.
8. Mercer VS, Gross MT, Sharma S, Weeks E. Comparison of gluteus medius muscle electromyographic activity during forward and lateral step-up exercises in older adults. *Phys Ther*. 2009;89:1205-1214.
9. Park J, Denning WM, Pitt JD, Francom D, Hopkins JT, Seeley MK. Effects of Experimental Anterior Knee Pain on Muscle Activation During Landing and Jumping Performed at Various Intensities. *J Sport Rehabil*. 2015. Epub ahead of print]
10. MacAskill MJ, Durant TJS, Wallace DA. Gluteal muscle activity during weightbearing and non-weightbearing exercise. *Int J Sports Phys Ther*. 2014;9(7):907-914.
11. Andersen LL, Magnusson SP, Nielsen M, Haleem J, Poulsen K, Aagaard P. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Phys Ther*. 2006;86(5):683-697.
12. Van Wingerden JP, Vleeming A, Buyruk HM, Raissadat K. Stabilization of the sacroiliac joint in vivo: verification of muscular contribution to force closure of the pelvis. *Eur Spine J*. 2004;13(3):199-205.
13. Richardson C a, Snijders CJ, Hides J a, Damen L, Pas MS, Storm J. The relation between the transversus abdominis muscles, sacroiliac joint mechanics, and low back pain. *Spine* (Phila Pa 1976). 2002;27(4):399-405.
14. Hodges P, Kaigle Holm A, Holm S, et al. Intervertebral stiffness of the spine is increased by evoked contraction of transversus abdominis and the diaphragm: in vivo porcine studies. *Spine* (Phila Pa 1976). 2003;28(23):2594-2601.
15. Pickard CM, Sullivan PE, Allison GT, Singer KP. Is there a difference in hip joint position sense between young and older groups? *J Gerontol A Biol Sci Med Sci*. 2003;58(7):631-635.
16. Krause DA, Jacobs RS, Pilger KE, Sather BR, Sibunka SP, Hollman JH. Electromyographic analysis of the gluteus medius in five weight-bearing exercises. *J Strength Cond Res*. 2009;23(9):2689.
17. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: Part 1, mechanisms and risk factors. *Am J Sports Med*. 2006;34(2):299-311.
18. Allet L, Kim H, Ashton-Miller J, De Mott T, Richardson JK. Frontal plane hip and ankle sensorimotor function, not age, predicts unipedal stance time. *Muscle Nerve*. 2012;45(4):578-585.
19. Monteiro RL, Facchin JH, de Freitas DG, Callegari B, Joao SM. Hip Rotations Influence Electromyographic Activity of Gluteus Medius Muscle During Pelvic Drop Exercise. *J Sport Rehabil*. 2015.[Epub ahead of print]
20. Hamstra-Wright KL, Huxel Bliven K. Effective exercises for targeting the gluteus medius. *J Sport Rehabil*. 2012;21(3):296-300.
21. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. Electromyographic analysis of gluteus medius and gluteus maximus during rehabilitation exercises. *Int J Sports Phys Ther*. 2011;6(3):206-223.
22. Kendall KD, Patel C, Wiley JP, Pohl MB, Emery CA, Ferber R. Steps toward the validation of the Trendelenburg test: the effect of experimentally reduced hip abductor muscle function on frontal plane mechanics. *Clin J Sport Med*. 2013;23(1):45-51.
23. Hortobagyi T, Devita P. Muscle pre- and coactivity during downward stepping are associated with leg stiffness in aging. *J Electromyogr Kinesiol*. 2000;10(2):117-126.
24. Hortobágyi T, DeVita P. Altered movement strategy increases lower extremity stiffness during stepping down in the aged. *J Gerontol A Biol Sci Med Sci*. 1999;54(2):B63-B70.
25. Lark SD, Buckley JG, Bennett S, Jones D, Sargeant AJ. Joint torques and dynamic joint stiffness in elderly and young men during stepping down. *Clin Biomech*. 2003;18(9):848-855.
26. DeVita P, Hortobagyi T. Age increases the skeletal versus muscular component of lower extremity stiffness during stepping down. *J Gerontol A Biol Sci Med Sci*. 2000;55(12):B593-B600.
27. Ketcham CJ, Stelmach GE. Movement Control in the Older Adult. In: RW P, Van Hemel SB, eds. *Technology for Adaptive Aging*. Washington, DC: National Academies Press; 2004.

REFERENCES

1. Presswood L, Cronin J, Keogh JWL, Whatman C. Gluteus Medius: Applied Anatomy, Dysfunction, Assessment, and Progressive Strengthening. *Strength Cond J*. 2008;30:41-53.
2. Patten C, Kamen G, Rowland DM. Adaptations in maximal motor unit discharge rate to strength training in young and older adults. *Muscle Nerve*. 2001;24(4):542-550.
3. French HP, Dunleavy M, Cusack T. Activation levels of gluteus medius during therapeutic exercise as measured with electromyography: a structured review. *Phys Ther Rev*. 2010;15(2):92-105.
4. McBeth JM, Earl-Boehm JE, Cobb SC, Huddleston WE. Hip muscle activity during 3 side-lying hip-strengthening exercises in distance runners. *J Athl Train*. 47(1):15-23.
5. Lee J, Cynn H, Choi S, Yoon T, Jeong H. Effects of different hip rotations on gluteus medius and tensor fasciae latae muscle activity during isometric side-lying hip abduction. *J Sport Rehabil*. 2013;22:301-307.
6. Distefano LJ, Blackburn JT, Marshall SW, Padua DA. Gluteal muscle activation during common therapeutic exercises. *J Orthop Sports Phys Ther*. 2009;39:532-540.
7. Ekstrom RA, Donatelli RA, Carp KC. Electromyographic analysis of core trunk, hip, and thigh muscles during 9 rehabilitation exercises. *J Orthop Sports Phys Ther*. 2007;37(12):754-762.
8. Bolgla LA, Uhl TL. Electromyographic Analysis of Hip Rehabilitation Exercises in a Group of Healthy Subjects. *J Orthop Sports Phys Ther*. 2005;35(8):487-94
9. Macadam P, Cronin J, Contreras B. An examination of the gluteal muscle activity associated with dynamic hip abduction and hip external rotation exercise: a systematic review. *Int J Sports Phys Ther*. 2015;10(5):573-591.
10. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. Electromyographic analysis of gluteus medius and gluteus maximus during rehabilitation exercises. *Int J Sports Phys Ther*. 2011;6(3):206-223.
11. Philippon MJ, Decker MJ, Giphart JE, Torry MR, Wahoff MS, LaPrade RF. Rehabilitation exercise progression for the gluteus medius muscle with consideration for iliopsoas tendinitis: an in vivo electromyography study. *Am J Sports Med*. 2011;39:1777-1785.
12. Lee KS, Ko E, Lim CG. Effect of Pelvic Belt on Gluteal Muscles Activity during Therapeutic Exercise. *J Phys Ther Sci*. 2013;25(3):337-340.
13. Fredericson M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreicher N, Sahrmann SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med*. 2000;10:169-175.
14. Mascal CL, Landel R, Powers C. Management of patellofemoral pain targeting hip, pelvis, and trunk muscle function: 2 case reports. *J Orthop Sports Phys Ther*. 2003;33(11):647-660.
15. Lepley AS, Strouse AM, Ericksen HM, Pfile KR, Gribble P a, Pietrosimone BG. Relationship between gluteal muscle strength, corticospinal excitability, and jump-landing biomechanics in healthy women. *J Sport Rehabil*. 2013;22(4):239-247.
16. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther*. 2013;43(2):54-64.
17. Ford KR, Nguyen A-D, Dischiavi SL, Hegedus EJ, Zuk EF, Taylor JB. An evidence-based review of hip-focused neuromuscular exercise interventions to address dynamic lower extremity valgus. *Open access J Sport Med*. 2015;6:291-303.
18. Embaby E, Abdallah A. Trunk and Gluteus-Medius Muscles' Fatigability During Occupational Standing in Clinical Instructors with Low Back Pain. *Br J Sports Med*. 2013;47(10):e3-e3.
19. Pope MH, Goh KL, Magnusson ML. Spine ergonomics. *Annu Rev Biomed Eng*. 2002;4:49-68.
20. Allen DG, Lamb GD, Westerblad H. Skeletal Muscle Fatigue: Cellular Mechanisms. *Physiol Rev*. 2008;88(1):287-332.
21. Hansen L, Winkel J, Jørgensen K. Significance of mat and shoe softness during prolonged work in upright position: Based on measurements of low back muscle EMG, foot volume changes, discomfort and ground force reactions. *Appl Ergon*. 1998;29(3):217-224.
22. Lafond D, Champagne A, Descarreaux M, Dubois J-D, Prado JM, Duarte M. Postural control during prolonged standing in persons with chronic low back pain. *Gait Posture*. 2009;29(3):421-427.
23. Taylor JL, Butler JE, Gandevia SC. Changes in muscle afferents, motoneurons and motor drive during muscle fatigue. *Eur J Appl Physiol*. 2000;83(2-3):106-115.
24. Lewis CL, Sahrmann SA. Muscle activation and movement patterns during prone hip extension exercise in women. *J Athl Train*. 2009;44(3):238-248.
25. Kang SY, Jeon HS, Kwon O, Cynn H seock, Choi B. Activation of the gluteus maximus and hamstring muscles during prone hip extension with knee flexion in three hip abduction positions. *Man Ther*. 2013;18(4):303-307.
26. MacAskill MJ, Durant TJS, Wallace DA. Gluteal muscle activity during weightbearing and non-weightbearing exercise. *Int J Sports Phys Ther*. 2014;9(7):907-914.
27. Yoon J-Y, Lee M-R, An D-H. Effect of initial position on the muscle activity of the hip extensors and amount of pelvic tilt during prone hip extension. *J Phys Ther Sci*. 2015;27(4):1195-1197.
28. Fisher BE, Southam AC, Kuo Y-L, Lee Y-Y, Powers CM. Evidence of altered corticomotor excitability following targeted activation of gluteus maximus training in healthy individuals. *Neuroreport*. 2016;27(6):415-421.

REFERENCES

1. Presswood L, Cronin J, Keogh JW, Whatman C. Gluteus Medius: Applied Anatomy, Dysfunction, Assessment, and Progressive Strengthening. *Strength Cond J*. 2008;30:41-53.
2. Patten C, Kamen G, Rowland DM. Adaptations in maximal motor unit discharge rate to strength training in young and older adults. *Muscle Nerve*. 2001;24(4):542-550.
3. French HP, Dunleavy M, Cusack T. Activation levels of gluteus medius during therapeutic exercise as measured with electromyography: a structured review. *Phys Ther Rev*. 2010;15(2):92-105.
4. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. Electromyographic analysis of gluteus medius and gluteus maximus during rehabilitation exercises. *Int J Sports Phys Ther*. 2011;6(3):206-223.
5. Van Dieën JH, Ogita F, De Haan A. Reduced neural drive in bilateral exertions: a performance-limiting factor? *Med Sci Sports Exerc*. 2003;35(1):111-118.
6. Ferbert A, Priori A, Rothwell JC, Day BL, Colebatch JG, Marsden CD. Interhemispheric inhibition of the human motor cortex. *J Physiol*. 1992;453(D):525-546.
7. Oda S, Moritani T. Movement-related cortical potentials during handgrip contractions with special reference to force and electromyogram bilateral deficit. *Eur J Appl Physiol Occup Physiol*. 1995;72(1):1-5.
8. Willy RW, Scholz JP, Davis IS. Mirror gait retraining for the treatment of patellofemoral pain in female runners. *Clin Biomech* (Bristol, Avon). 2012;27(10):1045-1051.
9. Proske U, Morgan DL. Muscle damage from eccentric exercise: Mechanism, mechanical signs, adaptation and clinical applications. *J Physiol*. 2001;537(2):333-345.
10. McHugh MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand J Med Sci Sports*. 2003;13(2):88-97.
11. Lamb T, Yang JF. Could different directions of infant stepping be controlled by the same locomotor central pattern generator? *J Neurophysiol*. 2000;83:2814-2824.
12. Marder E, Calabrese RL. Principles of rhythmic motor pattern generation. *Physiol Rev*. 1996;76(3):687-717.
13. Stein PS., Mortin LI. The forms of a task and their blends. In: *Neurobiology of Vertebrate Locomotion*. London: Macmillan; 1986:201-216.
14. Simenz CJ, Garceau LR, Lutsch BN, Suchomel TJ, Ebben WP. Electromyographical analysis of lower extremity muscle activation during variations of the loaded step-up exercise. *J Strength Cond Res*. 2012;26(12):3398-3405.
15. Krause DA, Jacobs RS, Pilger KE, Sather BR, Sibunka SP, Hollman JH. Electromyographic analysis of the gluteus medius in five weight-bearing exercises. *J Strength Cond Res*. 2009;23(9):2689.
16. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: Part 1, mechanisms and risk factors. *Am J Sports Med*. 2006;34(2):299-311.
17. Karim ME. A systematic approximation to the neural motor control of forward, backward and lateral toe-tapping in children and adults. *Digital Repository at the University of Maryland* 2014.
18. Bolgla LA, Uhl TL. Electromyographic Analysis of Hip Rehabilitation Exercises in a Group of Healthy Subjects. *J Orthop Sports Phys Ther*. 2005; 35(8):487-94.
19. Neumann D a. Hip abductor muscle activity as subjects with hip prostheses walk with different methods of using a cane. *Phys Ther*. 1998;78(5):490-501.
20. Berry JW, Lee TS, Foley HD, Lewis CL. Resisted Side Stepping: The Effect of Posture on Hip Abductor Muscle Activation. *J Orthop Sports Phys Ther*. 2015;45(9):675-682.
21. Schmitz RJ, Riemann BL, Thompson T. Gluteus medius activity during isometric closed-chain hip rotation. *J Sport Rehabil*. 2002;11(3):179-188.
22. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther*. 2013;43:54-64.
23. Sinsurin K, Pluemjai S, Srisangboriboon S, Suanshan S, Vachalathiti R. Gluteus medius muscle activities during standing hip abduction exercises in the transverse plane at different angles. *J Med Assoc Thail*. 2015;98:S42-S47.
24. Ferber R, Kendall KD, Farr L. Changes in knee biomechanics after a hip-abductor strengthening protocol for runners with patellofemoral pain syndrome. *J Athl Train*. 2011;46(2):142-149.
25. Akima H, Takahashi H, Kuno SY, et al. Early phase adaptations of muscle use and strength to isokinetic training. *Med Sci Sports Exerc*. 1999;31(4):588-594.
26. Hamstra-Wright KL, Huxel Bliven K. Effective exercises for targeting the gluteus medius. *J Sport Rehabil*. 2012;21(3):296-300.
27. Barton CJ, Kennedy A, Twycross-Lewis R, Woledge R, Malliaras P, Morrissey D. Gluteal muscle activation during the isometric phase of squatting exercises with and without a Swiss ball. *Phys Ther Sport*. 2014;15(1):39-46.
28. Blanpied P. Changes in muscles activation during wall slides and squat-machine exercise. *J Sport Rehabil*. 1999;(10):123.
29. Refshauge KM, Chan R, Taylor JL, McCloskey DI. Detection of movements imposed on human hip, knee, ankle and toe joints. *J Physiol*. 1995;488(1):231-241.

REFERENCES

1. Grimaldi A, Richardson C, Durbridge G, Donnelly W, Darnell R, Hides J. The association between degenerative hip joint pathology and size of the gluteus maximus and tensor fascia lata muscles. *Man Ther.* 2009;14(6):611-617.
2. Kankaanpää M, Taimela S, Laaksonen D, Hänninen O, Airaksinen O. Back and hip extensor fatigability in chronic low back pain patients and controls. *Arch Phys Med Rehabil.* 1998;79(4):412-417.
3. Leinonen V, Kankaanpää M, Airaksinen O, Hänninen O. Back and hip extensor activities during trunk flexion/extension: Effects of low back pain and rehabilitation. *Arch Phys Med Rehabil.* 2000;81(1):32-37.
4. Robinson RL, Nee RJ. Analysis of hip strength in females seeking physical therapy treatment for unilateral patellofemoral pain syndrome. *J Orthop Sport Phys Ther.* 2007;37(5):232-238.
5. Bolgla L a, Malone TR, Umberger BR, Uhl TL. Hip strength and hip and knee kinematics during stair descent in females with and without patellofemoral pain syndrome. *J Orthop Sports Phys Ther.* 2008;38(1):12-18.
6. Ferber R, Noehren B, Hamill J, Davis IS. Competitive female runners with a history of iliotibial band syndrome demonstrate atypical hip and knee kinematics. *J Orthop Sports Phys Ther.* 2010;40(2):52-58.
7. Noehren B, Davis I, Hamill J. ASB Clinical Biomechanics Award Winner 2006. Prospective study of the biomechanical factors associated with iliotibial band syndrome. *Clin Biomech.* 2007;22(9):951-956.
8. Fredericson M, Cookingham CL, Chaudhari a M, Dowdell BC, Oestreicher N, Sahrmann S a. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med.* 2000;10(3):169-175.
9. Fredericson M, Weir A. Practical management of iliotibial band friction syndrome in runners. *Clin J Sport Med.* 2006;16(3):261-268.
10. Webster KA, Gribble PA. A comparison of electromyography of gluteus medius and maximus in subjects with and without chronic ankle instability during two functional exercises. *Phys Ther Sport.* 2013;14(1):17-22.
11. Goo Y-M, Kim T-H, Lim J-Y. The effects of gluteus maximus and abductor hallucis strengthening exercises for four weeks on navicular drop and lower extremity muscle activity during gait with flatfoot. *J Phys Ther Sci.* 2016;28(3):911-915.
12. Mills M, Frank B, Goto S, et al. Effect of Restricted Hip Flexor Muscle Length on Hip Extensor Muscle Activity and Lower Extremity Biomechanics in College-Aged Female Soccer Players. *Int J Sports Phys Ther.* 2015;10(7):946-954.
13. Comyns T, Kenny I, Scales G. Effects of a Low-Load Gluteal Warm-Up on Explosive Jump Performance. *J Hum Kinet.* 2015;46(46):177-187.
14. Crow JF, Buttifant D, Kearny SG, Hrysomallis C. Low Load Exercises Targeting the Gluteal Muscle Group Acutely Enhance Explosive Power Output in Elite Athletes. *J Strength Cond Res.* 2012;26:438-442.
15. Reiman MP, Bolgla L a, Loudon JK. A literature review of studies evaluating gluteus maximus and gluteus medius activation during rehabilitation exercises. *Physiother Theory Pract.* 2012;28(4):257-268.
16. Neumann D a. Kinesiology of the hip: a focus on muscular actions. *J Orthop Sports Phys Ther.* 2010;40(2):82-94.
17. Khuu A, Foch E, Lewis CL. Not all single leg squats are equal: A biomechanical comparison of three variations. *Int J Sports Phys Ther.* 2016;11(2):201-211.
18. Escamilla RF, Zheng N, Macleod TD, et al. Patellofemoral joint force and stress between a short- and long-step forward lunge. *J Orthop Sports Phys Ther.* 2008;38(11):681-690.
19. Salsich GB. The Effects of Movement Pattern Modification on Lower Extremity Kinematics and Pain in Females With Patellofemoral Pain. *J Orthop Sports Phys Ther.* 2012;42(12):1017-1024.
20. Hwangbo P-N. The effects of squatting with visual feedback on the muscle activation of the vastus medialis oblique and the vastus lateralis in young adults with an increased quadriceps angle. *J Phys Ther Sci.* 2015;27(5):1507-1510.
21. Clohisy JC, Knaus ER, Hunt DM, Leshner JM, Harris-Hayes M, Prather H. Clinical presentation of patients with symptomatic anterior hip impingement. In: *Clinical Orthopaedics and Related Research.* Vol 467.; 2009:638-644.
22. Nepple JJ, Prather H, Trousdale RT, et al. Clinical diagnosis of femoroacetabular impingement. *J Am Acad Orthop Surg.* 2013;21 Suppl 1:S16-S19.
23. Kulas AS, Hortobogyi T, Devita P. Trunk position modulates anterior cruciate ligament forces and strains during a single-leg squat. *Clin Biomech.* 2012;27(1):16-21.
24. Lynn SK, Noffal GJ. Lower Extremity Biomechanics During a Regular and Counterbalanced Squat. *J Strength Cond Res.* 2012;26(9):2417-2425.
25. Kulas AS, Hortobogyi T, Devita P. Trunk position modulates anterior cruciate ligament forces and strains during a single-leg squat. *Clin Biomech.* 2012;27(1):16-21.
26. Boudreau SN, Dwyer MK, Mattacola CG, Lattermann C, Uhl TL, Mckeon JM. Hip-Muscle Activation During the Lunge, Single-Leg Squat, and Step-Up-and-Over Exercises. *J Sport Rehabil.* 2009;18:91-103.
27. Ayotte NW, Stetts DM, Keenan G, Greenway EH. Electromyographical analysis of selected lower extremity muscles during 5 unilateral weight-bearing exercises. *J Orthop Sports Phys Ther.* 2007;37:48-55.

28. Distefano LJ, Blackburn JT, Marshall SW, Padua DA. Gluteal muscle activation during common therapeutic exercises. *J Orthop Sports Phys Ther.* 2009;39:532-540.
29. Ekstrom RA, Donatelli RA, Carp KC. Electromyographic analysis of core trunk, hip, and thigh muscles during 9 rehabilitation exercises. *J Orthop Sports Phys Ther.* 2007;37(12):754-762.
30. Barton CJ, Kennedy A, Twycross-Lewis R, Woledge R, Malliaras P, Morrissey D. Gluteal muscle activation during the isometric phase of squatting exercises with and without a Swiss ball. *Phys Ther Sport.* 2014;15(1):39-46.
31. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. Electromyographic analysis of gluteus medius and gluteus maximus during rehabilitation exercises. *Int J Sports Phys Ther.* 2011;6(3):206-223.
32. Mccurdy KW, Langford GA, Doscher MW, Wiley LP, Mallard KG. The effects of short-term unilateral and bilateral lower-body resistance training on measures of strength and power. *J Strength Cond Res Natl Strength Cond Assoc.* 2005;19(1):9-15.
33. Mellor R, Hodges PW. Motor unit synchronization of the vasti muscles in closed and open chain tasks. *Arch Phys Med Rehabil.* 2005;86(4):716-721.
34. Mohr M, Nann M, Von Tscharnar V, Eskofier B, Nigg BM. Task-dependent intermuscular motor unit synchronization between medial and lateral Vastii muscles during dynamic and isometric squats. *PLoS One.* 2015;10(11).
35. Xu Y, Bach E, Orhede E. Work environment and low back pain: the influence of occupational activities. *Occup Environ Med.* 1997;54:741-745.
36. Hoogendoorn WE, Bongers PM, de Vet HC, et al. Flexion and rotation of the trunk and lifting at work are risk factors for low back pain: results of a prospective cohort study. *Spine (Phila Pa 1976).* 2000;25(23):3087-3092.
37. Macadam P, Cronin J, Contreras B. An examination of the gluteal muscle activity associated with dynamic hip abduction and hip external rotation exercise: a systematic review. *Int J Sports Phys Ther.* 2015;10(5):573-591.
38. Beckman SM, Buchanan TS. Ankle inversion injury and hypermobility: Effect on hip and ankle muscle electromyography onset latency. *Arch Phys Med Rehabil.* 1995;76(12):1138-1143.
39. Bullock-Saxton JE. Local sensation changes and altered hip muscle function following severe ankle sprain. *Phys Ther.* 1994;74(1):17-28; discussion 28-31.
40. Bullock-Saxton JE, Janda V, Bullock MI. The Influence of Ankle Sprain Injury on Muscle Activation during Hip Extension. *Int J Sports Med.* 1994;15(6):330-4
41. Caulfield BM, Garrett M. Functional instability of the ankle: Differences in patterns of ankle and knee movement prior to and post landing in a single leg jump. *Int J Sports Med.* 2002;23(31):64-68.
42. Gribble PA, Hertel J, Denegar CR, Buckley WE. The Effects of Fatigue and Chronic Ankle Instability on Dynamic Postural Control. *J Athl Train.* 2004;39(4):321-329.
43. Van Deun S, Staes FF, Stappaerts KH, Janssens L, Levin O, Peers KKH. Relationship of chronic ankle instability to muscle activation patterns during the transition from double-leg to single-leg stance. *Am J Sports Med.* 2007;35(2):274-281.
44. Leetun DT, Ireland ML, Willson JD, Ballantyne BT, Davis IM. Core stability measures as risk factors for lower extremity injury in athletes. *Med Sci Sports Exerc.* 2004;36(6):926-934.
45. Gottschalk F, Kourosh S, Leveau B. The functional anatomy of tensor fasciae latae and gluteus medius and minimus. *J Anat.* 1989;166:179-189.
46. Hase K, Stein RB. Turning strategies during human walking. *J Neurophysiol.* 1999;81(6):2914-2922.
47. Neumann DA. Kinesiology of the hip: A focus on muscular actions. *Joural Orthop Sport Phys Ther.* 2010;40(2):82-94.
48. Aminaka N, Pietrosimone BG, Armstrong CW, Meszaros A, Gribble PA. Patellofemoral pain syndrome alters neuromuscular control and kinetics during stair ambulation. *J Electromyogr Kinesiol.* 2011;21(4):645-651.
49. Brindle TJ, Mattacola C, McCrory J. Electromyographic changes in the gluteus medius during stair ascent and descent in subjects with anterior knee pain. *Knee Surg Sports Traumatol Arthrosc.* 2003;11(4):244-251.
50. Willson JD, Kernozek TW, Arndt RL, Reznichek DA, Scott Straker J. Gluteal muscle activation during running in females with and without patellofemoral pain syndrome. *Clin Biomech.* 2011;26(7):735-740.
51. Cowan SM, Crossley KM, Bennell KL. Altered hip and trunk muscle function in individuals with patellofemoral pain. *Br J Sports Med.* 2009;43(8):584-588.
52. Kim D, Unger J, Lanovaz JL, Oates AR. The Relationship of Anticipatory Gluteus Medius Activity to Pelvic and Knee Stability in the Transition to Single-Leg Stance. *PM R.* 2016;8(2):138-144.
53. Barati A, Safarcherati A, Aghayari A, Azizi F, Abbasi H. Evaluation of relationship between trunk muscle endurance and static balance in male students. *Asian J Sports Med.* 2013;4(4):289-294.
54. Balter SGT, Stokroos RJ, Akkermans E, Kingma H. Habituation to galvanic vestibular stimulation for analysis of postural control abilities in gymnasts. *Neurosci Lett.* 2004;366(1):71-75.
55. Ashton-Miller JA, Wojtys EM, Huston LJ, Fry-Welch D. Can proprioception really be improved by exercises? *Knee Surgery, Sport Traumatol Arthrosc.* 2001;9(3):128-136.
56. Youdas JW, Boor MMP, Darfler AL, Koenig MK, Mills KM, Hollman JH. Surface electromyographic analysis of core trunk and hip muscles during selected rehabilitation exercises in the side-bridge to neutral spine position. *Sports Health.* 2014;6(5):416-421.

57. Marshall PWM, Patel H, Callaghan JP. Gluteus medius strength, endurance, and co-activation in the development of low back pain during prolonged standing. *Hum Mov Sci.* 2011;30(1):63-73.
58. McGill SM. Low back exercises: evidence for improving exercise regimens. *Phys Ther.* 1998;78(7):754-765.
59. Cormier G, Berthelot JM, Maugars Y. Gluteus tendon rupture is under recognized by French orthopedic surgeons: results of a mail survey. *Jt Bone Spine.* 2006;73(4):411-413.
60. Kingzett-Taylor A, Tirman PFJ, Feller J, et al. Tendinosis and tears of gluteus medius and minimus muscles as a cause of hip pain: MR imaging findings. *Am J Roentgenol.* 1999;173(4):1123-1126.
61. Robertson WJ, Gardner MJ, Barker JU, Boraiah S, Lorich DG, Kelly BT. Anatomy and Dimensions of the Gluteus Medius Tendon Insertion. *Arthrosc - J Arthrosc Relat Surg.* 2008;24(2):130-136.
62. Chandrasekaran S, Vemula SP, Gui C, Suarez-Ahedo C, Lodhia P, Domb BG. Clinical Features That Predict the Need for Operative Intervention in Gluteus Medius Tears. *Orthop J Sport Med.* 2015;3(2):2325967115571079.
63. Ayotte NW, Stetts DM, Keenan G, Greenway EH. Electromyographical analysis of selected lower extremity muscles during 5 unilateral weight-bearing exercises. *J Orthop Sports Phys Ther.* 2007;37(2):48-55.
64. Cram JR, Criswell E. *Cram's Introduction to Surface Electromyography.* Jones & Bartlett Learning; 2011.
65. Yang JF, Winter DA. Electromyographic amplitude normalization methods: improving their sensitivity as diagnostic tools in gait analysis. *Arch Phys Med Rehabil.* 1984;65(9):517-521.
66. Dowling JJ, Konert E, Ljucovic P, Andrews DM. Are humans able to voluntarily elicit maximum muscle force? *Neurosci Lett.* 1994;179(1-2):25-28.