

Patient name:

INTERMEDIATE EXERCISES FOR THE GLUTEALS (LOW TENSOR FASCIA LATAE)

These exercises are aimed at strengthening your major buttock muscles (gluteals) while minimising the activation of the tensor fascia latae.

CLAMS WITH ELASTIC BAND

Step 1

Lie on your side with your knees bent and your hand supporting your head. Wrap an elastic band around your lower thighs.



Step 2

Slowly turn your top leg outwards keeping your feet together. You should not twist your body. You should feel your buttock muscles working.



Step 3

Return your leg to the step 1 position.

Repeat steps 2 and 3 ten times each side.

SINGLE LEG SUPINE BRIDGE

Step 1

Lie on your back with your knees bent, feet flat, hip width apart. Elevate one leg, keeping the knee bent to 90 degrees.



Step 2

Place pressure through the heel on the floor and gently squeeze your buttocks.

Step 3

Slowly raise your pelvis and torso until your body is straight. You should maintain the natural curve in your lower back.



Step 4

Slowly return to the step 1 position.

Repeat steps 2 to 4 ten times. You can straighten the elevated leg to increase the difficulty of this exercise.

DONKEY KICK (BENT LEG)

Step 1

On all fours, with your upper body on your elbows and forearms.



Step 3

Elevate the leg until your hip is straight.



Step 2

Elevate one leg slightly, keeping the knee bent to 90 degrees.

Step 4

Slowly lower your leg to the step 2 position.

Repeat steps 3 to 4 ten times.

The gluteal muscles provide essential support for your pelvic and hip joints.

DONKEY KICK (STRAIGHT LEG)

Step 1

On all fours with your upper body on your elbows and forearms.



Step 3

Elevate the leg until your hip is straight.



Step 2

Extend one leg behind you and elevate it slightly.

Step 4

Slowly lower your leg to the step 2 position.

Repeat steps 3 to 4 ten times.

Your gluteal muscles also play important role in the alignment of your hip, knee and ankle joints.

MONSTER WALKS

Step 1

Start in a squat position with an elastic band wrapped around your lower thighs. Feet should be hip width apart. Your knees should be in line with your toes.

The deep gluteal muscles provide essential feedback to the posture and movement centres of the brain.



Step 2

Take a side step maintaining a squat position.



Step 3

Bring your other leg back to the step 1 position.

Repeat steps 2 to 3 five times in one direction.

Repeat towards the other direction.



Performing specific gluteal exercises can reduce the risk of hip and knee joint injuries.

This material is copyright © Evidence for Exercise®. Not for re-sale. Reproduction and distribution is only permitted in the year 2022 under the Evidence for Exercise® registered provider terms as described at evidenceforexercise.org. You should seek professional advice from a relevantly qualified person to discuss any personal circumstances which may affect your ability to undertake particular exercise. Evidence for Exercise® shall not be liable for any injury caused, whether due to negligence or otherwise arising from use of or reliance on this information.