



INTERMEDIATE EXERCISES FOR THE GLUTEALS (LOW TENSOR FASCIA LATAE) PRACTITIONER NOTES

These Evidence for Exercise™ practitioner notes provide you with specific guidelines to prescribe intermediate exercises to strengthen the gluteal musculature while minimising activation of the tensor fascia latae.

What to prescribe them for

- Overactive tensor fascia latae¹

Excessive activation of the tensor fascia latae during exercises may be detrimental in patients with excessive hip internal rotation.¹ The tensor fascia latae is both a hip abductor and internal rotator.⁹ It can also exert a lateral force on the patella through its connections with the iliotibial band.¹⁰⁻¹² A small study found that “patients with abductor tendon tears showed hypertrophy of the tensor fascia latae muscle when compared to the contralateral healthy side and to patients without a tear.”¹³

- Hip pain

Tears of the gluteus medius and minimus are associated with hip pain^{14,15} although evidence relating to hip abduction exercises and hip pain reduction are lacking.

- Degenerative hip joint pathology²

Atrophy of the gluteus maximus relative to the tensor fascia latae has been observed in patients with advanced degenerative hip joint pathology.^{1,16} These patients also demonstrate increased gluteus medius activation during stepping activities which is considered a compensation for weakness.¹⁷ Both targeted hip programs and general exercise can improve gluteus minimus volume in patients with hip osteoarthritis.¹⁸ Interestingly, in the early

stages of hip joint pathology hypertrophy of the hip abductor muscles may be present and this should be considered when prescribing gluteal exercises.¹⁹

- Lower back pain³

Gluteus medius weakness and gluteal muscle tenderness are common symptoms in people with chronic non-specific lower back pain.²⁰⁻²² There is some association between gluteus medius and maximus weakness and lower back pain.²³⁻²⁷ While limited information regarding the effectiveness of hip strengthening exercises for lower back pain exists there is some indication they may be beneficial.^{28,3}

- Sacroiliac joint pain

Shear in the sacroiliac joint according to one model is prevented by two factors:

1. Form closure – joint anatomical features that increase the friction coefficient²⁹⁻³¹

2. Force closure - Tension of muscles and ligaments crossing the joint that lead to higher friction and therefore stiffness^{32,30}

Muscles that could increase force closure include gluteus maximus and biceps femoris³³⁻³⁵ (due to their attachments to the sacrotuberous ligament), latissimus dorsi³⁶ (due to its partial coupling with gluteus maximus by the posterior layer of the thoracolumbar fascia, creating a compressive force acting perpendicular to the sacroiliac joint) and the erector spinae³⁷ (which are closely linked to the sacrum and posterior superficial sacroiliac ligaments).

The erector spinae, biceps femoris and gluteus maximus muscles have been shown to have a significant effect on sacroiliac joint stiffness.³⁰ Both the sacroiliac joint³⁸⁻⁴¹ and the long dorsal sacroiliac ligament⁴²⁻⁴⁶ can be significant pain generators in those with pelvic girdle pain. Patients with sacroiliac joint pain have been shown to display a delayed onset of gluteus maximus on the stance leg during standing hip flexion compared with healthy subjects.⁴⁷ Due to a lack of investigation it is unclear if exercises for the gluteals improve sacroiliac joint pain.

- Groin pain

Athletes with groin pain are more likely to display enlarged tensor fascia latae on sonography.⁴

- Patellofemoral pain syndrome^{5,6}

Patellofemoral pain syndrome has been associated with weak hip abduction and external rotation,⁴⁸⁻⁵⁰ excessive internal rotation of the hip and lateral patella displacement⁵¹⁻⁵³ while general knee pathology has been associated with hip dysfunction that has ensued from gluteal weakness.⁵⁴ Alignment of the thigh and leg in the frontal plane can be heavily influenced by hip-abductor muscle weakness particularly during daily activities such as climbing/descending stairs, sitting or squatting.⁵⁵ Selkowitz et al. found that those with patellofemoral pain had higher EMG activity of tensor fascia latae and lower EMG activity of the superior portion of gluteus maximus and gluteus medius during different hip exercises.⁵⁶

Hip abduction strength exercises have demonstrated favourable outcomes for patellofemoral pain syndrome.⁵⁷⁻⁶¹

- Anterior cruciate ligament injury prevention

Increased attention has been given to neuromuscular exercise focused at the hip for anterior cruciate ligament ruptures.^{62,51} Poor hip strength and neuromuscular control has been associated with dynamic lower extremity valgus.^{63,64} In female athletes, future anterior cruciate ligament injury risk is significantly correlated with high knee abduction moments.^{63,65} This is reflected in the higher incidence of both anterior cruciate ligament ruptures in females who tend towards greater valgus alignment during landing and pivoting compared with men.⁶⁶⁻⁷⁸

Hip external rotation strength predicts hop performance after anterior cruciate ligament reconstruction.⁷⁹

- Iliotibial band syndrome⁷

Iliotibial band syndrome has been associated with greater hip adduction and knee internal rotation⁸⁰⁻⁸² as well as hip abductor weakness.⁸³ Hip abduction strength exercises have been recommended for these patients.^{84,85}

- Chronic ankle instability

Those with chronic ankle instability show a decreased onset latency of gluteus medius.⁸⁶ It is thought that weak hip abduction may limit the amount of time available to initiate the hip strategy required to counteract a sudden lateral external perturbation.⁸⁷ Hip abductor muscle strength deficit is a risk factor for inversion sprain in soccer players.⁸⁸

- Improving athletic performance
- Lower limb injury prevention⁸

Both the gluteus medius and maximus assist in load transference through the hip joint⁸⁹ providing local structural stability and contributing to alignment of the knee and hip joints.⁹⁰ The gluteal muscles can enhance athletic performance^{48,91-93} and contribute to the prevention or rehabilitation of lower extremity injuries.⁹⁴⁻⁹⁸

How to prescribe them

These exercises can be prescribed in a 20-minute, one-on-one consultation. Following the consultation they can be incorporated into a group session or performed as home exercises. Follow-up consultations are recommended to assess technique and outcomes. For optimal results they should be performed for a minimum of four to six weeks and ideally incorporated into a long-term daily routine.

What the patient can expect

This series of exercises progresses from beginner exercises for the hip. Patients should continue to experience improvement in their condition be it hip, pelvic, lower back or knee pain. They should gain further confidence in their daily activities and notice increases in strength.

It should be noted that this exercise prescription is based on two studies that compared multiple exercises to ascertain those that provided the most favourable activation of the gluteal muscles while

minimising the activity of tensor fascia latae.^{9,99} In the 2023 study only clams with elastic band was considered ideal (greater activation of both gluteals compared to tensor fascia latae), contradicting the earlier study. That study also identified single leg supine bridge, donkey kick (bent leg), donkey kick

(straight leg) and monster walk as producing greater than 50% higher normalised electromyographic amplitudes for gluteus medius and maximus compared to the tensor fascia lata, whereas sidelying hip abduction, supine bridge exercise, hip hike, lunge, squat and step-up step-down did not.

CLAMS WITH ELASTIC BAND

Teaching points

Many patients contract their quadriceps resulting in hip flexion.^{*100} Explain that the leg is to turn outwards and that it is the muscles of the buttock that do this. Have the patient palpate their gluteal muscles to identify the contraction.

If you wish to emphasise stability of the femoral head in the acetabulum you can ask the patient to 'suck your leg bone into its socket' before commencing the movement.

A small study indicated that the degree of hip flexion (30°, 45° or 60°) during clams did not significantly alter the predominance of gluteus medius activation over tensor fascia latae activation. The authors speculated that "Proper technique (e.g. no spine 'twisting' or rotation at the pelvis to initiate the movement) is likely a more substantial consideration when prescribing and performing these exercises."¹⁰¹

*In their study in runners McBeth et al.¹⁰⁰ found the anterior hip flexors were more active than the gluteus medius during clams. They questioned the relevance of the exercise for strengthening hip abductors and external rotators while others support its use.¹ It could be suggested that the patient perform clams in a more extended hip position if the quadriceps seems to dominate the movement.

Key teaching phrases

"You should not feel the muscles in the front of your leg working – if so you may be lifting your leg forwards."

Contraindications

- Neck pain in this position
- Excessive back or hip pain during this exercise
- Excessive trochanteric tenderness of the hip on the floor

Evidence

This exercise preferentially activates the gluteus medius and maximus while limiting recruitment of the tensor fascia latae.^{1,9} It is believed to activate the deep hip external rotators (quadratus femoris, obturator internus and externus, and the gemelli).^{102,103} These muscles appear to play a key role in hip joint stability by stabilising the femoral head in the acetabulum.¹⁰⁴ It is speculated they modulate hip joint stiffness and provide finely tuned adjustments to hip joint position.

Please note that when the hip is in the flexed position the gluteus medius can no longer act as a primary hip abductor¹⁰⁵ and therefore this exercise does not target this muscle as much as some other exercises (such as outer thigh exercise). However, if you wish to emphasise gluteus medius activity over tensor fascia latae, clams has a more favourable ratio.¹⁰¹

Neuro tip

Ballet dancers display greater hip external rotation strength and hip external rotation range of motion compared with non-dancers.¹⁰⁶⁻¹⁰⁸ This has been attributed to a number of factors, including "adaptive changes to the neural subsystems in response to the functional demands of ballet."¹⁰⁶ This could include increases in electromyographic amplitude and rate of electromyographic activation (a phenomenon found in other highly trained athletes) indicating that "improvements to the pattern of neural drive occurs with sport-specific resistance training."¹⁰⁹ Understandably this increase in external rotation may be related to injury. It appears that compensated turnout (where the turnout of the lower limbs is greater than the available range at the hip) increases this risk.¹¹⁰ Focusing on external rotation strength from the hip is recommended for dancers.

What to say to patients

"This exercise retrains small rotational movements which is important for stabilising your hip joint as well as providing feedback to your brain about the movements of your lower limb."

SINGLE LEG SUPINE BRIDGE

Teaching points

Anecdotally, asking the patient to place pressure through the heel (without lifting the toes) and keeping the foot close to the buttocks increases activation of the gluteus maximus. Verbal cues to maximally contract the gluteal muscles may also be beneficial – although some patients find this difficult to do without losing their natural lumbar lordosis. A sign of excessive hamstring activation is cramping or observation of a posterior pelvic tilt. You or the patient can palpate the hamstring to ensure it stays relatively relaxed.

Poor technique: posterior pelvic tilt



Alternative technique: you can increase the challenge of this exercise by straightening the free leg.



A variation on this exercise is to slightly abduct the stabilising leg, externally rotate the knee and externally rotate the foot. This may increase activity of the gluteals and emphasises lateral hamstring activation over medial hamstring activity.¹¹¹ Alternatively, the elevated leg can be straightened and abducted to increase rectus abdominis activity.¹¹²

Key teaching phrases

- “Maintain a natural curve in your lower back.”
- “Do not let your knee drift outwards.”
- “Do not use your hands to push up.”
- “You should feel the muscles in your buttocks working.”

Contraindications

- Excessive pain in the knees, hips or ankles
- Back pain during this exercise
- Acute disc episodes

Evidence

Gluteus maximus activation has been observed to range between 35% and 54% of maximum voluntary contraction during this exercise.^{113,1,9,114} Interestingly, standing gluteal squeeze has been shown to result in greater activation of the gluteus maximus. The authors speculated that substitution with the hamstrings during single leg supine bridge is a factor given that multiple subjects reported hamstring cramping during the single leg bridge and that their study was limited

by the fact that gluteal squeeze was the only exercise where verbal cues were given to maximally contract the gluteal muscles.¹¹³ In Selkowitz et al.’s 2023 study gluteus maximus was only activated at 17.7% MVC compared to 34% for tensor fascia latae.⁹

There is conflicting evidence regarding the level of activation of gluteus medius during this exercise; however, it is generally slightly less than levels observed in outer thigh exercise (side-lying hip abduction).^{1,113}

Neuro tip

The increased risk of anterior cruciate ligament injuries in women is not only attributed to dynamic genu valgus but also the tendency towards increased knee joint laxity, which appears to contribute to diminished joint proprioception.¹¹⁵ A common compensation for this is increased hamstring activity.¹¹⁶ Patients with a history of anterior cruciate ligament problems should be monitored for excessive hamstring activity during this exercise.

What to say to patients

- “This exercise requires coordinated movements to activate your buttock muscles while relaxing the back of your thigh.”

DONKEY KICK (BENT LEG)

Teaching points

This exercise can be surprisingly difficult for a patient to perform without inappropriately relying on the erector spinae to contribute to the movement.

Signs of this occurring include:

- Excessive lumbar lordosis
- Observation of excessive erector spinae activity
- Patient reports ‘lower back muscles working’
- Patient cannot ‘feel the buttock muscles working’

Poor technique: excessive lumbar lordosis



Poor technique: excessive erector spinae activity



Correct technique: relaxed erector spinae



To address poor technique you may wish to use the following techniques:

- Place your hands on the erector spinae and ask the patient to 'relax the muscles under your hands'
- Ask the patient to contract the gluteal muscles while lifting the leg¹¹⁷
- Prescribe the exercises with a smaller range of motion – stopping the leg movement at the point in which the patient starts to use their erector spinae
- Allow the patient to use a reduced lumbar lordosis initially (this is not ideal technique; however, some patients find it beneficial to identify their gluteus maximus. Once this has been achieved they can return to using a natural lumbar lordosis)

Modification: small leg movement



Modification: reduced lumbar lordosis



Prescribing exercises on all fours (refer to beginner and intermediate exercises for the lower back) prior to this exercise may also assist in patient awareness of their spine.

You can add an ankle weight to increase the challenge of this exercise. It can also be performed in the all-fours position for those who find it difficult to rest on their elbows.

Key teaching phrases

- “Maintain a natural curve in your lower back.”
- “Imagine your buttock muscles are elevating your leg not your lower back muscles.”
- “Keep your head in line with your body.”

Contraindications

- Excessive pain in the elbows, knees, hips or ankles
- Inability to maintain natural lumbar curve during this exercise
- Back pain during this exercise
- Acute disc episodes

Evidence

This exercise has demonstrated varying activation levels in different studies but can result in close to 60% of maximum voluntary contraction for gluteus maximus making it an appropriate strengthening exercise.^{113,118,119} In Selkowitz et al's 2023 study superior gluteus maximus was activated at 20 MVC compared with 28.5 for tensor fascia latae.⁹

Any decrease in force contribution from the gluteal muscles during hip extension can result in increased anterior hip force with potential for anterior hip pain, subtle hip instability or a tear of the acetabular labrum.^{120,121}

Neuro tip

At various times the focus for managing lower back and pelvic pain has moved from a strengthening approach to focusing on motor patterns. Janda developed a test to assess the muscular activation pattern during hip extension in the prone position.¹²² He theorised that the muscle activation pattern in this test could mimic those used during gait. Janda taught that the pattern of activation should be hamstring first, followed by gluteus maximus, contralateral lumbosacral erector spinae, ipsilateral lumbosacral erector spinae, contralateral thoracolumbar erector spinae, ipsilateral thoracolumbar erector spinae, and finally by the thoracic erector spinae.¹²³ However, when studied, a variety of activation patterns have been observed in healthy subjects.^{124,125} Even during gait the muscle recruitment patterns appear to be top-down in order.¹²⁶ Lehman et al. concluded: “The utility of the prone leg extension as a clinical and functional test is questionable due to the normal variability seen during the test and our current inability to determine what is normal and what is dysfunctional.”¹²⁵

What to say to patients

“This exercise is effective in strengthening the large buttock muscle, gluteus maximus. This muscle is important for controlling your hips and spine during typical daily activities such as bending and running.”

DONKEY KICK (STRAIGHT LEG)

Teaching points

See donkey kick (bent leg)

Patients who are likely to use their erector spinae may find this version of the exercise more difficult. You may wish to ensure the patient has mastered donkey kick (bent leg) prior to prescribing this exercise.

Key teaching phrases

- “Maintain a natural curve in your lower back.”
- “Imagine your buttock muscles are elevating your leg not your lower back muscles.”
- “Keep your head in line with your body.”

Contraindications

- Excessive pain in the elbows, knees, hips or ankles
- Inability to maintain natural lumbar curve during this exercise
- Back pain during this exercise
- Acute disc episodes

Evidence

See donkey kick (bent leg)

MONSTER WALKS

Teaching points

This exercise is best performed in the squat position as electromyographic activity is greater in the gluteus medius and maximus and lower in the tensor fascia latae compared with upright standing.¹²⁷

If you wish to increase the activity of the gluteus muscles you can place the band more distally – around the ankle (increases about 20%) or foot (increases about 40%).¹²⁸ It is thought that the lower position creates an external rotation moment that needs to be overcome by the gluteal muscles. In addition, it increases the activation of the gluteal muscles over the tensor fascia latae.

Alternative technique: elastic band around ankles

Alternative technique: elastic band around feet



Ensure the patient does not drift into internal rotation of the lower limb as this increases the activity of the erector spinae and may result in excessive hip adduction and internal rotation.^{129,130} Externally rotating the lower limbs can be used as a technique as the same level of gluteus medius activity is achieved in this position.¹²⁹ If the patient continues to struggle with technique you may need to prescribe the exercise with a lighter elastic band.

If you wish to favour one side it is important to note that electromyographic activity of the stance limb is higher than the moving limb.¹²⁹ This is because “the gluteus medius muscle must overcome band resistance in addition to the contralateral pelvic drop on the moving non-weight-bearing limb”

Key teaching phrases

- “Keep your knees in line with your second toe.”
- “Ensure your knees don’t move inwards.”

Contraindications

- Excessive pain in the knees, hips or ankles
- Limitations in hip, knee or ankle range of motion
- Posterior cruciate ligament injuries (healing needs to be well advanced)¹³¹

Evidence

Varying levels of gluteal activity has been observed in studies and presumably varying tensions and positioning of the elastic band could account for this.^{99,105,132} Youdis et al. found levels of greater than 50% of maximum voluntary contraction for gluteus medius and therefore monster walks could be effectively used for strengthening.¹²⁹ The activity of gluteus maximus was less than 40%.¹²⁹

In their study Begalle et al. found monster walks to have lower quadriceps-to-hamstrings coactivation ratios compared with lunge-type exercises suggesting it is a beneficial exercise for knee injury prevention and anterior cruciate ligament rehabilitation.¹³³

Neurotip

It has been evident for some time that the hip adductors and abductors play an important role in balance control in the medial lateral direction during standing and walking.¹³⁴ The ankle inverters and evertors only play a very small role (as opposed to anterior posterior balance control where the ankle flexors and extensors have a dominant role). This is because it is the muscles around the pelvis that control foot placement with some contribution from the plantar flexors and foot placement is critical to balance control.¹³⁵

What to say to patients

“This exercise strengthens the gluteus medius muscles on the side of your hip. They can help improve and prevent any inward movement of your knees. This could be beneficial to prevent injuries during jumping and landing activities.”

REFERENCES

1. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther*. 2013;43(2):54-64. doi:10.2519/jospt.2013.4116
2. Ackerman IN, Johansson MS, Grønne DT, et al. Are outcomes from an exercise therapy and patient education program for osteoarthritis associated with hip and knee replacement within two years? A register-based study of 9,339 patients with osteoarthritis. *Arthritis Care Res (Hoboken)*. Published online January 25, 2024. doi:10.1002/ACR.25303
3. Heald L, Dupree J, O'Brien M. The Correlation Between Gluteus Medius Strength and Lower Back Pain in Middle Age Adults: A Critically Appraised Topic. Published online February 18, 2022. Accessed February 16, 2024. <https://scholars.okstate.edu/en/publications/the-correlation-between-gluteus-medius-strength-and-lower-back-pa>
4. Bass CJ, Connell D a. Sonographic findings of tensor fascia lata tendinopathy: another cause of anterior groin pain. *Skeletal Radiol*. 2002;31(3):143-148. doi:10.1007/s00256-001-0459-7
5. Manojlović D, Kozinc Ž, Šarabon N. Trunk, Hip and Knee Exercise Programs for Pain Relief, Functional Performance and Muscle Strength in Patellofemoral Pain: Systematic Review and Meta-Analysis. *J Pain Res*. 2021;14:1431-1449. doi:10.2147/JPR.S301448
6. Na Y, Han C, Shi Y, Zhu Y, Ren Y, Liu W. Is Isolated Hip Strengthening or Traditional Knee-Based Strengthening More Effective in Patients With Patellofemoral Pain Syndrome? A Systematic Review With Meta-analysis. *Orthop J Sports Med*. 2021;9(7). doi:10.1177/23259671211017503
7. Friede MC, Innerhofer G, Fink C, Alegre LM, Csapo R. Conservative treatment of iliotibial band syndrome in runners: Are we targeting the right goals? *Physical Therapy in Sport*. 2022;54:44-52. doi:10.1016/J.PTSP.2021.12.006
8. Webster KA, Gribble PA. A comparison of electromyography of gluteus medius and maximus in subjects with and without chronic ankle instability during two functional exercises. *Phys Ther Sport*. 2013;14(1):17-22. doi:10.1016/j.ptsp.2012.02.002
9. Selkowitz DM, Beneck GJ, Powers CM. Comparison of electromyographic activity of the gluteal muscles and tensor fascia lata in persons with patellofemoral pain: evaluation of selected, hip-targeted exercises using indwelling fine-wire electrodes. *Physiother Theory Pract*. Published online 2023. doi:10.1080/09593985.2023.2215389
10. Kwak SD, Ahmad CS, Gardner TR, et al. Hamstrings and iliotibial band forces affect knee kinematics and contact pattern. *Journal of Orthopaedic Research*. 2000;18(1):101-108. doi:10.1002/jor.1100180115
11. Merican a M, Amis a a. Anatomy of the lateral retinaculum of the knee. *J Bone Joint Surg Br*. 2008;90(4):527-534. doi:10.1302/0301-620X.90B4.20085
12. Merican AM, Amis A a. Iliotibial band tension affects patellofemoral and tibiofemoral kinematics. *J Biomech*. 2009;42(10):1539-1546. doi:10.1016/j.jbiomech.2009.03.041
13. Sutter R, Kalberer F, Binkert CA, Graf N, Pfirrmann CWA, Gutzzeit A. Abductor tendon tears are associated with hypertrophy of the tensor fasciae latae muscle. *Skeletal Radiol*. 2013;42(5):627-633. doi:10.1007/s00256-012-1514-2
14. Kingzett-Taylor A, Tirman PFJ, Feller J, et al. Tendinosis and tears of gluteus medius and minimus muscles as a cause of hip pain: MR imaging findings. *American Journal of Roentgenology*. 1999;173(4):1123-1126. doi:10.2214/ajr.173.4.10511191
15. Meghpara MB, Bheem R, Shah S, et al. Prevalence of Gluteus Medius Pathology on Magnetic Resonance Imaging in Patients Undergoing Hip Arthroscopy for Femoroacetabular Impingement: Asymptomatic Tears Are Rare, Whereas Tendinosis Is Common. *Am J Sports Med*. 2020;48(12):2933-2938. doi:10.1177/0363546520952766
16. Grimaldi A, Richardson C, Durbridge G, Donnelly W, Darnell R, Hides J. The association between degenerative hip joint pathology and size of the gluteus maximus and tensor fascia lata muscles. *Man Ther*. 2009;14(6):611-617. doi:10.1016/j.math.2008.11.002
17. Dwyer MK, Stafford K, Mattacola CG, Uhl TL, Giordani M. Comparison of gluteus medius muscle activity during functional tasks in individuals with and without osteoarthritis of the hip joint. *Clinical Biomechanics*. 2013;28(7):757-761. doi:10.1016/j.clinbiomech.2013.07.007
18. Rostron ZPJ, Zacharias A, Semciw AI, et al. Effects of a targeted resistance intervention compared to a sham intervention on gluteal muscle hypertrophy, fatty infiltration and strength in people with hip osteoarthritis: analysis of secondary outcomes from a randomised clinical trial. *BMC Musculoskelet Disord*. 2022;23(1). doi:10.1186/s12891-022-05907-4
19. Grimaldi A, Richardson C, Stanton W, Durbridge G, Donnelly W, Hides J. The association between degenerative hip joint pathology and size of the gluteus medius, gluteus minimus and piriformis muscles. *Man Ther*. 2009;14(6):605-610. doi:10.1016/j.math.2009.07.004
20. Cooper NA, Scavo KM, Strickland KJ, et al. Prevalence of gluteus medius weakness in people with chronic low back pain compared to healthy controls. *Eur Spine J*. Published online May 26, 2015. doi:10.1007/s00586-015-4027-6

21. Simons DG, Travell JG. Myofascial origins of low back pain. 3. Pelvic and lower extremity muscles. *Postgrad Med.* 1983;73(2):99-105, 108.
22. Njoo KH, Van der Does E. The occurrence and inter-rater reliability of myofascial trigger points in the quadratus lumborum and gluteus medius: a prospective study in non-specific low back pain patients and controls in general practice. *Pain.* 1994;58(3):317-323.
23. Bewyer KJ, Bewyer DC, Messenger D, Kennedy CM. Pilot data: association between gluteus medius weakness and low back pain during pregnancy. *Iowa Orthopaedic Journal.* 2009;29:97-99.
24. Cooper NA, Scavo KM, Strickland KJ, et al. Prevalence of gluteus medius weakness in people with chronic low back pain compared to healthy controls. *European Spine Journal.* Published online 2015:1-8. doi:10.1007/s00586-015-4027-6
25. Embaby E, Abdallah A. Trunk and Gluteus-Medius Muscles' Fatigability During Occupational Standing in Clinical Instructors with Low Back Pain. *Br J Sports Med.* 2013;47(10):e3-e3. doi:10.1136/bjsports-2013-092558.85
26. Kankaanpää M, Taimela S, Laaksonen D, Hänninen O, Airaksinen O. Back and hip extensor fatigability in chronic low back pain patients and controls. *Arch Phys Med Rehabil.* 1998;79(4):412-417. doi:S0003-9993(98)90142-3 [pii]
27. Leinonen V, Kankaanpää M, Airaksinen O, Hänninen O. Back and hip extensor activities during trunk flexion/extension: Effects of low back pain and rehabilitation. *Arch Phys Med Rehabil.* 2000;81(1):32-37. doi:10.1016/S0003-9993(00)90218-1
28. Lee SW, Kim SY, L S, K SY. Effects of hip exercises for chronic low-back pain patients with lumbar instability. *J Phys Ther Sci.* 2015;27(2):345-348. doi:10.1589/jpts.27.345
29. Vleeming A, Stoeckart R, Volkers AC, Snijders CJ. Relation between form and function in the sacroiliac joint. Part I: Clinical anatomical aspects. *Spine (Phila Pa 1976).* 1990;15(2):130-132.
30. Van Wingerden JP, Vleeming A, Buyruk HM, Raissadat K. Stabilization of the sacroiliac joint in vivo: verification of muscular contribution to force closure of the pelvis. *Eur Spine J.* 2004;13(3):199-205. doi:10.1007/s00586-003-0575-2
31. Vleeming A, Volkers, Snijders CJ, Stoeckart R. Relation between form and function in the sacroiliac joint. Part II: Biomechanical aspects. *Spine (Phila Pa 1976).* 1990;15(2):133-136.
32. Snijders CJ, Vleeming A, Stoeckart R. Transfer of lumbosacral load to iliac bones and legs Part 1: Biomechanics of self-bracing of the sacroiliac joints and its significance for treatment and exercise. *Clin Biomech (Bristol, Avon).* 1993;8(6):285-294. doi:10.1016/0268-0033(93)90002-Y
33. Vleeming a., Van Wingerden JP, Snijders CJ, Stoeckart R, Stijnen T. Load application to the sacrotuberous ligament; influences on sacroiliac joint mechanics. *Clinical Biomechanics.* 1989;4(4):204-209. doi:10.1016/0268-0033(89)90003-X
34. Vleeming A, Stoeckart R, Snijders CJ. The sacrotuberous ligament: a conceptual approach to its dynamic role in stabilizing the sacroiliac joint. *Clinical Biomechanics.* 1989;4(4):201-203. doi:10.1016/0268-0033(89)90002-8
35. Van Wingerden JP, Vleeming A, Snijders CJ, Stoeckart R, Wingerden J, Vleeming A. A functional-anatomical approach to the spine-pelvis mechanism: interaction between the biceps femoris muscle and the sacrotuberous ligament. *Eur Spine J.* 1993;2(3):140-144. doi:10.1007/BF00301411
36. Mooney V, Pozos R, Vleeming a, Gulick J, Swenski D. Exercise treatment for sacroiliac pain. *Orthopedics.* 2001;24(1):29-32.
37. Vleeming A, Pool-Goudzwaard AL, Hammudoghlu D, Stoeckart R, Snijders CJ, Mens JM. The function of the long dorsal sacroiliac ligament: its implication for understanding low back pain. *Spine (Phila Pa 1976).* 1996;21(5):556-562. doi:10.1097/00007632-199603010-00005
38. Fortin JD, Aprill CN, Ponthieux B, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part II: Clinical evaluation. *Spine (Phila Pa 1976).* 1994;19(13):1483-1489. doi:PMID: 7939978
39. Fortin JD, Dwyer AP, West S, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/arthrography technique. Part I: Asymptomatic volunteers. *Spine (Phila Pa 1976).* 1994;19(13):1475-1482.
40. Fortin JD, Kissling RO, O'Connor BL, Vilensky JA. Sacroiliac joint innervation and pain. *Am J Orthop (Belle Mead NJ).* 1999;28(12):687-690.
41. Schwarzer a C, Aprill CN, Bogduk N. The sacroiliac joint in chronic low back pain. *Spine (Phila Pa 1976).* 1995;20(1):31-37.
42. Vleeming A, de Vries HJ, Mens JM a, van Wingerden JP. Possible role of the long dorsal sacroiliac ligament in women with peripartum pelvic pain. *Acta Obstet Gynecol Scand.* 2002;81(5):430-436. doi:10.1034/j.1600-0412.2002.810510.x
43. Ostgaard HC, Zetherstrom G, Roos-Hansson E. The posterior pelvic pain provocation test in pregnant women. *European Spine Journal.* 1994;3(5):258-260. doi:10.1111/j.1442-2018.2008.00406.x
44. Mens JM a, Vleeming A, Snijders CJ, Stam HJ, Ginai AZ. The active straight leg raising test and mobility of the pelvic joints. *European Spine Journal.* 1999;8(6):468-474. doi:10.1007/s005860050206

45. Jan M. A. Mens, Andry Vleeming CJS, Bart W. Koes, and Henk J. Stam M. Reliability and Validity of the Active Straight Leg Raise Test in Posterior Pelvic Pain Since Pregnancy. *Spine (Phila Pa 1976)*. 2001;26(10):1167-1171. doi:10.1097/00007632-200201150-00015
46. Mens JM a, Vleeming A, Snijders CJ, Koes BW, Stam HJ. Validity of the active straight leg raise test for measuring disease severity in patients with posterior pelvic pain after pregnancy. *Spine (Phila Pa 1976)*. 2002;27(2):196-200. doi:10.1097/00007632-200201150-00015
47. Hungerford B, Gilleard W, Hodges P. Evidence of altered lumbopelvic muscle recruitment in the presence of sacroiliac joint pain. *Spine (Phila Pa 1976)*. 2003;28(14):1593-1600. doi:10.1097/01.BRS.0000076821.41875.1C
48. Reiman MP, Bolgla L a, Loudon JK. A literature review of studies evaluating gluteus maximus and gluteus medius activation during rehabilitation exercises. *Physiother Theory Pract*. 2012;28(4):257-268. doi:10.3109/09593985.2011.604981
49. O'Sullivan K, Smith SM, Sainsbury D. Electromyographic analysis of the three subdivisions of gluteus medius during weight-bearing exercises. *Sports Med Arthrosc Rehabil Ther Technol*. 2010;2:17. doi:10.1186/1758-2555-2-17
50. Robinson RL, Nee RJ. Analysis of hip strength in females seeking physical therapy treatment for unilateral patellofemoral pain syndrome. *J Orthop Sports Phys Ther*. 2007;37(5):232-238. doi:10.2519/jospt.2007.2439
51. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther*. 2010;40(2):42-51. doi:10.2519/jospt.2010.3337
52. Powers CM, Ward SR, Fredericson M, Guillet M, Shellock FG. Patellofemoral kinematics during weight-bearing and non-weight-bearing knee extension in persons with lateral subluxation of the patella: a preliminary study. *J Orthop Sports Phys Ther*. 2003;33(11):677-685. doi:10.2519/jospt.2003.33.11.677
53. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther*. 2009;39(1):12-19. doi:10.2519/jospt.2009.2885
54. Reiman MP, Bolgla L a, Lorenz D. Hip functions influence on knee dysfunction: a proximal link to a distal problem. *J Sport Rehabil*. 2009;18(1):33-46.
55. Mascal CL, Landel R, Powers C. Management of patellofemoral pain targeting hip, pelvis, and trunk muscle function: 2 case reports. *J Orthop Sports Phys Ther*. 2003;33(11):647-660. doi:10.2519/jospt.2003.33.11.647
56. Selkowitz DM, Beneck GJ, Powers CM. Persons with patellofemoral pain exhibit altered hip abductor muscle recruitment while performing hip abductor exercises. *Physiother Theory Pract*. Published online 2022. doi:10.1080/09593985.2022.2101167
57. Bolgla L a, Malone TR, Umberger BR, Uhl TL. Hip strength and hip and knee kinematics during stair descent in females with and without patellofemoral pain syndrome. *J Orthop Sports Phys Ther*. 2008;38(1):12-18. doi:10.2519/jospt.2008.2462
58. Dolak KL, Silkman C, Medina McKeon J, Hosey RG, Lattermann C, Uhl TL. Hip strengthening prior to functional exercises reduces pain sooner than quadriceps strengthening in females with patellofemoral pain syndrome: a randomized clinical trial. *J Orthop Sports Phys Ther*. 2011;41(8):560-570. doi:10.2519/jospt.2011.3499
59. Nakagawa TH, Muniz TB, Baldon RDM, Dias Maciel C, de Menezes Reiff RB, Serrão FV. The effect of additional strengthening of hip abductor and lateral rotator muscles in patellofemoral pain syndrome: a randomized controlled pilot study. *Clin Rehabil*. 2008;22(12):1051-1060. doi:10.1177/0269215508095357
60. Ismail MM, Gamaleldein MH, Hassa KA. Closed Kinetic Chain exercises with or without additional hip strengthening exercises in management of Patellofemoral pain syndrome: A randomized controlled trial. *Eur J Phys Rehabil Med*. 2013;49(5):687-698. doi:R33Y9999N00A0249 [pii]
61. Fukuda TY. Hip Posterolateral Musculature Strengthening in Sedentary Women With Patellofemoral Pain Syndrome: A Randomized Controlled Clinical Trial With 1-Year Follow-up. *Journal of Orthopaedic and Sports Physical Therapy*. 2012;42(10):823-830. doi:10.2519/jospt.2012.4184
62. Ford KR, Nguyen AD, Dischiavi SL, Hegedus EJ, Zuk EF, Taylor JB. An evidence-based review of hip-focused neuromuscular exercise interventions to address dynamic lower extremity valgus. *Open Access J Sports Med*. 2015;6:291-303. doi:10.2147/OAJSM.S72432
63. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*. 2005;33(4):492-501. doi:10.1177/0363546504269591
64. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: Part 1, mechanisms and risk factors. *Am J Sports Med*. 2006;34(2):299-311. doi:10.1177/0363546505284183
65. Myer GD, Ford KR, Di Stasi SL, Foss KDB, Micheli LJ, Hewett TE. High knee abduction moments are common risk factors for patellofemoral pain (PFP) and anterior cruciate ligament (ACL) injury in girls: is PFP itself a predictor for subsequent ACL injury? *Br J Sports Med*. 2015;49(2):118-122.

doi:10.1136/bjsports-2013-092536

66. Ford KR, Myer GD, Hewett TE. Valgus knee motion during landing in high school female and male basketball players. *Med Sci Sports Exerc.* 2003;35(10):1745-1750. doi:10.1249/01.MSS.0000089346.85744.D9

67. Ford KR, Myer GD, Toms HE, Hewett TE. Gender differences in the kinematics of unanticipated cutting in young athletes. *Med Sci Sports Exerc.* 2005;37(1):124-129. doi:10.1249/01.MSS.0000150087.95953.C3

68. Ford KR, Myer GD, Smith RL, Vianello RM, Seiwert SL, Hewett TE. A comparison of dynamic coronal plane excursion between matched male and female athletes when performing single leg landings. *Clinical Biomechanics.* 2006;21(1):33-40. doi:10.1016/j.clinbiomech.2005.08.010

69. Ford KR, Shapiro R, Myer GD, Van Den Bogert AJ, Hewett TE. Longitudinal sex differences during landing in knee abduction in young athletes. *Med Sci Sports Exerc.* 2010;42(10):1923-1931. doi:10.1249/MSS.0b013e3181dc99b1

70. Malinzak RA, Colby SM, Kirkendall DT, Yu B, Garrett WE. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clinical Biomechanics.* 2001;16(5):438-445. doi:10.1016/S0268-0033(01)00019-5

71. Hewett TE, Myer GD, Ford KR. Decrease in neuromuscular control about the knee with maturation in female athletes. *J Bone Joint Surg Am.* 2004;86-A(8):1601-1608. doi:86/8/1601 [pii]

72. Chappell JD, Yu B, Kirkendall DT, Garrett WE. A comparison of knee kinetics between male and female recreational athletes in stop-jump tasks. *Am J Sports Med.* 2002;30(2):261-267.

73. McLean SG, Huang X, Su A, Van Den Bogert AJ. Sagittal plane biomechanics cannot injure the ACL during side-step cutting. *Clin Biomech (Bristol, Avon).* 2004;19(8):828-838. doi:10.1016/j.clinbiomech.2004.06.006

74. Kernozek TW, Torry MR, Van Hoof H, Cowley H, Tanner S. Gender differences in frontal and sagittal plane biomechanics during drop landings. *Med Sci Sports Exerc.* 2005;37(6):1003-1012. doi:10.1249/01.mss.0000171616.14640.2b

75. Zeller BL, McCrory JL, Kibler W Ben, Uhl TL. Differences in kinematics and electromyographic activity between men and women during the single-legged squat. *Am J Sports Med.* 31(3):449-456.

76. Pappas E, Hagins M, Sheikhzadeh A, Nordin M, Rose D. Biomechanical differences between unilateral and bilateral landings from a jump: gender differences. *Clin J Sport Med.* 2007;17(4):263-268. doi:10.1097/JSM.0b013e31811f415b

77. Hewett TE, Ford KR, Myer GD, Wanstrath K, Schep-er M. Gender differences in hip adduction motion and torque during a single-leg agility maneuver. *J Orthop Res.*

2006;24(3):416-421. doi:10.1002/jor.20056

78. Carson DW, Ford KR. Sex Differences in Knee Abduction During Landing: A Systematic Review. *Sports Health: A Multidisciplinary Approach.* 2011;3(4):373-382. doi:10.1177/1941738111410180

79. Kline PW, Burnham J, Yonz M, Johnson D, Ireland ML, Noehren B. Hip external rotation strength predicts hop performance after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(4):1137-1144. doi:10.1007/S00167-017-4534-6

80. Noehren B, Davis I, Hamill J. ASB Clinical Biomechanics Award Winner 2006. Prospective study of the biomechanical factors associated with iliotibial band syndrome. *Clinical Biomechanics.* 2007;22(9):951-956. doi:10.1016/j.clinbiomech.2007.07.001

81. Earl JE, Hertel JN, Denegar CR. Patterns of Dynamic Malalignment, Muscle Activation, Joint Motion, and Patellofemoral-Pain Syndrome. *J Sport Rehabil.* 2005;14:216-234.

82. Ferber R, Noehren B, Hamill J, Davis IS. Competitive female runners with a history of iliotibial band syndrome demonstrate atypical hip and knee kinematics. *J Orthop Sports Phys Ther.* 2010;40(2):52-58. doi:10.2519/jospt.2010.3028

83. Fredericson M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreich N, Sahrmann SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med.* 2000;10:169-175. doi:10.1097/00042752-200007000-00004

84. Fredericson M, Weir A. Practical management of iliotibial band friction syndrome in runners. *Clin J Sport Med.* 2006;16(3):261-268. doi:10.1097/00042752-200605000-00013

85. Fredericson M, Wolf C. Iliotibial band syndrome in runners: Innovations in treatment. *Sports Medicine.* 2005;35:451-459. doi:10.2165/00007256-200535050-00006

86. Beckman SM, Buchanan TS. Ankle inversion injury and hypermobility: Effect on hip and ankle muscle electromyography onset latency. *Arch Phys Med Rehabil.* 1995;76(12):1138-1143. doi:10.1016/S0003-9993(95)80123-5

87. Presswood L, Cronin J, Keogh JWL, Whatman C. Gluteus Medius: Applied Anatomy, Dysfunction, Assessment, and Progressive Strengthening. *Strength Cond J.* 2008;30:41-53. doi:10.1519/SSC.0b013e318187f19a

88. Kawaguchi K, Taketomi S, Mizutani Y, et al. Hip Abductor Muscle Strength Deficit as a Risk Factor for Inversion Ankle Sprain in Male College Soccer Players: A Prospective Cohort Study. *Orthop J Sports Med.* 2021;9(7).

doi:10.1177/23259671211020287

89. Lee D. Instability of the Sacroiliac Joint and the Consequences to Gait. *Journal of Manual & Manipulative Therapy*. 1996;4(1):22-29. doi:10.1179/jmt.1996.4.1.22

90. Presswood L, Cronin J, Keogh JWL, Whatman C. Gluteus Medius: Applied Anatomy, Dysfunction, Assessment, and Progressive Strengthening. *Strength Cond J*. 2008;30(5):41-53. doi:10.1519/SSC.0b013e318187f19a

91. Blazeovich AJ. Optimizing Hip Musculature For Greater Sprint Running Speed. *Strength Cond J*. 2000;22(2):22.

92. Schache AG, Blanch PD, Dorn TW, Brown N a T, Rosemond D, Pandy MG. Effect of running speed on lower limb joint kinetics. *Med Sci Sports Exerc*. 2011;43(7):1260-1271. doi:10.1249/MSS.0b013e3182084929

93. Secomb JL, Dascombe BJ, Nimphius S. Importance of joint angle-specific Hip strength for skating performance in semiprofessional ice hockey athletes. *J Strength Cond Res*. 2021;35(9):2599-2603. doi:10.1519/JSC.0000000000004087

94. Philippon MJ, Decker MJ, Giphart JE, Torry MR, Wahoff MS, LaPrade RF. Rehabilitation exercise progression for the gluteus medius muscle with consideration for iliopsoas tendinitis: an in vivo electromyography study. *Am J Sports Med*. 2011;39(8):1777-1785. doi:10.1177/0363546511406848

95. Robertson DGE, Wilson JMJ, St. Pierre T a. Lower extremity muscle functions during full squats. *J Appl Biomech*. 2008;24(4):333-339.

96. Hollman JH, Ginos BE, Kozuchowski J, Vaughn AS, Krause D a, Youdas JW. Relationships between knee valgus, hip-muscle strength, and hip-muscle recruitment during a single-limb step-down. *J Sport Rehabil*. 2009;18(1):104-117.

97. Hamstra-Wright KL, Huxel Bliven K. Effective exercises for targeting the gluteus medius. *J Sport Rehabil*. 2012;21(3):296-300.

98. Simenz CJ, Garceau LR, Lutsch BN, Suchomel TJ, Ebben WP. Electromyographical analysis of lower extremity muscle activation during variations of the loaded step-up exercise. *J Strength Cond Res*. 2012;26(12):3398-3405. doi:10.1519/JSC.0b013e3182472fad

99. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimizing activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther*. 2013;43:54-64. doi:10.2519/jospt.2013.4116

100. McBeth JM, Earl-Boehm JE, Cobb SC, Huddleston WE. Hip muscle activity during 3 side-lying hip-strengthening exercises in distance runners. *J Athl Train*. 47(1):15-23.

101. Sidorkewicz N, Cambridge EDJ, McGill SM. Examining the effects of altering hip orientation on gluteus medius

and tensor fasciae latae interplay during common non-weight-bearing hip rehabilitation exercises. *Clin Biomech (Bristol, Avon)*. 2014;29(9):971-976. doi:10.1016/j.clinbiomech.2014.09.002

102. Dostal WF, Soderberg GL, Andrews JG. Actions of hip muscles. *Phys Ther*. 1986;66:351-361. doi:10.1097/01241398-198703000-00046

103. Delp SL, Hess WE, Hungerford DS, Jones LC. Variation of rotation moment arms with hip flexion. *J Biomech*. 1999;32:493-501. doi:10.1016/S0021-9290(99)00032-9

104. Retchford T, Crossley KM, Grimaldi A, Kemp JL, Cowan SM. Can local muscles augment stability in the hip? A narrative literature review. *Journal of Musculoskeletal Neuronal Interactions*. 2013;13:1-12. doi:10.1136/bjism.2009.066944

105. Distefano LJ, Blackburn JT, Marshall SW, Padua DA. Gluteal muscle activation during common therapeutic exercises. *J Orthop Sports Phys Ther*. 2009;39:532-540. doi:10.2519/jospt.2009.2796

106. Gupta a, Fernihough B, Bailey G, Bombeck P, Clarke a, Hopper D. An evaluation of differences in hip external rotation strength and range of motion between female dancers and non-dancers. *Br J Sports Med*. 2004;38(6):778-783. doi:10.1136/bjism.2003.010827

107. Bennell KL, Khan KM, Matthews BL, Singleton C. Changes in hip and ankle range of motion and hip muscle strength in 8–11 year old novice female ballet dancers and controls: a 12 month follow up study. *Br J Sports Med*. 2001;35:54-59. doi:10.1136/bjism.35.1.54

108. Hamilton WG, Hamilton LH, Marshall P, Molnar M. A profile of the musculoskeletal characteristics of elite professional ballet dancers. *Am J Sports Med*. 1992;20(3):267-273. doi:10.1177/036354659202000306

109. Judge LW, Moreau C, Burke JR. Neural adaptations with sport-specific resistance training in highly skilled athletes. *J Sports Sci*. 2003;21(5):419-427. doi:10.1080/0264041031000071173

110. Parr VA, Pettitt ML, Krzyzanowicz R, Krzyzanowicz S, Pryor RR. Type of Turnout and Injury Rates in Elite Dancers: A Critically Appraised Topic. *J Sport Rehabil*. 2022;31(6):815-818. doi:10.1123/JSR.2021-0243

111. Tobey K, Mike J. Single-leg glute bridge. *Strength Cond J*. 2018;40(2):110-114. doi:10.1519/SSC.0000000000000323

112. Yoon JO, Kang MH, Kim JS, Oh JS. Effect of modified bridge exercise on trunk muscle activity in healthy adults: a cross sectional study. *Braz J Phys Ther*. 2018;22(2):161. doi:10.1016/J.BJPT.2017.09.005

113. Boren K, Conrey C, Le Coguic J, Paprocki L, Voight M, Robinson TK. Electromyographic analysis of gluteus medius and gluteus maximus during rehabilitation exercises. *Int J*

Sports Phys Ther. 2011;6(3):206-223.

114. Ekstrom RA, Donatelli RA, Carp KC. Electromyographic analysis of core trunk, hip, and thigh muscles during 9 rehabilitation exercises. *J Orthop Sports Phys Ther.* 2007;37(12):754-762. doi:10.2519/jospt.2007.2471

115. Somerson JS, Isby IJ, Hagen MS, Kweon CY, Gee AO. The Menstrual Cycle May Affect Anterior Knee Laxity and the Rate of Anterior Cruciate Ligament Rupture: A Systematic Review and Meta-Analysis. *JBJS Rev.* 2019;7(9). doi:10.2106/JBJS.RVW.18.00198

116. Rozzi SL, Lephart SM, Gear WS, Fu FH. Knee joint laxity and neuromuscular characteristics of male and female soccer and basketball players. *Am J Sports Med.* 1999;27(3):312-319.

117. Lewis CL, Sahrmann SA. Muscle activation and movement patterns during prone hip extension exercise in women. *J Athl Train.* 2009;44(3):238-248. doi:10.4085/1062-6050-44.3.238

118. Anders M. Glutes to the max. *ACE FitnessMatters.* Published online 2006:7-9.

119. Bishop B, Greenstein J. EMG Activation of Gluteal Musculature During Exercises With and Without Resistance. *13th Annual TRAC meeting.* 2011;Sport and.

120. Lewis CL, Sahrmann SA, Moran DW. Anterior hip joint force increases with hip extension, decreased gluteal force, or decreased iliopsoas force. *J Biomech.* 2007;40(16):3725-3731. doi:10.1016/j.jbiomech.2007.06.024

121. Shindle MK, Ranawat AS, Kelly BT. Diagnosis and Management of Traumatic and Atraumatic Hip Instability in the Athletic Patient. *Clin Sports Med.* 2006;25(2):309-326. doi:10.1016/j.csm.2005.12.003

122. Janda V. On the concept of postural muscles and posture in man. *Aust J Physiother.* 1983;29(3):83-84. doi:10.1016/S0004-9514(14)60665-6

123. Murphy DR, Byfield D, McCarthy P, Humphreys K, Gregory AA, Rochon R. Interexaminer Reliability of the Hip Extension Test for Suspected Impaired Motor Control of the Lumbar Spine. *J Manipulative Physiol Ther.* 2006;29(5):374-377. doi:10.1016/j.jmpt.2006.04.012

124. Lehman GJ, Lennon D, Tresidder B, Rayfield B, Poschar M. Muscle recruitment patterns during the prone leg extension. *BMC Musculoskelet Disord.* 2004;5:3. doi:10.1186/1471-2474-5-3

125. Lehman GJ. Trunk and hip muscle recruitment patterns during the prone leg extension following a lateral ankle sprain: a prospective case study pre and post injury. *Chiropr Osteopat.* 2006;14:4. doi:10.1186/1746-1340-14-4

126. Prince F, Winter D, Stergiou P, Walt S. Anticipatory control of upper body balance during human locomotion. *Gait Posture.* 1994;2(1):19-25. doi:10.1016/0966-

6362(94)90013-2

127. Berry JW, Lee TS, Foley HD, Lewis CL. Resisted Side Stepping: The Effect of Posture on Hip Abductor Muscle Activation. *J Orthop Sports Phys Ther.* 2015;45(9):675-682. doi:10.2519/jospt.2015.5888

128. Cambridge EDJ, Sidorkewicz N, Ikeda DM, McGill SM. Progressive hip rehabilitation: The effects of resistance band placement on gluteal activation during two common exercises. *Clinical Biomechanics.* 2012;27(7):719-724. doi:10.1016/j.clinbiomech.2012.03.002

129. Youdas JW, Foley BM, Kruger BL, et al. Electromyographic analysis of trunk and hip muscles during resisted lateral band walking. *Physiother Theory Pract.* 2013;29(2):113-123. doi:10.3109/09593985.2012.70449

2

130. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther.* 2010;40:42-51. doi:10.2519/jospt.2010.3337

131. Toutoungi DE, Lu TW, Leardini A, Catani F, O'Connor JJ. Cruciate ligament forces in the human knee during rehabilitation exercises. *Clin Biomech (Bristol, Avon).* 2000;15(3):176-187.

132. Goller M, Quittmann OJ, Alt T. How to activate the glutes best? Peak muscle activity of acceleration-specific pre-activation and traditional strength training exercises. *Eur J Appl Physiol.* Published online January 27, 2024:1-13. doi:10.1007/S00421-023-05400-3/FIGURES/4

133. Begalle RL, DiStefano LJ, Blackburn T, Padua DA. Quadriceps and hamstrings coactivation during common therapeutic exercises. *J Athl Train.* 2012;47(4):396-405. doi:10.4085/1062-6050-47.4.01

134. Winter DA. Human balance and posture control during standing and walking. *Gait Posture.* 1995;3(4):193-214. doi:10.1016/0966-6362(96)82849-9

135. Roelker SA, Kautz SA, Neptune RR. Muscle contributions to mediolateral and anteroposterior foot placement during walking. *J Biomech.* 2019;95. doi:10.1016/j.jbiomech.2019.08.004